

71622 U.S. PTO
05/23/97

810./101 A

PATENT
ATTORNEY DOCKET NO. 07334/004002

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant : Andrew W. Shyjan
Serial No.:
Filed : May 23, 1997
Title : COMPOSITIONS AND METHODS FOR THE DIAGNOSIS,
PREVENTION AND TREATMENT OF TUMOR PROGRESSION

Art Unit:
Examiner:

Box Patent Application

Assistant Commissioner for Patents
Washington, DC 20231


REQUEST FOR FILING CONTINUATION/DIVISIONAL APPLICATION

This is a request for filing a divisional application under 37 CFR 1.60, of pending prior application Serial No. 08/623,679 filed on March 29, 1996 by Andrew W. Shyjan for COMPOSITIONS AND METHODS FOR THE DIAGNOSIS, PREVENTION AND TREATMENT OF TUMOR PROGRESSION, which is a continuation-in-part of pending prior application Serial No. 08/412,431 filed on March 29, 1995 by Andrew W. Shyjan.

1. Enclosed is a true copy of the above identified prior application, including the declaration as originally filed and a copy of the signed declaration filed June 24, 1996 to complete the prior application. The true copy of the prior application is as follows: 161 pages of specification; five (5) page of claims; one (1) page of abstract; thirteen (13) sheets of drawings; and four (2) pages of declaration and power of attorney; Sequence Listing Statement under 37 CFR §1.821(f) and 29 pages of sequence listing.

"EXPRESS MAIL" Mailing Label Number EM518286979US

Date of Deposit May 23, 1997
I hereby certify under 37 CFR 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office To Addressee" with sufficient postage on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.



Lisa G. Gray

Applicants also include a copy of the sequence listing in paper and computer readable form (updated to include parent application data) and a new statement under 37 CFR §1.821(f). The new Sequence Listing contains no new matter.

2. Cancel in this application claims 1 through 5 and 7 through 13 of the prior application before calculating the filing fee. At least one original independent claim has been retained for the purpose of filing this application.

3. The filing fee, based on the original claims in the prior application less any claims cancelled herein, is calculated below:

CLAIMS AS FILED				
	Number Filed	Number Extra	Rate	Basic Fee
Total Claims	[14] - 20 =	[0] x	\$ 11 =	\$770
Independent Claims	[4] - 3 =	[1] x	\$ 40 =	[***]
Multiple Dependent Claims			\$130 =	[40]
				[***]
			Total Filing Fee =	\$810.00

4. Please amend the specification by replacing lines 5 and 6, on page 1, with the sentence: --This is a divisional of copending application Serial No. 08/632,679, filed March 29, 1996, which is a continuation-in-part of U.S. Serial No. 08/412,431, filed March 29, 1995.--

5. With respect to the prior application from which this application claims benefit under 35 U.S.C. 120, the inventor named in this application is the same as the one named in the prior application. The inventorship for the claims in this application is the same.

6. The prior application is assigned of record to Millennium Pharmaceuticals, Inc., a Delaware corporation, by virtue of an assignment submitted to the Patent and Trademark Office for recording on June 26, 1996 at Reel 8003, Frame 0981.

7. Enclosed is a check for the filing fee of \$810.00.

8. The Commissioner is hereby authorized to charge any additional fees which may be required by this paper, or credit any overpayment, to Deposit Account No. 06-1050. A duplicate of this request is enclosed.

9. The power of attorney in the prior application is to J. Peter Fasse, Reg. No. 32,983. The power appears in the true copy of the prior application. Address all future communications to J. Peter Fasse at Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804.

Respectfully submitted,

Date: 5-23-97


J. Peter Fasse
Reg. No. 32,983

Fish & Richardson P.C.
225 Franklin Street
Boston, MA 02110-2804

Telephone: 617/542-5070
Facsimile: 617/542-8906
241351.B11

PATENT
ATTORNEY DOCKET NO. 07334/004002

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant : Andrew W. Shyjan
Serial No.:
Filed : May 23, 1997
Title : COMPOSITIONS AND METHODS FOR THE DIAGNOSIS,
PREVENTION AND TREATMENT OF TUMOR PROGRESSION

Box Patent Application
Assistant Commissioner for Patents
Washington, DC 20231

VERIFIED STATEMENT UNDER 37 CFR § 1.821 (f)

I, Maureen Ruttle, declare that I personally prepared the paper and the computer-readable copies of the Sequence Listing filed herewith in the above-entitled case and that the content of both is the same.

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of The United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Date: May 23, 1997

Maureen Ruttle
Maureen Ruttle

Fish & Richardson P.C.
225 Franklin Street
Boston, MA 02110-2804
Telephone: 617/542-5070
Facsimile: 617/542-8906
245302.B11

"EXPRESS MAIL" Mailing Label Number EM51828697903

Date of Deposit May 23, 1997
I hereby certify under 37 CFR 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office To Addressee" with sufficient postage on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

Lisa G. Gray
Lisa G. Gray

468350 24429830

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Shyjan, Andrew W.
- (ii) TITLE OF INVENTION: COMPOSITIONS AND METHODS FOR THE
DIAGNOSIS, PREVENTION AND TREATMENT OF TUMOR
PROGRESSION
- (iii) NUMBER OF SEQUENCES: 9
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Fish & Richardson P.C.
 - (B) STREET: 225 Franklin Street
 - (C) CITY: Boston
 - (D) STATE: MA
 - (E) COUNTRY: USA
 - (F) ZIP: 02110-2804
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE: 23-MAY-1997
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/623,679
 - (B) FILING DATE: 29-MAR-1996
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/412,431
 - (B) FILING DATE: 29-MAR-1995
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Fasse, J. Peter
 - (B) REGISTRATION NUMBER: 32,983
 - (C) REFERENCE/DOCKET NUMBER: 07334/004002
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 617/542-5070
 - (B) TELEFAX: 617/542-8906
 - (C) TELEX: 200154

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 186 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GGTGCTGGAG TACCTCATGG GCGGTGCCTA CCGCTGCAAC TACTCTCGGA AAAGCTTCCG	60
GACTCTCTAC AACAACTTGT TTGGCCCTAA GAGGGTAGAG CTCAGCAGAC ACACAGTGTC	120

24429880

CTGTGCCTCC CAGAGTAACA TGTGGTTCCT TGATGTGCTT CCCCAAAAGC CCACCTGTGC 180
AGAATG 186

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2729 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

AAGGAGGCTA GGCTGCACCC TTCCCGCTTG CTCAGCAGCT GAGGCAGGGT CAGAAAGCAT 60
 GGATAGAGAA GACATTTTGC AAAAGGGAAT GCATCTTTGT AATTCCCAGT ACAAAGACC 120
 CTAACAGATG TTGCTGTGGT CAGCTCACTA ACCAGCACAT CCCCCCTTG CCGAGTGGGG 180
 CTCCCAGCAC AACAGGAGAG GACACCAAGC AGGCAGACAC GCAGTCCGGG AAATGGTCTG 240
 TCAGCAAACA CCCCCAGAGC TACCCAACAG ACTCCTATGG GATTCTTGAA TTCCAGGGTG 300
 GGGGTTACTC CAATAAAGCC ATG TAC ATC CGA GTC TCC TAC GAC ACC AAG 350
 Met Tyr Ile Arg Val Ser Tyr Asp Thr Lys
 1 5 10
 CCA GAT TCC CTG CTC CAC CTC ATG GTG AAG GAC TGG CAG CTG GAG CTC 398
 Pro Asp Ser Leu Leu His Leu Met Val Lys Asp Trp Gln Leu Glu Leu
 15 20 25
 CCG AAG CTC TTG ATA TCT GTG CAC GGA GGC CTC CAA AGC TTC GAG ATG 446
 Pro Lys Leu Leu Ile Ser Val His Gly Gly Leu Gln Ser Phe Glu Met
 30 35 40
 CAG TCC AAA CTG AAG CAG GTG TTT GGG AAA GGT CTG ATC AAG GCT GCC 494
 Gln Ser Lys Leu Lys Gln Val Phe Gly Lys Gly Leu Ile Lys Ala Ala
 45 50 55
 ATG ACC ACG GGG GCG TGG ATC TTC ACC GGG GGT GTG AGC ACT GGT GTC 542
 Met Thr Thr Gly Ala Trp Ile Phe Thr Gly Gly Val Ser Thr Gly Val
 60 65 70
 GTC AGC CAT GTG GGG GAT GCC TTG AAA GAC CAC TCC TCC AAG TCC AGA 590
 Val Ser His Val Gly Asp Ala Leu Lys Asp His Ser Ser Lys Ser Arg
 75 80 85 90
 GGC CGG CTC TGT GCT ATA GGA ATT GCT CCC TGG GGC ATG GTG GAG AAC 638
 Gly Arg Leu Cys Ala Ile Gly Ile Ala Pro Trp Gly Met Val Glu Asn
 95 100 105
 AAG GAA GAC CTG ATT GGA AAA GAT GTA ACA AGA GTC TAT CAG ACC ATG 686
 Lys Glu Asp Leu Ile Gly Lys Asp Val Thr Arg Val Tyr Gln Thr Met
 110 115 120
 TCC AAC CCT CTG AGC AAG CTC TCT GTG CTC AAC AAT TCC CAC ACT CAC 734
 Ser Asn Pro Leu Ser Lys Leu Ser Val Leu Asn Asn Ser His Thr His
 125 130 135
 TTC ATC TTG GCT GAC AAC GGC ACC CTG GGC AAG TAT GGT GCT GAG GTG 782
 Phe Ile Leu Ala Asp Asn Gly Thr Leu Gly Lys Tyr Gly Ala Glu Val
 140 145 150

46350 "24429880

AAG CTT CGA AGA CAG CTG GAA AAA CAC ATC TCC CTG CAG AAG ATC AAC Lys Leu Arg Arg Gln Leu Glu Lys His Ile Ser Leu Gln Lys Ile Asn 155 160 165 170	830
ACA AGG CTG GGC CAG GGT GTA CCT GTC GTG GGC CTA GTG GTA GAA GGT Thr Arg Leu Gly Gln Gly Val Pro Val Val Gly Leu Val Val Glu Gly 175 180 185	878
GGT CCT AAC GTG GTT TCT ATC GTC CTG GAG TAT CTC AAA GAA GAC CCT Gly Pro Asn Val Val Ser Ile Val Leu Glu Tyr Leu Lys Glu Asp Pro 190 195 200	926
CCT GTC CCT GTG GTG GTT TGC GAT GGC AGT GGA CGT GCC TCT GAC ATT Pro Val Pro Val Val Val Cys Asp Gly Ser Gly Arg Ala Ser Asp Ile 205 210 215	974
TTG TCC TTC GCA CAC AAA TAC TGC GAC GAA GGA GGA GTC ATA AAC GAG Leu Ser Phe Ala His Lys Tyr Cys Asp Glu Gly Gly Val Ile Asn Glu 220 225 230	1022
TCC CTG CGG GAC CAG CTT CTA GTT ACC ATT CAG AAA ACA TTT AAT TAC Ser Leu Arg Asp Gln Leu Val Thr Ile Gln Lys Thr Phe Asn Tyr 235 240 245 250	1070
AGC AAG TCC CAG TCG TAT CAG CTG TTT GCA ATT ATC ATG GAG TGC ATG Ser Lys Ser Gln Ser Tyr Gln Leu Phe Ala Ile Ile Met Glu Cys Met 255 260 265	1118
AAG AAG AAA GAA CTC GTC ACT GTG TTT CGG ATG GGT TCC GAG GGT CAG Lys Lys Lys Glu Leu Val Thr Val Phe Arg Met Gly Ser Glu Gly Gln 270 275 280	1166
CAA GAT GTC GAG ATG GCA ATT TTA ACT GCC TTG CTC AAA GGA ACC AAC Gln Asp Val Glu Met Ala Ile Leu Thr Ala Leu Leu Lys Gly Thr Asn 285 290 295	1214
GCA TCA GCT CCA GAT CAG CTG AGC TTG GCC CTG GCT TGG AAC CGG GTC Ala Ser Ala Pro Asp Gln Leu Ser Leu Ala Leu Ala Trp Asn Arg Val 300 305 310	1262
GAC ATA GCG CGA AGC CAG ATC TTC GTC TTT GGC CCA CAC TGG CCG CCA Asp Ile Ala Arg Ser Gln Ile Phe Val Phe Gly Pro His Trp Pro Pro 315 320 325 330	1310
CTG GGA AGC CTG GCC CCT CCT GTG GAC ACC AAA GCC GCA GAG AAG GAA Leu Gly Ser Leu Ala Pro Pro Val Asp Thr Lys Ala Ala Glu Lys Glu 335 340 345	1358
AAG AAG CCA CCC ACA GCC ACC ACC AAG GGG AGA GGA AAA GGA AAA GGC Lys Lys Pro Pro Thr Ala Thr Thr Lys Gly Arg Gly Lys Gly Lys Gly 350 355 360	1406
AAG AAG AAA GGC AAA GTG AAA GAG GAA GTG GAG GAA GAG ACG GAC CCC Lys Lys Lys Gly Lys Val Lys Glu Glu Val Glu Glu Glu Thr Asp Pro 365 370 375	1454
CGG AAG CTT GAG CTG CTC AAC TGG GTG AAT GCC CTG GAG CAA GCC ATG Arg Lys Leu Glu Leu Leu Asn Trp Val Asn Ala Leu Glu Gln Ala Met 380 385 390	1502
CTG GAT GCT CTT GTC CTA GAT CGG GTG GAC TTT GTA AAG CTC CTG ATT Leu Asp Ala Leu Val Leu Asp Arg Val Asp Phe Val Lys Leu Leu Ile 395 400 405 410	1550

GAA AAC GGA GTG AAC ATG CAG CAT TTC CTC ACC ATC CCG AGG CTG GAG Glu Asn Gly Val Asn Met Gln His Phe Leu Thr Ile Pro Arg Leu Glu 415 420 425	1598
GAG CTA TAC AAC ACC AGA CTG GGC CCA CCA AAC ACC CTT CAT CTG CTG Glu Leu Tyr Asn Thr Arg Leu Gly Pro Pro Asn Thr Leu His Leu Leu 430 435 440	1646
GTG CGG GAT GTA AAG AAG AGC AAC CTT CCA CCT GAT TAC CAC ATC AGC Val Arg Asp Val Lys Lys Ser Asn Leu Pro Pro Asp Tyr His Ile Ser 445 450 455	1694
CTC ATT GAT ATA GGA CTG GTG CTG GAG TAC CTC ATG GGC GGT GCC TAC Leu Ile Asp Ile Gly Leu Val Leu Glu Tyr Leu Met Gly Gly Ala Tyr 460 465 470	1742
CGC TGC AAC TAC ACT CGG AAA AGC TTC CGG ACT CTC TAC AAC AAC TTG Arg Cys Asn Tyr Thr Arg Lys Ser Phe Arg Thr Leu Tyr Asn Asn Leu 475 480 485 490	1790
TTT GGC CCT AAG AGG GTA GAG CTC AGC AGA CAC ACA GTG TCC TGT GCC Phe Gly Pro Lys Arg Val Glu Leu Ser Arg His Thr Val Ser Cys Ala 495 500 505	1838
TCC CAG AGT AAC ATG TGG TTC CTT GAT GTG CTT CCC CAA AAG CCC ACC Ser Gln Ser Asn Met Trp Phe Leu Asp Val Leu Pro Gln Lys Pro Thr 510 515 520	1886
TGT GCA GAA TGC AAC TCT TCA CCT CAC CTG TCC CAA ACT GAC ATC ACC Cys Ala Glu Cys Asn Ser Ser Pro His Leu Ser Gln Thr Asp Ile Thr 525 530 535	1934
CCA CCT CTG CCC T GACACCCAGT GCAGGGCCTC CTAGCTTTCA CATGCAGCCA Pro Pro Leu Pro 540	1987
TTCACATCGC CTCTCAAGAC TGGGCCAGGC AGTGCAACCT GTCAAGCATG TCTGTCCTCC	2047
CCTCCTTCCT ACAATAGCCC CCCCTCTGGG CCCCATGCCT CTGCTCTCTC AGCCCGTTCT	2107
CCTCCCCACT GATCACTGGC GCTCCTGTTG TCTTCCAAGG CAAGGAACAA GGAAAAGCAT	2167
CTTTTTGCCC AAAAAAGTTT AGGGCTCCCC GCTGTTCAAC CATAGCCAAC CTCACTGTAC	2227
ATCGGAGTCA TCCAGGCCAG CTGCCACACA CAAGCCTTCC CCACCCTATC CCAATAGACC	2287
CTATTCCTCC ATCAAAATCA AAGCTAACTC CTGGCCTGCC ACATTGCTTC TTCTTGCTCC	2347
AGCCTGTTAA ACCTCCAATA AATGTCAGAT CTGTGGGAAG CCTTCCTCAC TCTCACTCCA	2407
CAGTTTGTAC AGAGAGCGAG AGCCTCGTTT GGTTCCTACTT ACAAGGAAGG CTTTGTGTCT	2467
GTCTGTCCTT CCCAACTGAC TTCTGTTGAC AGAAGCAGTT TCCACATGAA AGCGTTGACT	2527
CACCTGGATG TTGTCATTAA TTAATAGTGA TACAAAATAT TGACACTTCT TTTCTGCTT	2587
CTTTGTTATG CAGCCGAAAG CACTTAAGCT TCTGGGAATG GAAGTAAGTA GGACATGTTT	2647
GTGGCAGTTT ATTTACTATA TATACCTTTG TCATTCTGTG GAAGCAAAAA TTGCAATGTT	2707
TTCCATGAAT AAAGCTCGTG CC	2729

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 542 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met	Tyr	Ile	Arg	Val	Ser	Tyr	Asp	Thr	Lys	Pro	Asp	Ser	Leu	Leu	His
1				5					10					15	
Leu	Met	Val	Lys	Asp	Trp	Gln	Leu	Glu	Leu	Pro	Lys	Leu	Leu	Ile	Ser
			20					25					30		
Val	His	Gly	Gly	Leu	Gln	Ser	Phe	Glu	Met	Gln	Ser	Lys	Leu	Lys	Gln
		35					40					45			
Val	Phe	Gly	Lys	Gly	Leu	Ile	Lys	Ala	Ala	Met	Thr	Thr	Gly	Ala	Trp
	50					55					60				
Ile	Phe	Thr	Gly	Gly	Val	Ser	Thr	Gly	Val	Val	Ser	His	Val	Gly	Asp
65					70					75					80
Ala	Leu	Lys	Asp	His	Ser	Ser	Lys	Ser	Arg	Gly	Arg	Leu	Cys	Ala	Ile
				85					90					95	
Gly	Ile	Ala	Pro	Trp	Gly	Met	Val	Glu	Asn	Lys	Glu	Asp	Leu	Ile	Gly
			100					105					110		
Lys	Asp	Val	Thr	Arg	Val	Tyr	Gln	Thr	Met	Ser	Asn	Pro	Leu	Ser	Lys
	115						120					125			
Leu	Ser	Val	Leu	Asn	Asn	Ser	His	Thr	His	Phe	Ile	Leu	Ala	Asp	Asn
	130					135					140				
Gly	Thr	Leu	Gly	Lys	Tyr	Gly	Ala	Glu	Val	Lys	Leu	Arg	Arg	Gln	Leu
145				150						155					160
Glu	Lys	His	Ile	Ser	Leu	Gln	Lys	Ile	Asn	Thr	Arg	Leu	Gly	Gln	Gly
				165					170					175	
Val	Pro	Val	Val	Gly	Leu	Val	Val	Glu	Gly	Gly	Pro	Asn	Val	Val	Ser
			180					185					190		
Ile	Val	Leu	Glu	Tyr	Leu	Lys	Glu	Asp	Pro	Pro	Val	Pro	Val	Val	Val
	195						200					205			
Cys	Asp	Gly	Ser	Gly	Arg	Ala	Ser	Asp	Ile	Leu	Ser	Phe	Ala	His	Lys
	210					215					220				
Tyr	Cys	Asp	Glu	Gly	Gly	Val	Ile	Asn	Glu	Ser	Leu	Arg	Asp	Gln	Leu
225					230					235					240
Leu	Val	Thr	Ile	Gln	Lys	Thr	Phe	Asn	Tyr	Ser	Lys	Ser	Gln	Ser	Tyr
				245					250					255	
Gln	Leu	Phe	Ala	Ile	Ile	Met	Glu	Cys	Met	Lys	Lys	Lys	Glu	Leu	Val
			260				265						270		
Thr	Val	Phe	Arg	Met	Gly	Ser	Glu	Gly	Gln	Gln	Asp	Val	Glu	Met	Ala
			275				280					285			

Ile Leu Thr Ala Leu Leu Lys Gly Thr Asn Ala Ser Ala Pro Asp Gln
 290 295 300
 Leu Ser Leu Ala Leu Ala Trp Asn Arg Val Asp Ile Ala Arg Ser Gln
 305 310 315 320
 Ile Phe Val Phe Gly Pro His Trp Pro Pro Leu Gly Ser Leu Ala Pro
 325 330 335
 Pro Val Asp Thr Lys Ala Ala Glu Lys Glu Lys Lys Pro Pro Thr Ala
 340 345 350
 Thr Thr Lys Gly Arg Gly Lys Gly Lys Gly Lys Lys Lys Gly Lys Val
 355 360 365
 Lys Glu Glu Val Glu Glu Glu Thr Asp Pro Arg Lys Leu Glu Leu Leu
 370 375 380
 Asn Trp Val Asn Ala Leu Glu Gln Ala Met Leu Asp Ala Leu Val Leu
 385 390 395 400
 Asp Arg Val Asp Phe Val Lys Leu Leu Ile Glu Asn Gly Val Asn Met
 405 410 415
 Gln His Phe Leu Thr Ile Pro Arg Leu Glu Glu Leu Tyr Asn Thr Arg
 420 425 430
 Leu Gly Pro Pro Asn Thr Leu His Leu Leu Val Arg Asp Val Lys Lys
 435 440 445
 Ser Asn Leu Pro Pro Asp Tyr His Ile Ser Leu Ile Asp Ile Gly Leu
 450 455 460
 Val Leu Glu Tyr Leu Met Gly Gly Ala Tyr Arg Cys Asn Tyr Thr Arg
 465 470 475 480
 Lys Ser Phe Arg Thr Leu Tyr Asn Asn Leu Phe Gly Pro Lys Arg Val
 485 490 495
 Glu Leu Ser Arg His Thr Val Ser Cys Ala Ser Gln Ser Asn Met Trp
 500 505 510
 Phe Leu Asp Val Leu Pro Gln Lys Pro Thr Cys Ala Glu Cys Asn Ser
 515 520 525
 Ser Pro His Leu Ser Gln Thr Asp Ile Thr Pro Pro Leu Pro
 530 535 540

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GGGGAAGCAC ATCAAGGAAC

20

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GCAACTACTA CACTCGGAAA AGC

23

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 4944 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 346..4837

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

ACTCATTATA GGGNTCGAGC GGCCGCCCCG GCAGGTTTGA GCTGTGCCCT CTCCATTCCA	60
CTGCTGTGGC AGGGTCAGAA ATCTTGGATA GAGAAAACCT TTTGCAAACG GGAATGTATC	120
TTTGTAATTC CTAGCACGAA AGACTCTAAC AGGTGTTGCT GTGGCCAGTT CACCAACCAG	180
CATATCCCCC CTCTGCCAAG TGCAACACCC AGCAAAAATG AAGAGGAAAG CAAACAGGTG	240
GAGACTCAGC CTGAGAAATG GTCTGTTGCC AAGCACACCC AGAGCTACCC AACAGATTCC	300
TATGGAGTTC TTGAATTCCA GGGTGCGGGA TATTCCAATA AAGCC ATG TAT ATC	354
Met Tyr Ile	
1	
CGT GTA TCC TAT GAC ACC AAG CCA GAC TCA CTG CTC CAT CTC ATG GTG	402
Arg Val Ser Tyr Asp Thr Lys Pro Asp Ser Leu Leu His Leu Met Val	
5 10 15	
AAA GAT TGG CAG CTG GAA CTC CCC AAG CTC TTA ATA TCT GTG CAT GGA	450
Lys Asp Trp Gln Leu Glu Leu Pro Lys Leu Leu Ile Ser Val His Gly	
20 25 30 35	
GGC CTC CAG AAC TTT GAG ATG CAG CCC AAG CTG AAA CAA GTC TTT GGG	498
Gly Leu Gln Asn Phe Glu Met Gln Pro Lys Leu Lys Gln Val Phe Gly	
40 45 50	
AAA GGC CTG ATC AAG GCT GCT ATG ACC ACC GGG GCC TGG ATC TTC ACC	546
Lys Gly Leu Ile Lys Ala Ala Met Thr Thr Gly Ala Trp Ile Phe Thr	
55 60 65	
GGG GGT GTC AGC ACA GGT GTT ATC AGC CAC GTA GGG GAT GCC TTG AAA	594
Gly Gly Val Ser Thr Gly Val Ile Ser His Val Gly Asp Ala Leu Lys	
70 75 80	
GAC CAC TCC TCC AAG TCC AGA GGC CGG GTT TGT GCT ATA GGA ATT GCT	642
Asp His Ser Ser Lys Ser Arg Gly Arg Val Cys Ala Ile Gly Ile Ala	
85 90 95	

268250 2442980

CCA TGG GGC ATC GTG GAG AAT AAG GAA GAC CTG GTT GGA AAG GAT GTA Pro Trp Gly Ile Val Glu Asn Lys Glu Asp Leu Val Gly Lys Asp Val 100 105 110 115	690
ACA AGA GTG TAC CAG ACC ATG TCC AAC CCT CTA AGT AAG CTC TCT GTG Thr Arg Val Tyr Gln Thr Met Ser Asn Pro Leu Ser Lys Leu Ser Val 120 125 130	738
CTC AAC AAC TCC CAC ACC CAC TTC ATC CTG GCT GAC AAT GGC ACC CTG Leu Asn Asn Ser His Thr His Phe Ile Leu Ala Asp Asn Gly Thr Leu 135 140 145	786
GGC AAG TAT GGC GCC GAG GTG AAG CTG CGA AGG CTG CTG GAA AAG CAC Gly Lys Tyr Gly Ala Glu Val Lys Leu Arg Arg Leu Leu Glu Lys His 150 155 160	834
ATC TCC CTC CAG AAG ATC AAC ACA AGA CTG GGG CAG GGC GTG CCC CTC Ile Ser Leu Gln Lys Ile Asn Thr Arg Leu Gly Gln Gly Val Pro Leu 165 170 175	882
GTG GGT CTC GTG GTG GAG GGG GGC CCT AAC GTG GTG TCC ATC GTC TTG Val Gly Leu Val Val Glu Gly Gly Pro Asn Val Val Ser Ile Val Leu 180 185 190 195	930
GAA TAC CTG CAA GAA GAG CCT CCC ATC CCT GTG GTG ATT TGT GAT GGC Glu Tyr Leu Gln Glu Glu Pro Pro Ile Pro Val Val Ile Cys Asp Gly 200 205 210	978
AGC GGA CGT GCC TCG GAC ATC CTG TCC TTT GCG CAC AAG TAC TGT GAA Ser Gly Arg Ala Ser Asp Ile Leu Ser Phe Ala His Lys Tyr Cys Glu 215 220 225	1026
GAA GGC GGA ATA ATA AAT GAG TCC CTC AGG GAG CAG CTT CTA GTT ACC Glu Gly Gly Ile Ile Asn Glu Ser Leu Arg Glu Gln Leu Leu Val Thr 230 235 240	1074
ATT CAG AAA ACA TTT AAT TAT AAT AAG GCA CAA TCA CAT CAG CTG TTT Ile Gln Lys Thr Phe Asn Tyr Asn Lys Ala Gln Ser His Gln Leu Phe 245 250 255	1122
GCA ATT ATA ATG GAG TGC ATG AAG AAG AAA GAA CTC GTC ACT GTG TTC Ala Ile Ile Met Glu Cys Met Lys Lys Lys Glu Leu Val Thr Val Phe 260 265 270 275	1170
AGA ATG GGT TCT GAG GGC CAG CAG GAC ATC GAG ATG GCA ATT TTA ACT Arg Met Gly Ser Glu Gly Gln Gln Asp Ile Glu Met Ala Ile Leu Thr 280 285 290	1218
GCC CTG CTG AAA GGA ACA AAC GTA TCT GCT CCA GAT CAG CTG AGC TTG Ala Leu Leu Lys Gly Thr Asn Val Ser Ala Pro Asp Gln Leu Ser Leu 295 300 305	1266
GCA CTG GCT TGG AAC CGC GTG GAC ATA GCA CGA AGC CAG ATC TTT GTC Ala Leu Ala Trp Asn Arg Val Asp Ile Ala Arg Ser Gln Ile Phe Val 310 315 320	1314
TTT GGG CCC CAC TGG ACG CCC CTG GGA AGC CTG GCA CCC CCG ACG GAC Phe Gly Pro His Trp Thr Pro Leu Gly Ser Leu Ala Pro Pro Thr Asp 325 330 335	1362
AGC AAA GCC ACG GAG AAG GAG AAG AAG CCA CCC ATG GCC ACC ACC AAG Ser Lys Ala Thr Glu Lys Glu Lys Lys Pro Pro Met Ala Thr Thr Lys 340 345 350 355	1410

GGA	GGA	AGA	GGA	AAA	GGG	AAA	GGC	AAG	AAG	AAA	GGG	AAA	GTG	AAA	GAG	1458
Gly	Gly	Arg	Gly	Lys	Gly	Lys	Gly	Lys	Lys	Lys	Gly	Lys	Val	Lys	Glu	
				360					365					370		
GAA	GTG	GAG	GAA	GAA	ACT	GAC	CCC	CGG	AAG	ATA	GAG	CTG	CTG	AAC	TGG	1506
Glu	Val	Glu	Glu	Glu	Thr	Asp	Pro	Arg	Lys	Ile	Glu	Leu	Leu	Asn	Trp	
			375					380					385			
GTG	AAT	GCT	TTG	GAG	CAA	GCG	ATG	CTA	GAT	GCT	TTA	GTG	TTA	GAT	CGT	1554
Val	Asn	Ala	Leu	Glu	Gln	Ala	Met	Leu	Asp	Ala	Leu	Val	Leu	Asp	Arg	
		390					395					400				
GTC	GAC	TTT	GTG	AAG	CTC	CTG	ATT	GAA	AAC	GGA	GTG	AAC	ATG	CAA	CAC	1602
Val	Asp	Phe	Val	Lys	Leu	Leu	Ile	Glu	Asn	Gly	Val	Asn	Met	Gln	His	
	405					410					415					
TTT	CTG	ACC	ATT	CCG	AGG	CTG	GAG	GAG	CTT	TAT	AAC	ACA	AGA	CTG	GGT	1650
Phe	Leu	Thr	Ile	Pro	Arg	Leu	Glu	Glu	Leu	Tyr	Asn	Thr	Arg	Leu	Gly	
420					425					430					435	
CCA	CCA	AAC	ACA	CTT	CAT	CTG	CTG	GTG	AGG	GAT	GTG	AAA	AAG	AGC	AAC	1698
Pro	Pro	Asn	Thr	Leu	His	Leu	Leu	Val	Arg	Asp	Val	Lys	Lys	Ser	Asn	
				440					445					450		
CTT	CCG	CCT	GAT	TAC	CAC	ATC	AGC	CTC	ATA	GAC	ATC	GGG	CTC	GTG	CTG	1746
Leu	Pro	Pro	Asp	Tyr	His	Ile	Ser	Leu	Ile	Asp	Ile	Gly	Leu	Val	Leu	
			455					460					465			
GAG	TAC	CTC	ATG	GGA	GGA	GCC	TAC	CGC	TGC	AAC	TAC	ACT	CGG	AAA	AAC	1794
Glu	Tyr	Leu	Met	Gly	Gly	Ala	Tyr	Arg	Cys	Asn	Tyr	Thr	Arg	Lys	Asn	
		470					475					480				
TTT	CGG	ACC	CTT	TAC	AAC	AAC	TTG	TTT	GGA	CCA	AAG	AGG	CCT	AAA	GCT	1842
Phe	Arg	Thr	Leu	Tyr	Asn	Asn	Leu	Phe	Gly	Pro	Lys	Arg	Pro	Lys	Ala	
	485					490					495					
CTT	AAA	CTT	CTG	GGA	ATG	GAA	GAT	GAT	GAG	CCT	CCA	GCT	AAA	GGG	AAG	1890
Leu	Lys	Leu	Leu	Gly	Met	Glu	Asp	Asp	Glu	Pro	Pro	Ala	Lys	Gly	Lys	
500					505					510				515		
AAA	AAA	AAA	AAA	AAG	AAA	AAG	GAG	GAA	GAG	ATC	GAC	ATT	GAT	GTG	GAC	1938
Lys	Lys	Lys	Lys	Lys	Lys	Lys	Glu	Glu	Glu	Ile	Asp	Ile	Asp	Val	Asp	
				520					525					530		
GAC	CCT	GCC	GTG	AGT	CGG	TTC	CAG	TAT	CCC	TTC	CAC	GAG	CTG	ATG	GTG	1986
Asp	Pro	Ala	Val	Ser	Arg	Phe	Gln	Tyr	Pro	Phe	His	Glu	Leu	Met	Val	
			535					540					545			
TGG	GCA	GTG	CTG	ATG	AAA	CGC	CAG	AAA	ATG	GCA	GTG	TTC	CTC	TGG	CAG	2034
Trp	Ala	Val	Leu	Met	Lys	Arg	Gln	Lys	Met	Ala	Val	Phe	Leu	Trp	Gln	
		550					555					560				
CGA	GGG	GAA	GAG	AGC	ATG	GCC	AAG	GCC	CTG	GTG	GCC	TGC	AAG	CTC	TAC	2082
Arg	Gly	Glu	Glu	Ser	Met	Ala	Lys	Ala	Leu	Val	Ala	Cys	Lys	Leu	Tyr	
	565					570					575					
AAG	GCC	ATG	GCC	CAC	GAG	TCC	TCC	GAG	AGT	GAT	CTG	GTG	GAT	GAC	ATC	2130
Lys	Ala	Met	Ala	His	Glu	Ser	Ser	Glu	Ser	Asp	Leu	Val	Asp	Asp	Ile	
580					585				590						595	
TCC	CAG	GAC	TTG	GAT	AAC	AAT	TCC	AAA	GAC	TTC	GGC	CAG	CTT	GCT	TTG	2178
Ser	Gln	Asp	Leu	Asp	Asn	Asn	Ser	Lys	Asp	Phe	Gly	Gln	Leu	Ala	Leu	
				600					605					610		

GAG TTA TTA GAC CAG TCC TAT AAG CAT GAC GAG CAG ATC GCT ATG AAA Glu Leu Leu Asp Gln Ser Tyr Lys His Asp Glu Gln Ile Ala Met Lys 615 620 625	2226
CTC CTG ACC TAC GAG CTG AAA AAC TGG AGC AAC TCG ACC TGC CTC AAA Leu Leu Thr Tyr Glu Leu Lys Asn Trp Ser Asn Ser Thr Cys Leu Lys 630 635 640	2274
CTG GCC GTG GCA GCC AAA CAC CGG GAC TTC ATT GCT CAC ACC TGC AGC Leu Ala Val Ala Ala Lys His Arg Asp Phe Ile Ala His Thr Cys Ser 645 650 655	2322
CAG ATG CTG CTG ACC GAT ATG TGG ATG GGA AGA CTG CGG ATG CGG AAG Gln Met Leu Leu Thr Asp Met Trp Met Gly Arg Leu Arg Met Arg Lys 660 665 670 675	2370
AAC CCC GGC CTG AAG GTT ATC ATG GGG ATT CTT CTA CCC CCC ACC ATC Asn Pro Gly Leu Lys Val Ile Met Gly Ile Leu Leu Pro Pro Thr Ile 680 685 690	2418
TTG TTT TTG GAA TTT CGC ACA TAT GAT GAT TTC TCG TAT CAA ACA TCC Leu Phe Leu Glu Phe Arg Thr Tyr Asp Asp Phe Ser Tyr Gln Thr Ser 695 700 705	2466
AAG GAA AAC GAG GAT GGC AAA GAA AAA GAA GAG GAA AAT ACG GAT GCA Lys Glu Asn Glu Asp Gly Lys Glu Lys Glu Glu Glu Asn Thr Asp Ala 710 715 720	2514
AAT GCA GAT GCT GGC TCA AGA AAG GGG GAT GAG GAG AAC GAG CAT AAA Asn Ala Asp Ala Gly Ser Arg Lys Gly Asp Glu Glu Asn Glu His Lys 725 730 735	2562
AAA CAG AGA ATT ATC CCC ATC GGA ACA AAA ATC TGT AAA TTC TAT AAC Lys Gln Arg Ile Ile Pro Ile Gly Thr Lys Ile Cys Lys Phe Tyr Asn 740 745 750 755	2610
GCG CCC ATT GTC AAG TTC TGG TTT TAC ACA ATA TCA TAC TTG GGC TAC Ala Pro Ile Val Lys Phe Trp Phe Tyr Thr Ile Ser Tyr Leu Gly Tyr 760 765 770	2658
CTG CTG CTG TTT AAC TAC GTC ATC CTG GTG CGG ATG GAT GGC TGG CCG Leu Leu Leu Phe Asn Tyr Val Ile Leu Val Arg Met Asp Gly Trp Pro 775 780 785	2706
TCC CTC CAG GAG TGG ATC GTC ATC TCC TAC ATC GTG AGC CTG GCG TTA Ser Leu Gln Glu Trp Ile Val Ile Ser Tyr Ile Val Ser Leu Ala Leu 790 795 800	2754
GAG AAG ATA CGA GAG ATC CTC ATG TCA GAA CCA GGC AAA CTC AGC CAG Glu Lys Ile Arg Glu Ile Leu Met Ser Glu Pro Gly Lys Leu Ser Gln 805 810 815	2802
AAA ATC AAA GTT TGG CTT CAG GAG TAC TGG AAC ATC ACA GAT CTC GTG Lys Ile Lys Val Trp Leu Gln Glu Tyr Trp Asn Ile Thr Asp Leu Val 820 825 830 835	2850
GCC ATT TCC ACA TTC ATG ATT GGA GCA ATG GCC ACG AGA TCT GTG ATG Ala Ile Ser Thr Phe Met Ile Gly Ala Met Ala Thr Arg Ser Val Met 840 845 850	2898
ATG ATT GGA AAG ATG ATG ATC GAC ATG CTG TAC TTT GTG GTC ATC ATG Met Ile Gly Lys Met Met Ile Asp Met Leu Tyr Phe Val Val Ile Met 855 860 865	2946

CTG	GTC	GTG	CTC	ATG	AGT	TTC	GGA	GTA	GCC	CGT	CAA	GCC	ATT	CTG	CAT	2994
Leu	Val	Val	Leu	Met	Ser	Phe	Gly	Val	Ala	Arg	Gln	Ala	Ile	Leu	His	
		870					875					880				
CCA	GAG	GAG	AAG	CCC	TCT	TGG	AAA	CTG	GCC	CGA	AAC	ATC	TTC	TAC	ATG	3042
Pro	Glu	Glu	Lys	Pro	Ser	Trp	Lys	Leu	Ala	Arg	Asn	Ile	Phe	Tyr	Met	
	885					890					895					
CCC	TAC	TGG	ATG	ATC	TAT	GGA	GAG	GTG	TTT	GCA	GAC	CAG	ATA	GAC	CTC	3090
Pro	Tyr	Trp	Met	Ile	Tyr	Gly	Glu	Val	Phe	Ala	Asp	Gln	Ile	Asp	Leu	
900					905					910					915	
TAC	GCC	ATG	GAA	ATT	AAT	CCT	CCT	TGT	GGT	GAG	AAC	CTA	TAT	GAT	GAG	3138
Tyr	Ala	Met	Glu	Ile	Asn	Pro	Pro	Cys	Gly	Glu	Asn	Leu	Tyr	Asp	Glu	
				920					925					930		
GAG	GGC	AAG	CGG	CTT	CCT	CCC	TGT	ATC	CCC	GGC	GCC	TGG	CTC	ACT	CCA	3186
Glu	Gly	Lys	Arg	Leu	Pro	Pro	Cys	Ile	Pro	Gly	Ala	Trp	Leu	Thr	Pro	
			935					940					945			
GCA	CTC	ATG	GCG	TGC	TAT	CTA	CTG	GTC	GCC	AAC	ATC	CTG	CTG	GTG	AAC	3234
Ala	Leu	Met	Ala	Cys	Tyr	Leu	Leu	Val	Ala	Asn	Ile	Leu	Leu	Val	Asn	
		950					955					960				
CTG	CTG	ATT	GCT	GTG	TTC	AAC	AAT	ACC	TTC	TTT	GAA	GTA	AAA	TCA	ATA	3282
Leu	Leu	Ile	Ala	Val	Phe	Asn	Asn	Thr	Phe	Phe	Glu	Val	Lys	Ser	Ile	
	965					970					975					
TCC	AAC	CAG	GTG	TGG	AAG	TTC	CAG	CGA	TAT	CAG	CTG	ATT	ATG	ACA	TTT	3330
Ser	Asn	Gln	Val	Trp	Lys	Phe	Gln	Arg	Tyr	Gln	Leu	Ile	Met	Thr	Phe	
980					985					990					995	
CAT	GAC	AGG	CCA	GTC	CTG	CCC	CCA	CCG	ATG	ATC	ATT	TTA	AGC	CAC	ATC	3378
His	Asp	Arg	Pro	Val	Leu	Pro	Pro	Pro	Met	Ile	Ile	Leu	Ser	His	Ile	
				1000					1005					1010		
TAC	ATC	ATC	ATT	ATG	CGT	CTC	AGC	GGC	CGC	TGC	AGG	AAA	AAG	AGA	GAA	3426
Tyr	Ile	Ile	Ile	Met	Arg	Leu	Ser	Gly	Arg	Cys	Arg	Lys	Lys	Arg	Glu	
			1015					1020					1025			
GGG	GAC	CAA	GAG	GAA	CGG	GAT	CGT	GGA	TTG	AAG	CTC	TTC	CTT	AGC	GAC	3474
Gly	Asp	Gln	Glu	Glu	Arg	Asp	Arg	Gly	Leu	Lys	Leu	Phe	Leu	Ser	Asp	
		1030					1035					1040				
GAG	GAG	CTA	AAG	AGG	CTG	CAT	GAG	TTC	GAG	GAG	CAG	TGC	GTG	CAG	GAG	3522
Glu	Glu	Leu	Lys	Arg	Leu	His	Glu	Phe	Glu	Glu	Gln	Cys	Val	Gln	Glu	
	1045					1050					1055					
CAC	TTC	CGG	GAG	AAG	GAG	GAT	GAG	CAG	CAG	TCG	TCC	AGC	GAC	GAG	CGC	3570
His	Phe	Arg	Glu	Lys	Glu	Asp	Glu	Gln	Gln	Ser	Ser	Ser	Asp	Glu	Arg	
	1060				1065					1070					1075	
ATC	CGG	GTC	ACT	TCT	GAA	AGA	GTT	GAA	AAT	ATG	TCA	ATG	AGG	TTG	GAA	3618
Ile	Arg	Val	Thr	Ser	Glu	Arg	Val	Glu	Asn	Met	Ser	Met	Arg	Leu	Glu	
				1080					1085					1090		
GAA	ATC	AAT	GAA	AGA	GAA	ACT	TTT	ATG	AAA	ACT	TCC	CTG	CAG	ACT	GTT	3666
Glu	Ile	Asn	Glu	Arg	Glu	Thr	Phe	Met	Lys	Thr	Ser	Leu	Gln	Thr	Val	
			1095					1100					1105			
GAC	CTT	CGA	CTT	GCT	CAG	CTA	GAA	GAA	TTA	TCT	AAC	AGA	ATG	GTG	AAT	3714
Asp	Leu	Arg	Leu	Ala	Gln	Leu	Glu	Glu	Leu	Ser	Asn	Arg	Met	Val	Asn	
		1110					1115					1120				

GCT CTT GAA AAT CTT GCG GGA ATC GAC AGG TCT GAC CTG ATC CAG GCA Ala Leu Glu Asn Leu Ala Gly Ile Asp Arg Ser Asp Leu Ile Gln Ala 1125 1130 1135	3762
CGG TCC CGG GCT TCT TCT GAA TGT GAG GCA ACG TAT CTT CTC CGG CAA Arg Ser Arg Ala Ser Ser Glu Cys Glu Ala Thr Tyr Leu Leu Arg Gln 1140 1145 1150 1155	3810
AGC AGC ATC AAT AGC GCT GAT GGC TAC AGC TTG TAT CGA TAT CAT TTT Ser Ser Ile Asn Ser Ala Asp Gly Tyr Ser Leu Tyr Arg Tyr His Phe 1160 1165 1170	3858
AAC GGA GAA GAG TTA TTA TTT GAG GAT ACA TCT CTC TCC ACG TCA CCA Asn Gly Glu Glu Leu Leu Phe Glu Asp Thr Ser Leu Ser Thr Ser Pro 1175 1180 1185	3906
GGG ACA GGA GTC AGG AAA AAA ACC TGT TCC TTC CGT ATA AAG GAA GAG Gly Thr Gly Val Arg Lys Lys Thr Cys Ser Phe Arg Ile Lys Glu Glu 1190 1195 1200	3954
AAG GAC GTG AAA ACG CAC CTA GTC CCA GAA TGT CAG AAC AGT CTT CAC Lys Asp Val Lys Thr His Leu Val Pro Glu Cys Gln Asn Ser Leu His 1205 1210 1215	4002
CTT TCA CTG GGC ACA AGC ACA TCA GCA ACC CCA GAT GGC AGT CAC CTT Leu Ser Leu Gly Thr Ser Thr Ser Ala Thr Pro Asp Gly Ser His Leu 1220 1225 1230 1235	4050
GCA GTA GAT GAC TTA AAG AAC GCT GAA GAG TCA AAA TTA GGT CCA GAT Ala Val Asp Asp Leu Lys Asn Ala Glu Glu Ser Lys Leu Gly Pro Asp 1240 1245 1250	4098
ATT GGG ATT TCA AAG GAA GAT GAT GAA AGA CAG ACA GAC TCT AAA AAA Ile Gly Ile Ser Lys Glu Asp Asp Glu Arg Gln Thr Asp Ser Lys Lys 1255 1260 1265	4146
GAA GAA ACT ATT TCC CCA AGT TTA AAT AAA ACA GAT GTG ATA CAT GGA Glu Glu Thr Ile Ser Pro Ser Leu Asn Lys Thr Asp Val Ile His Gly 1270 1275 1280	4194
CAG GAC AAA TCA GAT GTT CAA AAC ACT CAG CTA ACA GTG GAA ACG ACA Gln Asp Lys Ser Asp Val Gln Asn Thr Gln Leu Thr Val Glu Thr Thr 1285 1290 1295	4242
AAT ATA GAA GGC ACT ATT TCC TAT CCC CTG GAA GAA ACC AAA ATT ACA Asn Ile Glu Gly Thr Ile Ser Tyr Pro Leu Glu Glu Thr Lys Ile Thr 1300 1305 1310 1315	4290
CGC TAT TTC CCC GAT GAA ACG ATC AAT GCT TGT AAA ACA ATG AAG TCC Arg Tyr Phe Pro Asp Glu Thr Ile Asn Ala Cys Lys Thr Met Lys Ser 1320 1325 1330	4338
AGA AGC TTC GTC TAT TCC CGG GGA AGA AAG CTG GTC GGT GGG GTT AAC Arg Ser Phe Val Tyr Ser Arg Gly Arg Lys Leu Val Gly Gly Val Asn 1335 1340 1345	4386
CAG GAT GTA GAG TAC AGT TCA ATC ACG GAC CAG CAA TTG ACG ACG GAA Gln Asp Val Glu Tyr Ser Ser Ile Thr Asp Gln Gln Leu Thr Thr Glu 1350 1355 1360	4434
TGG CAA TGC CAA GTT CAA AAG ATC ACG CGC TCT CAT AGC ACA GAT ATT Trp Gln Cys Gln Val Gln Lys Ile Thr Arg Ser His Ser Thr Asp Ile 1365 1370 1375	4482

CCT TAC ATT GTG TCG GAA GCT GCA GTG CAA GCT GAG CAA AAA GAG CAG Pro Tyr Ile Val Ser Glu Ala Ala Val Gln Ala Glu Gln Lys Glu Gln 1380 1385 1390 1395	4530
TTT GCA GAT ATG CAA GAT GAA CAC CAT GTC GCT GAA GCA ATT CCT CGA Phe Ala Asp Met Gln Asp Glu His His Val Ala Glu Ala Ile Pro Arg 1400 1405 1410	4578
ATC CCT CGC TTG TCC CTA ACC ATT ACT GAC AGA AAT GGG ATG GAA AAC Ile Pro Arg Leu Ser Leu Thr Ile Thr Asp Arg Asn Gly Met Glu Asn 1415 1420 1425	4626
TTA CTG TCT GTG AAG CCA GAT CAA ACT TTG GGA TTC CCA TCT CTC AGG Leu Leu Ser Val Lys Pro Asp Gln Thr Leu Gly Phe Pro Ser Leu Arg 1430 1435 1440	4674
TCA AAA AGT TTA CAT GGA CAT CCT AGG AAT GTG AAA TCC ATT CAG GGA Ser Lys Ser Leu His Gly His Pro Arg Asn Val Lys Ser Ile Gln Gly 1445 1450 1455	4722
AAG TTA GAC AGA TCT GGA CAT GCC AGT AGT GTA AGC AGC TTA GTA ATT Lys Leu Asp Arg Ser Gly His Ala Ser Ser Val Ser Ser Leu Val Ile 1460 1465 1470 1475	4770
GTG TCT GGA ATG ACA GCA GAA GAA AAA AAG GTT AAG AAA GAG AAA GCT Val Ser Gly Met Thr Ala Glu Glu Lys Lys Val Lys Lys Glu Lys Ala 1480 1485 1490	4818
TCC ACA GAA ACT GAA TGC T AGTCTGTTTT GTTCTTTTAA TTTTTTTTTT Ser Thr Glu Thr Glu Cys 1495	4867
TAACAGTCAG AACCACTAAT GGGTGTCTATC TTGGCCATCC TAAACATCCA TCCAATTTCC	4927
TAAAAACATT TTCCCTT	4944

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1497 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met Tyr Ile Arg Val Ser Tyr Asp Thr Lys Pro Asp Ser Leu Leu His 1 5 10 15
Leu Met Val Lys Asp Trp Gln Leu Glu Leu Pro Lys Leu Leu Ile Ser 20 25 30
Val His Gly Gly Leu Gln Asn Phe Glu Met Gln Pro Lys Leu Lys Gln 35 40 45
Val Phe Gly Lys Gly Leu Ile Lys Ala Ala Met Thr Thr Gly Ala Trp 50 55 60
Ile Phe Thr Gly Gly Val Ser Thr Gly Val Ile Ser His Val Gly Asp 65 70 75 80
Ala Leu Lys Asp His Ser Ser Lys Ser Arg Gly Arg Val Cys Ala Ile 85 90 95

Gly Ile Ala Pro Trp Gly Ile Val Glu Asn Lys Glu Asp Leu Val Gly
 100 105 110
 Lys Asp Val Thr Arg Val Tyr Gln Thr Met Ser Asn Pro Leu Ser Lys
 115 120 125
 Leu Ser Val Leu Asn Asn Ser His Thr His Phe Ile Leu Ala Asp Asn
 130 135 140
 Gly Thr Leu Gly Lys Tyr Gly Ala Glu Val Lys Leu Arg Arg Leu Leu
 145 150 155 160
 Glu Lys His Ile Ser Leu Gln Lys Ile Asn Thr Arg Leu Gly Gln Gly
 165 170 175
 Val Pro Leu Val Gly Leu Val Val Glu Gly Gly Pro Asn Val Val Ser
 180 185 190
 Ile Val Leu Glu Tyr Leu Gln Glu Glu Pro Pro Ile Pro Val Val Ile
 195 200 205
 Cys Asp Gly Ser Gly Arg Ala Ser Asp Ile Leu Ser Phe Ala His Lys
 210 215 220
 Tyr Cys Glu Glu Gly Gly Ile Ile Asn Glu Ser Leu Arg Glu Gln Leu
 225 230 235 240
 Leu Val Thr Ile Gln Lys Thr Phe Asn Tyr Asn Lys Ala Gln Ser His
 245 250 255
 Gln Leu Phe Ala Ile Ile Met Glu Cys Met Lys Lys Lys Glu Leu Val
 260 265 270
 Thr Val Phe Arg Met Gly Ser Glu Gly Gln Gln Asp Ile Glu Met Ala
 275 280 285
 Ile Leu Thr Ala Leu Leu Lys Gly Thr Asn Val Ser Ala Pro Asp Gln
 290 295 300
 Leu Ser Leu Ala Leu Ala Trp Asn Arg Val Asp Ile Ala Arg Ser Gln
 305 310 315 320
 Ile Phe Val Phe Gly Pro His Trp Thr Pro Leu Gly Ser Leu Ala Pro
 325 330 335
 Pro Thr Asp Ser Lys Ala Thr Glu Lys Glu Lys Lys Pro Pro Met Ala
 340 345 350
 Thr Thr Lys Gly Gly Arg Gly Lys Gly Lys Gly Lys Lys Gly Lys
 355 360 365
 Val Lys Glu Glu Val Glu Glu Glu Thr Asp Pro Arg Lys Ile Glu Leu
 370 375 380
 Leu Asn Trp Val Asn Ala Leu Glu Gln Ala Met Leu Asp Ala Leu Val
 385 390 395 400
 Leu Asp Arg Val Asp Phe Val Lys Leu Leu Ile Glu Asn Gly Val Asn
 405 410 415
 Met Gln His Phe Leu Thr Ile Pro Arg Leu Glu Glu Leu Tyr Asn Thr
 420 425 430

Arg	Leu	Gly	Pro	Pro	Asn	Thr	Leu	His	Leu	Leu	Val	Arg	Asp	Val	Lys
		435					440					445			
Lys	Ser	Asn	Leu	Pro	Pro	Asp	Tyr	His	Ile	Ser	Leu	Ile	Asp	Ile	Gly
	450					455					460				
Leu	Val	Leu	Glu	Tyr	Leu	Met	Gly	Gly	Ala	Tyr	Arg	Cys	Asn	Tyr	Thr
465					470					475					480
Arg	Lys	Asn	Phe	Arg	Thr	Leu	Tyr	Asn	Asn	Leu	Phe	Gly	Pro	Lys	Arg
				485					490					495	
Pro	Lys	Ala	Leu	Lys	Leu	Leu	Gly	Met	Glu	Asp	Asp	Glu	Pro	Pro	Ala
			500					505					510		
Lys	Gly	Lys	Lys	Lys	Lys	Lys	Lys	Lys	Lys	Glu	Glu	Glu	Ile	Asp	Ile
		515					520					525			
Asp	Val	Asp	Asp	Pro	Ala	Val	Ser	Arg	Phe	Gln	Tyr	Pro	Phe	His	Glu
530						535					540				
Leu	Met	Val	Trp	Ala	Val	Leu	Met	Lys	Arg	Gln	Lys	Met	Ala	Val	Phe
545					550					555					560
Leu	Trp	Gln	Arg	Gly	Glu	Glu	Ser	Met	Ala	Lys	Ala	Leu	Val	Ala	Cys
				565					570					575	
Lys	Leu	Tyr	Lys	Ala	Met	Ala	His	Glu	Ser	Ser	Glu	Ser	Asp	Leu	Val
			580					585					590		
Asp	Asp	Ile	Ser	Gln	Asp	Leu	Asp	Asn	Asn	Ser	Lys	Asp	Phe	Gly	Gln
		595					600					605			
Leu	Ala	Leu	Glu	Leu	Leu	Asp	Gln	Ser	Tyr	Lys	His	Asp	Glu	Gln	Ile
	610					615					620				
Ala	Met	Lys	Leu	Leu	Thr	Tyr	Glu	Leu	Lys	Asn	Trp	Ser	Asn	Ser	Thr
625					630					635					640
Cys	Leu	Lys	Leu	Ala	Val	Ala	Ala	Lys	His	Arg	Asp	Phe	Ile	Ala	His
				645					650					655	
Thr	Cys	Ser	Gln	Met	Leu	Leu	Thr	Asp	Met	Trp	Met	Gly	Arg	Leu	Arg
			660					665					670		
Met	Arg	Lys	Asn	Pro	Gly	Leu	Lys	Val	Ile	Met	Gly	Ile	Leu	Leu	Pro
		675					680					685			
Pro	Thr	Ile	Leu	Phe	Leu	Glu	Phe	Arg	Thr	Tyr	Asp	Asp	Phe	Ser	Tyr
	690					695					700				
Gln	Thr	Ser	Lys	Glu	Asn	Glu	Asp	Gly	Lys	Glu	Lys	Glu	Glu	Glu	Asn
705					710					715					720
Thr	Asp	Ala	Asn	Ala	Asp	Ala	Gly	Ser	Arg	Lys	Gly	Asp	Glu	Glu	Asn
				725					730					735	
Glu	His	Lys	Lys	Gln	Arg	Ile	Ile	Pro	Ile	Gly	Thr	Lys	Ile	Cys	Lys
			740					745					750		
Phe	Tyr	Asn	Ala	Pro	Ile	Val	Lys	Phe	Trp	Phe	Tyr	Thr	Ile	Ser	Tyr
		755					760					765			

[illegible]

Leu	Gly	Tyr	Leu	Leu	Leu	Phe	Asn	Tyr	Val	Ile	Leu	Val	Arg	Met	Asp		
	770					775					780						
Gly	Trp	Pro	Ser	Leu	Gln	Glu	Trp	Ile	Val	Ile	Ser	Tyr	Ile	Val	Ser		
785					790					795						800	
Leu	Ala	Leu	Glu	Lys	Ile	Arg	Glu	Ile	Leu	Met	Ser	Glu	Pro	Gly	Lys		
				805					810					815			
Leu	Ser	Gln	Lys	Ile	Lys	Val	Trp	Leu	Gln	Glu	Tyr	Trp	Asn	Ile	Thr		
			820					825					830				
Asp	Leu	Val	Ala	Ile	Ser	Thr	Phe	Met	Ile	Gly	Ala	Met	Ala	Thr	Arg		
		835					840					845					
Ser	Val	Met	Met	Ile	Gly	Lys	Met	Met	Ile	Asp	Met	Leu	Tyr	Phe	Val		
	850					855					860						
Val	Ile	Met	Leu	Val	Val	Leu	Met	Ser	Phe	Gly	Val	Ala	Arg	Gln	Ala		
865					870					875					880		
Ile	Leu	His	Pro	Glu	Glu	Lys	Pro	Ser	Trp	Lys	Leu	Ala	Arg	Asn	Ile		
				885					890					895			
Phe	Tyr	Met	Pro	Tyr	Trp	Met	Ile	Tyr	Gly	Glu	Val	Phe	Ala	Asp	Gln		
			900					905					910				
Ile	Asp	Leu	Tyr	Ala	Met	Glu	Ile	Asn	Pro	Pro	Cys	Gly	Glu	Asn	Leu		
	915						920					925					
Tyr	Asp	Glu	Glu	Gly	Lys	Arg	Leu	Pro	Pro	Cys	Ile	Pro	Gly	Ala	Trp		
	930					935					940						
Leu	Thr	Pro	Ala	Leu	Met	Ala	Cys	Tyr	Leu	Leu	Val	Ala	Asn	Ile	Leu		
945					950					955					960		
Leu	Val	Asn	Leu	Leu	Ile	Ala	Val	Phe	Asn	Asn	Thr	Phe	Phe	Glu	Val		
			965					970					975				
Lys	Ser	Ile	Ser	Asn	Gln	Val	Trp	Lys	Phe	Gln	Arg	Tyr	Gln	Leu	Ile		
			980					985					990				
Met	Thr	Phe	His	Asp	Arg	Pro	Val	Leu	Pro	Pro	Pro	Met	Ile	Ile	Leu		
		995					1000					1005					
Ser	His	Ile	Tyr	Ile	Ile	Ile	Met	Arg	Leu	Ser	Gly	Arg	Cys	Arg	Lys		
	1010					1015					1020						
Lys	Arg	Glu	Gly	Asp	Gln	Glu	Glu	Arg	Asp	Arg	Gly	Leu	Lys	Leu	Phe		
1025					1030					1035					1040		
Leu	Ser	Asp	Glu	Glu	Leu	Lys	Arg	Leu	His	Glu	Phe	Glu	Glu	Gln	Cys		
				1045					1050					1055			
Val	Gln	Glu	His	Phe	Arg	Glu	Lys	Glu	Asp	Glu	Gln	Gln	Ser	Ser	Ser		
			1060					1065					1070				
Asp	Glu	Arg	Ile	Arg	Val	Thr	Ser	Glu	Arg	Val	Glu	Asn	Met	Ser	Met		
		1075					1080					1085					
Arg	Leu	Glu	Glu	Ile	Asn	Glu	Arg	Glu	Thr	Phe	Met	Lys	Thr	Ser	Leu		
	1090					1095					1100						

Gln Thr Val Asp Leu Arg Leu Ala Gln Leu Glu Glu Leu Ser Asn Arg	1105	1110	1115	1120
Met Val Asn Ala Leu Glu Asn Leu Ala Gly Ile Asp Arg Ser Asp Leu	1125	1130	1135	
Ile Gln Ala Arg Ser Arg Ala Ser Ser Glu Cys Glu Ala Thr Tyr Leu	1140	1145	1150	
Leu Arg Gln Ser Ser Ile Asn Ser Ala Asp Gly Tyr Ser Leu Tyr Arg	1155	1160	1165	
Tyr His Phe Asn Gly Glu Glu Leu Leu Phe Glu Asp Thr Ser Leu Ser	1170	1175	1180	
Thr Ser Pro Gly Thr Gly Val Arg Lys Lys Thr Cys Ser Phe Arg Ile	1185	1190	1195	1200
Lys Glu Glu Lys Asp Val Lys Thr His Leu Val Pro Glu Cys Gln Asn	1205	1210	1215	
Ser Leu His Leu Ser Leu Gly Thr Ser Thr Ser Ala Thr Pro Asp Gly	1220	1225	1230	
Ser His Leu Ala Val Asp Asp Leu Lys Asn Ala Glu Glu Ser Lys Leu	1235	1240	1245	
Gly Pro Asp Ile Gly Ile Ser Lys Glu Asp Asp Glu Arg Gln Thr Asp	1250	1255	1260	
Ser Lys Lys Glu Glu Thr Ile Ser Pro Ser Leu Asn Lys Thr Asp Val	1265	1270	1275	1280
Ile His Gly Gln Asp Lys Ser Asp Val Gln Asn Thr Gln Leu Thr Val	1285	1290	1295	
Glu Thr Thr Asn Ile Glu Gly Thr Ile Ser Tyr Pro Leu Glu Glu Thr	1300	1305	1310	
Lys Ile Thr Arg Tyr Phe Pro Asp Glu Thr Ile Asn Ala Cys Lys Thr	1315	1320	1325	
Met Lys Ser Arg Ser Phe Val Tyr Ser Arg Gly Arg Lys Leu Val Gly	1330	1335	1340	
Gly Val Asn Gln Asp Val Glu Tyr Ser Ser Ile Thr Asp Gln Gln Leu	1345	1350	1355	1360
Thr Thr Glu Trp Gln Cys Gln Val Gln Lys Ile Thr Arg Ser His Ser	1365	1370	1375	
Thr Asp Ile Pro Tyr Ile Val Ser Glu Ala Ala Val Gln Ala Glu Gln	1380	1385	1390	
Lys Glu Gln Phe Ala Asp Met Gln Asp Glu His His Val Ala Glu Ala	1395	1400	1405	
Ile Pro Arg Ile Pro Arg Leu Ser Leu Thr Ile Thr Asp Arg Asn Gly	1410	1415	1420	
Met Glu Asn Leu Leu Ser Val Lys Pro Asp Gln Thr Leu Gly Phe Pro	1425	1430	1435	1440

Ser Leu Arg Ser Lys Ser Leu His Gly His Pro Arg Asn Val Lys Ser
1445 1450 1455

Ile Gln Gly Lys Leu Asp Arg Ser Gly His Ala Ser Ser Val Ser Ser
1460 1465 1470

Leu Val Ile Val Ser Gly Met Thr Ala Glu Glu Lys Lys Val Lys Lys
1475 1480 1485

Glu Lys Ala Ser Thr Glu Thr Glu Cys
1490 1495

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5055 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

- (ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 346..4945

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

ACTCATTATA GGGNTCGAGC GGCCGCCCCG GCAGGTTTGA GCTGTGCCCT CTCCATTCCA	60
CTGCTGTGGC AGGGTCAGAA ATCTTGATA GAGAAAACCT TTTGCAAACG GGAATGTATC	120
TTTGTAATTC CTAGCACGAA AGACTCTAAC AGGTGTTGCT GTGGCCAGTT CACCAACCAG	180
CATATCCCCC CTCTGCCAAG TGCAACACCC AGCAAAAATG AAGAGGAAAG CAAACAGGTG	240
GAGACTCAGC CTGAGAAATG GTCTGTTGCC AAGCACACCC AGAGCTACCC AACAGATTCC	300
TATGGAGTTC TTGAATTCCA GGGTGGCGGA TATTCCAATA AAGCC ATG TAT ATC	354
Met Tyr Ile	
1	
CGT GTA TCC TAT GAC ACC AAG CCA GAC TCA CTG CTC CAT CTC ATG GTG	402
Arg Val Ser Tyr Asp Thr Lys Pro Asp Ser Leu Leu His Leu Met Val	
5 10 15	
AAA GAT TGG CAG CTG GAA CTC CCC AAG CTC TTA ATA TCT GTG CAT GGA	450
Lys Asp Trp Gln Leu Glu Leu Pro Lys Leu Leu Ile Ser Val His Gly	
20 25 30 35	
GGC CTC CAG AAC TTT GAG ATG CAG CCC AAG CTG AAA CAA GTC TTT GGG	498
Gly Leu Gln Asn Phe Glu Met Gln Pro Lys Leu Lys Gln Val Phe Gly	
40 45 50	
AAA GGC CTG ATC AAG GCT GCT ATG ACC ACC GGG GCC TGG ATC TTC ACC	546
Lys Gly Leu Ile Lys Ala Ala Met Thr Thr Gly Ala Trp Ile Phe Thr	
55 60 65	
GGG GGT GTC AGC ACA GGT GTT ATC AGC CAC GTA GGG GAT GCC TTG AAA	594
Gly Gly Val Ser Thr Gly Val Ile Ser His Val Gly Asp Ala Leu Lys	
70 75 80	

GAC Asp	CAC His 85	TCC Ser	TCC Ser	AAG Lys	TCC Ser	AGA Arg 90	GGC Gly	CGG Arg	GTT Val	TGT Cys	GCT Ala 95	ATA Ile	GGA Gly	ATT Ile	GCT Ala	642
CCA Pro 100	TGG Trp	GGC Gly	ATC Ile	GTG Val	GAG Glu 105	AAT Asn	AAG Lys	GAA Glu	GAC Asp	CTG Leu 110	GTT Val	GGA Gly	AAG Lys	GAT Asp	GTA Val 115	690
ACA Thr	AGA Arg	GTG Val	TAC Tyr	CAG Gln 120	ACC Thr	ATG Met	TCC Ser	AAC Asn	CCT Pro 125	CTA Leu	AGT Ser	AAG Lys	CTC Leu	TCT Ser 130	GTG Val	738
CTC Leu	AAC Asn	AAC Asn	TCC Ser 135	CAC His	ACC Thr	CAC His	TTC Phe	ATC Ile 140	CTG Leu	GCT Ala	GAC Asp	AAT Asn	GGC Gly 145	ACC Thr	CTG Leu	786
GGC Gly	AAG Lys	TAT Tyr 150	GGC Gly	GCC Ala	GAG Glu	GTG Val	AAG Lys 155	CTG Leu	CGA Arg	AGG Arg	CTG Leu	CTG Leu 160	GAA Glu	AAG Lys	CAC His	834
ATC Ile 165	TCC Ser	CTC Leu	CAG Gln	AAG Lys	ATC Ile	AAC Asn 170	ACA Thr	AGA Arg	CTG Leu	GGG Gly 175	CAG Gln	GGC Gly	GTG Val	CCC Pro	CTC Leu	882
GTG Val 180	GGT Gly	CTC Leu	GTG Val	GTG Val	GAG Glu 185	GGG Gly	GGC Gly	CCT Pro	AAC Asn	GTG Val 190	GTG Val	TCC Ser	ATC Ile	GTC Val	TTG Leu 195	930
GAA Glu	TAC Tyr	CTG Leu	CAA Gln	GAA Glu 200	GAG Glu	CCT Pro	CCC Pro	ATC Ile	CCT Pro 205	GTG Val	GTG Val	ATT Ile	TGT Cys	GAT Asp 210	GGC Gly	978
AGC Ser	GGA Gly	CGT Arg	GCC Ala 215	TCG Ser	GAC Asp	ATC Ile	CTG Leu	TCC Ser 220	TTT Phe	GCG Ala	CAC His	AAG Lys	TAC Tyr 225	TGT Cys	GAA Glu	1026
GAA Glu	GGC Gly	GGA Gly 230	ATA Ile	ATA Ile	AAT Asn	GAG Glu	TCC Ser 235	CTC Leu	AGG Arg	GAG Glu	CAG Gln	CTT Leu 240	CTA Leu	GTT Val	ACC Thr	1074
ATT Ile	CAG Gln 245	AAA Lys	ACA Thr	TTT Phe	AAT Asn	TAT Tyr 250	AAT Asn	AAG Lys	GCA Ala	CAA Gln	TCA Ser 255	CAT His	CAG Gln	CTG Leu	TTT Phe	1122
GCA Ala 260	ATT Ile	ATA Ile	ATG Met	GAG Glu	TGC Cys 265	ATG Met	AAG Lys	AAG Lys	AAA Lys	GAA Glu 270	CTC Leu	GTC Val	ACT Thr	GTG Val	TTC Phe 275	1170
AGA Arg	ATG Met	GGT Gly	TCT Ser	GAG Glu 280	GGC Gly	CAG Gln	CAG Gln	GAC Asp	ATC Ile 285	GAG Glu	ATG Met	GCA Ala	ATT Ile	TTA Leu 290	ACT Thr	1218
GCC Ala	CTG Leu	CTG Leu	AAA Lys 295	GGA Gly	ACA Thr	AAC Asn	GTA Val	TCT Ser 300	GCT Ala	CCA Pro	GAT Asp	CAG Gln	CTG Leu 305	AGC Ser	TTG Leu	1266
GCA Ala	CTG Leu	GCT Ala 310	TGG Trp	AAC Asn	CGC Arg	GTG Val	GAC Asp 315	ATA Ile	GCA Ala	CGA Arg	AGC Ser	CAG Gln 320	ATC Ile	TTT Phe	GTC Val	1314
TTT Phe	GGG Gly 325	CCC Pro	CAC His	TGG Trp	ACG Thr	CCC Pro 330	CTG Leu	GGA Gly	AGC Ser	CTG Leu	GCA Ala 335	CCC Pro	CCG Pro	ACG Thr	GAC Asp	1362

AGC Ser 340	AAA Lys	GCC Ala	ACG Thr	GAG Glu	AAG Lys 345	GAG Glu	AAG Lys	AAG Lys	CCA Pro	CCC Pro 350	ATG Met	GCC Ala	ACC Thr	ACC Thr	AAG Lys 355	1410
GGA Gly	GGA Gly	AGA Arg	GGA Gly	AAA Lys 360	GGG Gly	AAA Lys	GGC Gly	AAG Lys	AAG Lys 365	AAA Lys	GGG Gly	AAA Lys	GTG Val	AAA Lys 370	GAG Glu	1458
GAA Glu	GTG Val	GAG Glu	GAA Glu	GAA Glu	ACT Thr	GAC Asp	CCC Pro	CGG Arg 380	AAG Lys	ATA Ile	GAG Glu	CTG Leu	CTG Leu	AAC Asn 385	TGG Trp	1506
GTG Val	AAT Asn	GCT Ala 390	TTG Leu	GAG Glu	CAA Gln	GCG Ala	ATG Met 395	CTA Leu	GAT Asp	GCT Ala	TTA Leu	GTC Val 400	TTA Leu	GAT Asp	CGT Arg	1554
GTC Val	GAC Asp 405	TTT Phe	GTG Val	AAG Lys	CTC Leu	CTG Leu	ATT Ile 410	GAA Glu	AAC Asn	GGA Gly	GTG Val 415	AAC Asn	ATG Met	CAA Gln	CAC His	1602
TTT Phe 420	CTG Leu	ACC Thr	ATT Ile	CCG Pro	AGG Arg 425	CTG Leu	GAG Glu	GAG Glu	CTT Leu	TAT Tyr 430	AAC Asn	ACA Thr	AGA Arg	CTG Leu	GGT Gly 435	1650
CCA Pro	CCA Pro	AAC Asn	ACA Thr	CTT Leu 440	CAT His	CTG Leu	CTG Leu	GTG Val	AGG Arg 445	GAT Asp	GTG Val	AAA Lys	AAG Lys	AGC Ser 450	AAC Asn	1698
CTT Leu	CCG Pro	CCT Pro	GAT Asp 455	TAC Tyr	CAC His	ATC Ile	AGC Ser	CTC Leu	ATA Ile	GAC Asp	ATC Ile	GGG Gly	CTC Leu 465	GTG Val	CTG Leu	1746
GAG Glu	TAC Tyr	CTC Leu 470	ATG Met	GGA Gly	GGA Gly	GCC Ala	TAC Tyr 475	CGC Arg	TGC Cys	AAC Asn	TAC Tyr	ACT Thr 480	CGG Arg	AAA Lys	AAC Asn	1794
TTT Phe 485	CGG Arg	ACC Thr	CTT Leu	TAC Tyr	AAC Asn	AAC Asn 490	TTG Leu	TTT Phe	GGA Gly	CCA Pro	AAG Lys 495	AGG Arg	CCT Pro	AAA Lys	GCT Ala	1842
CTT Leu 500	AAA Lys	CTT Leu	CTG Leu	GGA Gly	ATG Met 505	GAA Glu	GAT Asp	GAT Asp	GAG Glu	CCT Pro 510	CCA Pro	GCT Ala	AAA Lys	GGG Gly 515	AAG Lys	1890
AAA Lys	AAA Lys	AAA Lys	AAA Lys	AAG Lys 520	AAA Lys	AAG Lys	GAG Glu	GAA Glu	GAG Glu 525	ATC Ile	GAC Asp	ATT Ile	GAT Asp	GTG Val 530	GAC Asp	1938
GAC Asp	CCT Pro	GCC Ala	GTG Val 535	AGT Ser	CGG Arg	TTC Phe	CAG Gln	TAT Tyr 540	CCC Pro	TTC Phe	CAC His	GAG Glu	CTG Leu 545	ATG Met	GTG Val	1986
TGG Trp	GCA Ala	GTG Val 550	CTG Leu	ATG Met	AAA Lys	CGC Arg	CAG Gln	AAA Lys	ATG Met 555	GCA Ala	GTG Val	TTC Phe 560	CTC Leu	TGG Trp	CAG Gln	2034
CGA Arg	GGG Gly 565	GAA Glu	GAG Glu	AGC Ser	ATG Met	GCC Ala 570	AAG Lys	GCC Ala	CTG Leu	GTG Val	GCC Ala 575	TGC Cys	AAG Lys	CTC Leu	TAC Tyr	2082
AAG Lys 580	GCC Ala	ATG Met	GCC Ala	CAC His	GAG Glu 585	TCC Ser	TCC Ser	GAG Glu	AGT Ser 590	GAT Asp	CTG Leu	GTG Val	GAT Asp	GAC Asp	ATC Ile 595	2130

TCC Ser	CAG Gln	GAC Asp	TTG Leu	GAT Asp 600	AAC Asn	AAT Asn	TCC Ser	AAA Lys	GAC Asp 605	TTC Phe	GGC Gly	CAG Gln	CTT Leu	GCT Ala 610	TTG Leu	2178
GAG Glu	TTA Leu	TTA Leu	GAC Asp 615	CAG Gln	TCC Ser	TAT Tyr	AAG Lys	CAT His	GAC Asp 620	GAG Glu	CAG Gln	ATC Ile	GCT Ala 625	ATG Met	AAA Lys	2226
CTC Leu	CTG Leu	ACC Thr 630	TAC Tyr	GAG Glu	CTG Leu	AAA Lys	AAC Asn 635	TGG Trp	AGC Ser	AAC Asn	TCG Ser	ACC Thr 640	TGC Cys	CTC Leu	AAA Lys	2274
CTG Leu	GCC Ala 645	GTG Val	GCA Ala	GCC Ala	AAA Lys	CAC His 650	CGG Arg	GAC Asp	TTC Phe	ATT Ile	GCT Ala 655	CAC His	ACC Thr	TGC Cys	AGC Ser	2322
CAG Gln 660	ATG Met	CTG Leu	CTG Leu	ACC Thr	GAT Asp 665	ATG Met	TGG Trp	ATG Met	GGA Gly 670	AGA Arg	CTG Leu	CGG Arg	ATG Met	CGG Arg	AAG Lys 675	2370
AAC Asn	CCC Pro	GGC Gly	CTG Leu 680	AAG Lys	GTT Val	ATC Ile	ATG Met	GGG Gly 685	ATT Ile	CTT Leu	CTA Leu	CCC Pro	CCC Pro	ACC Thr 690	ATC Ile	2418
TTG Leu	TTT Phe	TTG Leu	GAA Glu 695	TTT Phe	CGC Arg	ACA Thr	TAT Tyr	GAT Asp 700	GAT Asp	TTC Phe	TCG Ser	TAT Tyr	CAA Gln 705	ACA Thr	TCC Ser	2466
AAG Lys	GAA Glu 710	AAC Asn	GAG Glu	GAT Asp	GGC Gly	AAA Lys	GAA Glu 715	AAA Lys	GAA Glu	GAG Glu	GAA Glu	AAT Asn 720	ACG Thr	GAT Asp	GCA Ala	2514
AAT Asn	GCA Ala 725	GAT Asp	GCT Ala	GGC Gly	TCA Ser	AGA Arg 730	AAG Lys	GGG Gly	GAT Asp	GAG Glu	GAG Glu	AAC Asn 735	GAG Glu	CAT His	AAA Lys	2562
AAA Lys 740	CAG Gln	AGA Arg	AGT Ser	ATT Ile	CCC Pro 745	ATC Ile	GGA Gly	ACA Thr	AAG Lys 750	ATC Ile	TGT Cys	GAA Glu	TTC Phe	TAT Tyr	AAC Asn 755	2610
GCG Ala	CCC Pro	ATT Ile	GTC Val 760	AAG Lys	TTC Phe	TGG Trp	TTT Phe	TAC Tyr	ACA Thr 765	ATA Ile	TCA Ser	TAC Tyr	TTG Leu	GGC Gly 770	TAC Tyr	2658
CTG Leu	CTG Leu	CTG Leu	TTT Phe 775	AAC Asn	TAC Tyr	GTC Val	ATC Ile 780	CTG Leu	GTG Val	CGG Arg	ATG Met	GAT Asp	GGC Gly 785	TGG Trp	CCG Pro	2706
TCC Ser	CTC Leu	CAG Gln 790	GAG Glu	TGG Trp	ATC Ile	GTC Val 795	ATC Ile 795	TCC Ser	TAC Tyr	ATC Ile	GTG Val	AGC Ser 800	CTG Leu	GCG Ala	TTA Leu	2754
GAG Glu 805	AAG Lys	ATA Ile	CGA Arg	GAG Glu	ATC Ile	CTC Leu 810	ATG Met	TCA Ser	GAA Glu	CCA Pro	GGC Gly 815	AAA Lys	CTC Leu	AGC Ser	CAG Gln	2802
AAA Lys 820	ATC Ile	AAA Lys	GTT Val	TGG Trp	CTT Leu 825	CAG Gln	GAG Glu	TAC Tyr	TGG Trp	AAC Asn 830	ATC Ile	ACA Thr	GAT Asp	CTC Leu	GTG Val 835	2850
GCC Ala	ATT Ile	TCC Ser	ACA Thr 840	TTC Phe	ATG Met	ATT Ile	GGA Gly	GCA Ala	ATT Ile 845	CTT Leu	CGC Arg	CTA Leu	CAG Gln	AAC Asn 850	CAG Gln	2898

CCC	TAC	ATG	GGC	TAT	GGC	CGG	GTG	ATC	TAC	TGT	GTG	GAT	ATC	ATC	TTC	2946
Pro	Tyr	Met	Gly	Tyr	Gly	Arg	Val	Ile	Tyr	Cys	Val	Asp	Ile	Ile	Phe	
			855					860					865			
TGG	TAC	ATC	CGT	GTC	CTG	GAC	ATC	TTT	GGT	GTC	AAC	AAG	TAT	CTG	GGG	2994
Trp	Tyr	Ile	Arg	Val	Leu	Asp	Ile	Phe	Gly	Val	Asn	Lys	Tyr	Leu	Gly	
		870					875					880				
CCA	TAC	GTG	ATG	ATG	ATT	GGA	AAG	ATG	ATG	ATC	GAC	ATG	CTG	TAC	TTT	3042
Pro	Tyr	Val	Met	Met	Ile	Gly	Lys	Met	Met	Ile	Asp	Met	Leu	Tyr	Phe	
		885				890					895					
GTG	GTC	ATC	ATG	CTG	GTC	GTG	CTC	ATG	AGT	TTC	GGA	GTA	GCC	CGT	CAA	3090
Val	Val	Ile	Met	Leu	Val	Val	Leu	Met	Ser	Phe	Gly	Val	Ala	Arg	Gln	
900					905					910					915	
GCC	ATT	CTG	CAT	CCA	GAG	GAG	AAG	CCC	TCT	TGG	AAA	CTG	GCC	CGA	AAC	3138
Ala	Ile	Leu	His	Pro	Glu	Glu	Lys	Pro	Ser	Trp	Lys	Leu	Ala	Arg	Asn	
			920					925						930		
ATC	TTC	TAC	ATG	CCC	TAC	TGG	ATG	ATC	TAT	GGA	GAG	GTG	TTT	GCA	GAC	3186
Ile	Phe	Tyr	Met	Pro	Tyr	Trp	Met	Ile	Tyr	Gly	Glu	Val	Phe	Ala	Asp	
			935					940					945			
CAG	ATA	GAC	CTC	TAC	GCC	ATG	GAA	ATT	AAT	CCT	CCT	TGT	GGT	GAG	AAC	3234
Gln	Ile	Asp	Leu	Tyr	Ala	Met	Glu	Ile	Asn	Pro	Pro	Cys	Gly	Glu	Asn	
		950					955					960				
CTA	TAT	GAT	GAG	GAG	GGC	AAG	CGG	CTT	CCT	CCC	TGT	ATC	CCC	GGC	GCC	3282
Leu	Tyr	Asp	Glu	Glu	Gly	Lys	Arg	Leu	Pro	Pro	Cys	Ile	Pro	Gly	Ala	
		965				970					975					
TGG	CTC	ACT	CCA	GCA	CTC	ATG	GCG	TGC	TAT	CTA	CTG	GTC	GCC	AAC	ATC	3330
Trp	Leu	Thr	Pro	Ala	Leu	Met	Ala	Cys	Tyr	Leu	Leu	Val	Ala	Asn	Ile	
980					985					990					995	
CTG	CTG	GTG	AAC	CTG	CTG	ATT	GCT	GTG	TTC	AAC	AAT	ACT	TTC	TTT	GAA	3378
Leu	Leu	Val	Asn	Leu	Leu	Ile	Ala	Val	Phe	Asn	Asn	Thr	Phe	Phe	Glu	
				1000					1005					1010		
GTA	AAA	TCA	ATA	TCC	AAC	CAG	GTG	TGG	AAG	TTC	CAG	CGA	TAT	CAG	CTG	3426
Val	Lys	Ser	Ile	Ser	Asn	Gln	Val	Trp	Lys	Phe	Gln	Arg	Tyr	Gln	Leu	
			1015					1020					1025			
ATT	ATG	ACA	TTT	CAT	GAC	AGG	CCA	GTC	CTG	CCC	CCA	CCG	ATG	ATC	ATT	3474
Ile	Met	Thr	Phe	His	Asp	Arg	Pro	Val	Leu	Pro	Pro	Pro	Met	Ile	Ile	
			1030				1035						1040			
TTA	AGC	CAC	ATC	TAC	ATC	ATC	ATT	ATG	CGT	CTC	AGC	GGC	CGC	TGC	AGG	3522
Leu	Ser	His	Ile	Tyr	Ile	Ile	Ile	Met	Arg	Leu	Ser	Gly	Arg	Cys	Arg	
		1045				1050					1055					
AAA	AAG	AGA	GAA	GGG	GAC	CAA	GAG	GAA	CGG	GAT	CGT	GGA	TTG	AAG	CTC	3570
Lys	Lys	Arg	Glu	Gly	Asp	Gln	Glu	Glu	Arg	Asp	Arg	Gly	Leu	Lys	Leu	
1060					1065				1070						1075	
TTC	CTT	AGC	GAC	GAG	GAG	CTA	AAG	AGG	CTG	CAT	GAG	TTC	GAG	GAG	CAG	3618
Phe	Leu	Ser	Asp	Glu	Glu	Leu	Lys	Arg	Leu	His	Glu	Phe	Glu	Glu	Gln	
				1080					1085					1090		
TGC	GTG	CAG	GAG	CAC	TTC	CGG	GAG	AAG	GAG	GAT	GAG	CAG	CAG	TCG	TCC	3666
Cys	Val	Gln	Glu	His	Phe	Arg	Glu	Lys	Glu	Asp	Glu	Gln	Gln	Ser	Ser	
			1095					1100					1105			

AGC Ser	GAC Asp	GAG Glu	CGC Arg	ATC Ile	CGG Arg	GTC Val	ACT Thr	TCT Ser	GAA Glu	AGA Arg	GTT Val	GAA Glu	AAT Asn	ATG Met	TCA Ser	3714
		1110					1115					1120				
ATG Met	AGG Arg	TTG Leu	GAA Glu	GAA Glu	ATC Ile	AAT Asn	GAA Glu	AGA Arg	GAA Glu	ACT Thr	TTT Phe	ATG Met	AAA Lys	ACT Thr	TCC Ser	3762
		1125				1130					1135					
CTG Leu	CAG Gln	ACT Thr	GTT Val	GAC Asp	CTT Leu	CGA Arg	CTT Leu	GCT Ala	CAG Gln	CTA Leu	GAA Glu	GAA Glu	TTA Leu	TCT Ser	AAC Asn	3810
					1140	1145			1150						1155	
AGA Arg	ATG Met	GTG Val	AAT Asn	GCT Ala	CTT Leu	GAA Glu	AAT Asn	CTT Leu	GCG Ala	GGA Gly	ATC Ile	GAC Asp	AGG Arg	TCT Ser	GAC Asp	3858
				1160					1165					1170		
CTG Leu	ATC Ile	CAG Gln	GCA Ala	CGG Arg	TCC Ser	CGG Arg	GCT Ala	TCT Ser	TCT Ser	GAA Glu	TGT Cys	GAG Glu	GCA Ala	ACG Thr	TAT Tyr	3906
			1175				1180						1185			
CTT Leu	CTC Leu	CGG Arg	CAA Gln	AGC Ser	AGC Ser	ATC Ile	AAT Asn	AGC Ser	GCT Ala	GAT Asp	GGC Gly	TAC Tyr	AGC Ser	TTG Leu	TAT Tyr	3954
		1190					1195					1200				
CGA Arg	TAT Tyr	CAT His	TTT Phe	AAC Asn	GGA Gly	GAA Glu	GAG Glu	TTA Leu	TTA Leu	TTT Phe	GAG Glu	GAT Asp	ACA Thr	TCT Ser	CTC Leu	4002
		1205				1210					1215					
TCC Ser	ACG Thr	TCA Ser	CCA Pro	GGG Gly	ACA Thr	GGA Gly	GTC Val	AGG Arg	AAA Lys	AAA Lys	ACC Thr	TGT Cys	TCC Ser	TTC Phe	CGT Arg	4050
				1220		1225			1230						1235	
ATA Ile	AAG Lys	GAA Glu	GAG Glu	AAG Lys	GAC Asp	GTG Val	AAA Lys	ACG Thr	CAC His	CTA Leu	GTC Val	CCA Pro	GAA Glu	TGT Cys	CAG Gln	4098
				1240					1245					1250		
AAC Asn	AGT Ser	CTT Leu	CAC His	CTT Leu	TCA Ser	CTG Leu	GGC Gly	ACA Thr	AGC Ser	ACA Thr	TCA Ser	GCA Ala	ACC Thr	CCA Pro	GAT Asp	4146
			1255				1260					1265				
GGC Gly	AGT Ser	CAC His	CTT Leu	GCA Ala	GTA Val	GAT Asp	GAC Asp	TTA Leu	AAG Lys	AAC Asn	GCT Ala	GAA Glu	GAG Glu	TCA Ser	AAA Lys	4194
		1270				1275					1280					
TTA Leu	GGT Gly	CCA Pro	GAT Asp	ATT Ile	GGG Gly	ATT Ile	TCA Ser	AAG Lys	GAA Glu	GAT Asp	GAT Asp	GAA Glu	AGA Arg	CAG Gln	ACA Thr	4242
		1285				1290					1295					
GAC Asp	TCT Ser	AAA Lys	AAA Lys	GAA Glu	GAA Glu	ACT Thr	ATT Ile	TCC Ser	CCA Pro	AGT Ser	TTA Leu	AAT Asn	AAA Lys	ACA Thr	GAT Asp	4290
		1300				1305			1310					1315		
GTG Val	ATA Ile	CAT His	GGA Gly	CAG Gln	GAC Asp	AAA Lys	TCA Ser	GAT Asp	GTT Val	CAA Gln	AAC Asn	ACT Thr	CAG Gln	CTA Leu	ACA Thr	4338
				1320				1325						1330		
GTG Val	GAA Glu	ACG Thr	ACA Thr	AAT Asn	ATA Ile	GAA Glu	GGC Gly	ACT Thr	ATT Ile	TCC Ser	TAT Tyr	CCC Pro	CTG Leu	GAA Glu	GAA Glu	4386
			1335				1340						1345			
ACC Thr	AAA Lys	ATT Ile	ACA Thr	CGC Arg	TAT Tyr	TTC Phe	CCC Pro	GAT Asp	GAA Glu	ACG Thr	ATC Ile	AAT Asn	GCT Ala	TGT Cys	AAA Lys	4434
			1350				1355					1360				

ACA ATG AAG TCC AGA AGC TTC GTC TAT TCC CGG GGA AGA AAG CTG GTC Thr Met Lys Ser Arg Ser Phe Val Tyr Ser Arg Gly Arg Lys Leu Val 1365 1370 1375	4482
GGT GGG GTT AAC CAG GAT GTA GAG TAC AGT TCA ATC ACG GAC CAG CAA Gly Gly Val Asn Gln Asp Val Glu Tyr Ser Ser Ile Thr Asp Gln Gln 1380 1385 1390 1395	4530
TTG ACG ACG GAA TGG CAA TGC CAA GTT CAA AAG ATC ACG CGC TCT CAT Leu Thr Thr Glu Trp Gln Cys Gln Val Gln Lys Ile Thr Arg Ser His 1400 1405 1410	4578
AGC ACA GAT ATT CCT TAC ATT GTG TCG GAA GCT GCA GTG CAA GCT GAG Ser Thr Asp Ile Pro Tyr Ile Val Ser Glu Ala Ala Val Gln Ala Glu 1415 1420 1425	4626
CAA AAA GAG CAG TTT GCA GAT ATG CAA GAT GAA CAC CAT GTC GCT GAA Gln Lys Glu Gln Phe Ala Asp Met Gln Asp Glu His His Val Ala Glu 1430 1435 1440	4674
GCA ATT CCT CGA ATC CCT CGC TTG TCC CTA ACC ATT ACT GAC AGA AAT Ala Ile Pro Arg Ile Pro Arg Leu Ser Leu Thr Ile Thr Asp Arg Asn 1445 1450 1455	4722
GGG ATG GAA AAC TTA CTG TCT GTG AAG CCA GAT CAA ACT TTG GGA TTC Gly Met Glu Asn Leu Leu Ser Val Lys Pro Asp Gln Thr Leu Gly Phe 1460 1465 1470 1475	4770
CCA TCT CTC AGG TCA AAA AGT TTA CAT GGA CAT CCT AGG AAT GTG AAA Pro Ser Leu Arg Ser Lys Ser Leu His Gly His Pro Arg Asn Val Lys 1480 1485 1490	4818
TCC ATT CAG GGA AAG TTA GAC AGA TCT GGA CAT GCC AGT AGT GTA AGC Ser Ile Gln Gly Lys Leu Asp Arg Ser Gly His Ala Ser Ser Val Ser 1495 1500 1505	4866
AGC TTA GTA ATT GTG TCT GGA ATG ACA GCA GAA GAA AAA AAG GTT AAG Ser Leu Val Ile Val Ser Gly Met Thr Ala Glu Glu Lys Lys Val Lys 1510 1515 1520	4914
AAA GAG AAA GCT TCC ACA GAA ACT GAA TGC T AGTCTGTTTT GTTCTTTAA Lys Glu Lys Ala Ser Thr Glu Thr Glu Cys 1525 1530	4965
TTTTTTTTTTT TAACAGTCAG AAACCCACTA ATGGGTGTCA TCTTGGCCCA TCCTAAACAC	5025
ATMTCCAATT TCCTAAAAAC ATTTTCCCTT	5055

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1533 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Met Tyr Ile Arg Val Ser Tyr Asp Thr Lys Pro Asp Ser Leu Leu His 1 5 10 15
Leu Met Val Lys Asp Trp Gln Leu Glu Leu Pro Lys Leu Leu Ile Ser 20 25 30

Val His Gly Gly Leu Gln Asn Phe Glu Met Gln Pro Lys Leu Lys Gln
35 40 45
Val Phe Gly Lys Gly Leu Ile Lys Ala Ala Met Thr Thr Gly Ala Trp
50 55 60
Ile Phe Thr Gly Gly Val Ser Thr Gly Val Ile Ser His Val Gly Asp
65 70 75 80
Ala Leu Lys Asp His Ser Ser Lys Ser Arg Gly Arg Val Cys Ala Ile
85 90 95
Gly Ile Ala Pro Trp Gly Ile Val Glu Asn Lys Glu Asp Leu Val Gly
100 105 110
Lys Asp Val Thr Arg Val Tyr Gln Thr Met Ser Asn Pro Leu Ser Lys
115 120 125
Leu Ser Val Leu Asn Asn Ser His Thr His Phe Ile Leu Ala Asp Asn
130 135 140
Gly Thr Leu Gly Lys Tyr Gly Ala Glu Val Lys Leu Arg Arg Leu Leu
145 150 155 160
Glu Lys His Ile Ser Leu Gln Lys Ile Asn Thr Arg Leu Gly Gln Gly
165 170 175
Val Pro Leu Val Gly Leu Val Val Glu Gly Gly Pro Asn Val Val Ser
180 185 190
Ile Val Leu Glu Tyr Leu Gln Glu Glu Pro Pro Ile Pro Val Val Ile
195 200 205
Cys Asp Gly Ser Gly Arg Ala Ser Asp Ile Leu Ser Phe Ala His Lys
210 215 220
Tyr Cys Glu Glu Gly Gly Ile Ile Asn Glu Ser Leu Arg Glu Gln Leu
225 230 235 240
Leu Val Thr Ile Gln Lys Thr Phe Asn Tyr Asn Lys Ala Gln Ser His
245 250 255
Gln Leu Phe Ala Ile Ile Met Glu Cys Met Lys Lys Lys Glu Leu Val
260 265 270
Thr Val Phe Arg Met Gly Ser Glu Gly Gln Gln Asp Ile Glu Met Ala
275 280 285
Ile Leu Thr Ala Leu Leu Lys Gly Thr Asn Val Ser Ala Pro Asp Gln
290 295 300
Leu Ser Leu Ala Leu Ala Trp Asn Arg Val Asp Ile Ala Arg Ser Gln
305 310 315 320
Ile Phe Val Phe Gly Pro His Trp Thr Pro Leu Gly Ser Leu Ala Pro
325 330 335
Pro Thr Asp Ser Lys Ala Thr Glu Lys Glu Lys Lys Pro Pro Met Ala
340 345 350
Thr Thr Lys Gly Gly Arg Gly Lys Gly Lys Gly Lys Lys Lys Gly Lys
355 360 365

26E250" 24429830

Val	Lys	Glu	Glu	Val	Glu	Glu	Glu	Thr	Asp	Pro	Arg	Lys	Ile	Glu	Leu
370							375				380				
Leu	Asn	Trp	Val	Asn	Ala	Leu	Glu	Gln	Ala	Met	Leu	Asp	Ala	Leu	Val
385					390					395					400
Leu	Asp	Arg	Val	Asp	Phe	Val	Lys	Leu	Leu	Ile	Glu	Asn	Gly	Val	Asn
				405					410					415	
Met	Gln	His	Phe	Leu	Thr	Ile	Pro	Arg	Leu	Glu	Glu	Leu	Tyr	Asn	Thr
			420					425					430		
Arg	Leu	Gly	Pro	Pro	Asn	Thr	Leu	His	Leu	Leu	Val	Arg	Asp	Val	Lys
		435					440					445			
Lys	Ser	Asn	Leu	Pro	Pro	Asp	Tyr	His	Ile	Ser	Leu	Ile	Asp	Ile	Gly
	450					455					460				
Leu	Val	Leu	Glu	Tyr	Leu	Met	Gly	Gly	Ala	Tyr	Arg	Cys	Asn	Tyr	Thr
465					470					475					480
Arg	Lys	Asn	Phe	Arg	Thr	Leu	Tyr	Asn	Asn	Leu	Phe	Gly	Pro	Lys	Arg
				485					490					495	
Pro	Lys	Ala	Leu	Lys	Leu	Leu	Gly	Met	Glu	Asp	Asp	Glu	Pro	Pro	Ala
			500					505					510		
Lys	Gly	Lys	Lys	Lys	Lys	Lys	Lys	Lys	Lys	Glu	Glu	Glu	Ile	Asp	Ile
		515						520				525			
Asp	Val	Asp	Asp	Pro	Ala	Val	Ser	Arg	Phe	Gln	Tyr	Pro	Phe	His	Glu
	530					535					540				
Leu	Met	Val	Trp	Ala	Val	Leu	Met	Lys	Arg	Gln	Lys	Met	Ala	Val	Phe
545					550					555					560
Leu	Trp	Gln	Arg	Gly	Glu	Glu	Ser	Met	Ala	Lys	Ala	Leu	Val	Ala	Cys
				565					570					575	
Lys	Leu	Tyr	Lys	Ala	Met	Ala	His	Glu	Ser	Ser	Glu	Ser	Asp	Leu	Val
			580					585					590		
Asp	Asp	Ile	Ser	Gln	Asp	Leu	Asp	Asn	Asn	Ser	Lys	Asp	Phe	Gly	Gln
		595					600					605			
Leu	Ala	Leu	Glu	Leu	Leu	Asp	Gln	Ser	Tyr	Lys	His	Asp	Glu	Gln	Ile
	610					615					620				
Ala	Met	Lys	Leu	Leu	Thr	Tyr	Glu	Leu	Lys	Asn	Trp	Ser	Asn	Ser	Thr
625					630					635					640
Cys	Leu	Lys	Leu	Ala	Val	Ala	Ala	Lys	His	Arg	Asp	Phe	Ile	Ala	His
				645					650					655	
Thr	Cys	Ser	Gln	Met	Leu	Leu	Thr	Asp	Met	Trp	Met	Gly	Arg	Leu	Arg
			660					665					670		
Met	Arg	Lys	Asn	Pro	Gly	Leu	Lys	Val	Ile	Met	Gly	Ile	Leu	Leu	Pro
		675					680					685			
Pro	Thr	Ile	Leu	Phe	Leu	Glu	Phe	Arg	Thr	Tyr	Asp	Asp	Phe	Ser	Tyr
	690					695					700				

26E250" 24429880

Gln Thr Ser Lys Glu Asn Glu Asp Gly Lys Glu Lys Glu Glu Glu Asn
 705 710 715 720
 Thr Asp Ala Asn Ala Asp Ala Gly Ser Arg Lys Gly Asp Glu Glu Asn
 725 730 735
 Glu His Lys Lys Gln Arg Ser Ile Pro Ile Gly Thr Lys Ile Cys Glu
 740 745 750
 Phe Tyr Asn Ala Pro Ile Val Lys Phe Trp Phe Tyr Thr Ile Ser Tyr
 755 760 765
 Leu Gly Tyr Leu Leu Leu Phe Asn Tyr Val Ile Leu Val Arg Met Asp
 770 775 780
 Gly Trp Pro Ser Leu Gln Glu Trp Ile Val Ile Ser Tyr Ile Val Ser
 785 790 795 800
 Leu Ala Leu Glu Lys Ile Arg Glu Ile Leu Met Ser Glu Pro Gly Lys
 805 810 815
 Leu Ser Gln Lys Ile Lys Val Trp Leu Gln Glu Tyr Trp Asn Ile Thr
 820 825 830
 Asp Leu Val Ala Ile Ser Thr Phe Met Ile Gly Ala Ile Leu Arg Leu
 835 840 845
 Gln Asn Gln Pro Tyr Met Gly Tyr Gly Arg Val Ile Tyr Cys Val Asp
 850 855 860
 Ile Ile Phe Trp Tyr Ile Arg Val Leu Asp Ile Phe Gly Val Asn Lys
 865 870 875 880
 Tyr Leu Gly Pro Tyr Val Met Met Ile Gly Lys Met Met Ile Asp Met
 885 890 895
 Leu Tyr Phe Val Val Ile Met Leu Val Val Leu Met Ser Phe Gly Val
 900 905 910
 Ala Arg Gln Ala Ile Leu His Pro Glu Glu Lys Pro Ser Trp Lys Leu
 915 920 925
 Ala Arg Asn Ile Phe Tyr Met Pro Tyr Trp Met Ile Tyr Gly Glu Val
 930 935 940
 Phe Ala Asp Gln Ile Asp Leu Tyr Ala Met Glu Ile Asn Pro Pro Cys
 945 950 955 960
 Gly Glu Asn Leu Tyr Asp Glu Glu Gly Lys Arg Leu Pro Pro Cys Ile
 965 970 975
 Pro Gly Ala Trp Leu Thr Pro Ala Leu Met Ala Cys Tyr Leu Leu Val
 980 985 990
 Ala Asn Ile Leu Leu Val Asn Leu Leu Ile Ala Val Phe Asn Asn Thr
 995 1000 1005
 Phe Phe Glu Val Lys Ser Ile Ser Asn Gln Val Trp Lys Phe Gln Arg
 1010 1015 1020
 Tyr Gln Leu Ile Met Thr Phe His Asp Arg Pro Val Leu Pro Pro Pro
 1025 1030 1035 1040

266350 24429830

Met	Ile	Ile	Leu	Ser	His	Ile	Tyr	Ile	Ile	Ile	Met	Arg	Leu	Ser	Gly	
				1045					1050						1055	
Arg	Cys	Arg	Lys	Lys	Arg	Glu	Gly	Asp	Gln	Glu	Glu	Arg	Asp	Arg	Gly	
			1060					1065					1070			
Leu	Lys	Leu	Phe	Leu	Ser	Asp	Glu	Glu	Leu	Lys	Arg	Leu	His	Glu	Phe	
		1075					1080					1085				
Glu	Glu	Gln	Cys	Val	Gln	Glu	His	Phe	Arg	Glu	Lys	Glu	Asp	Glu	Gln	
	1090					1095					1100					
Gln	Ser	Ser	Ser	Asp	Glu	Arg	Ile	Arg	Val	Thr	Ser	Glu	Arg	Val	Glu	
1105					1110					1115					1120	
Asn	Met	Ser	Met	Arg	Leu	Glu	Glu	Ile	Asn	Glu	Arg	Glu	Thr	Phe	Met	
				1125					1130					1135		
Lys	Thr	Ser	Leu	Gln	Thr	Val	Asp	Leu	Arg	Leu	Ala	Gln	Leu	Glu	Glu	
			1140					1145					1150			
Leu	Ser	Asn	Arg	Met	Val	Asn	Ala	Leu	Glu	Asn	Leu	Ala	Gly	Ile	Asp	
		1155					1160					1165				
Arg	Ser	Asp	Leu	Ile	Gln	Ala	Arg	Ser	Arg	Ala	Ser	Ser	Glu	Cys	Glu	
	1170					1175					1180					
Ala	Thr	Tyr	Leu	Leu	Arg	Gln	Ser	Ser	Ile	Asn	Ser	Ala	Asp	Gly	Tyr	
1185					1190					1195					1200	
Ser	Leu	Tyr	Arg	Tyr	His	Phe	Asn	Gly	Glu	Glu	Leu	Leu	Phe	Glu	Asp	
				1205					1210					1215		
Thr	Ser	Leu	Ser	Thr	Ser	Pro	Gly	Thr	Gly	Val	Arg	Lys	Lys	Thr	Cys	
			1220					1225					1230			
Ser	Phe	Arg	Ile	Lys	Glu	Glu	Lys	Asp	Val	Lys	Thr	His	Leu	Val	Pro	
		1235					1240					1245				
Glu	Cys	Gln	Asn	Ser	Leu	His	Leu	Ser	Leu	Gly	Thr	Ser	Thr	Ser	Ala	
	1250					1255					1260					
Thr	Pro	Asp	Gly	Ser	His	Leu	Ala	Val	Asp	Asp	Leu	Lys	Asn	Ala	Glu	
1265					1270					1275					1280	
Glu	Ser	Lys	Leu	Gly	Pro	Asp	Ile	Gly	Ile	Ser	Lys	Glu	Asp	Asp	Glu	
				1285					1290					1295		
Arg	Gln	Thr	Asp	Ser	Lys	Lys	Glu	Glu	Thr	Ile	Ser	Pro	Ser	Leu	Asn	
			1300					1305					1310			
Lys	Thr	Asp	Val	Ile	His	Gly	Gln	Asp	Lys	Ser	Asp	Val	Gln	Asn	Thr	
		1315					1320					1325				
Gln	Leu	Thr	Val	Glu	Thr	Thr	Asn	Ile	Glu	Gly	Thr	Ile	Ser	Tyr	Pro	
	1330					1335					1340					
Leu	Glu	Glu	Thr	Lys	Ile	Thr	Arg	Tyr	Phe	Pro	Asp	Glu	Thr	Ile	Asn	
1345					1350					1355					1360	
Ala	Cys	Lys	Thr	Met	Lys	Ser	Arg	Ser	Phe	Val	Tyr	Ser	Arg	Gly	Arg	
				1365					1370					13		

[illegible]

Lys	Leu	Val	Gly	Gly	Val	Asn	Gln	Asp	Val	Glu	Tyr	Ser	Ser	Ile	Thr
			1380					1385						1390	
Asp	Gln	Gln	Leu	Thr	Thr	Glu	Trp	Gln	Cys	Gln	Val	Gln	Lys	Ile	Thr
			1395				1400						1405		
Arg	Ser	His	Ser	Thr	Asp	Ile	Pro	Tyr	Ile	Val	Ser	Glu	Ala	Ala	Val
			1410				1415					1420			
Gln	Ala	Glu	Gln	Lys	Glu	Gln	Phe	Ala	Asp	Met	Gln	Asp	Glu	His	His
					1430					1435					1440
Val	Ala	Glu	Ala	Ile	Pro	Arg	Ile	Pro	Arg	Leu	Ser	Leu	Thr	Ile	Thr
				1445					1450					1455	
Asp	Arg	Asn	Gly	Met	Glu	Asn	Leu	Leu	Ser	Val	Lys	Pro	Asp	Gln	Thr
			1460					1465					1470		
Leu	Gly	Phe	Pro	Ser	Leu	Arg	Ser	Lys	Ser	Leu	His	Gly	His	Pro	Arg
			1475					1480				1485			
Asn	Val	Lys	Ser	Ile	Gln	Gly	Lys	Leu	Asp	Arg	Ser	Gly	His	Ala	Ser
			1490				1495					1500			
Ser	Val	Ser	Ser	Leu	Val	Ile	Val	Ser	Gly	Met	Thr	Ala	Glu	Glu	Lys
					1510					1515					1520
Lys	Val	Lys	Lys	Glu	Lys	Ala	Ser	Thr	Glu	Thr	Glu	Cys			
				1525						1530					

[illegible]

APPLICATION
FOR
UNITED STATES LETTERS PATENT

TITLE: **COMPOSITIONS AND METHODS FOR THE DIAGNOSIS,
PREVENTION AND TREATMENT OF TUMOR PROGRESSION**

APPLICANT: **ANDREW W. SHYJAN**

"EXPRESS MAIL" Mailing Label Number RB656730558US

Date of Deposit March 29, 1996

I hereby certify under 37 CFR 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office To Addressee" with sufficient postage on the date indicated above and is addressed to the Commissioner of Patents and Trademarks, Washington, D.C. 20231.

Mary Jane DiPalma

Mary Jane DiPalma

**COMPOSITIONS AND METHODS FOR THE DIAGNOSIS,
PREVENTION AND TREATMENT OF TUMOR PROGRESSION**

5 This is a continuation-in-part of U.S. Serial No.
08/412,431.

1. INTRODUCTION

268244-0330
468250-24429880

The present invention relates to methods and compositions for the diagnosis, prevention and treatment of tumor progression in mammals, for example, humans. The different types of tumors may include, but are not limited to, human melanomas, breast, gastrointestinal tumors such as esophageal, stomach, duodenal, colon, colorectal and rectal cancers, prostate, bladder, testicular, ovarian, uterine, cervical, brain, lung, bronchial, larynx, pharynx, liver, pancreatic, thyroid, bone, various types of skin cancers and neoplastic conditions such as leukemias and lymphomas. Specifically, genes which are differentially expressed in tumor cells relative to normal cells and/or relative to tumor cells at a different stage of tumor progression are identified. For example, genes are identified which are differentially expressed in benign (e.g., non-malignant) tumor cells relative to malignant tumor cells exhibiting a high metastatic potential. Genes are also identified via the ability of their gene products to interact with gene products involved in the progression to and/or aggressiveness of neoplastic tumor disease states. The genes identified can be used diagnostically or as targets for therapeutic intervention. In this regard, the present invention provides methods for the identification of compounds useful in the diagnosis, prevention and therapeutic treatment of tumor progression, including, for example, metastatic neoplastic disorders. The present invention also provides methods for the identification of compounds useful in the diagnosis, prevention and

10
15
20
25
30
35

therapeutic treatment of tumor progression, including, for example, pre-neoplastic and/or benign states. Additionally, methods are provided for the diagnostic evaluation and prognosis of conditions involving tumor progression, for the identification of subjects exhibiting a predisposition to such conditions, for monitoring patients undergoing clinical evaluation for the prevention and treatment of tumor progression disorders, and for monitoring the efficacy of compounds used in clinical trials.

2. BACKGROUND OF THE INVENTION

Cancer is the second leading cause of death in the United States, after heart disease (Boring, C.C. et al., 1993, CA Cancer J. Clin. 43:7), and develops in one in three Americans, and one of every four Americans dies of cancer. Cancer is characterized primarily by an increase in the number of abnormal, or neoplastic, cells derived from a given normal tissue which proliferate to form a tumor mass, the invasion of adjacent tissues by these neoplastic tumor cells, and the generation of malignant cells which spread via the blood or lymphatic system to regional lymph nodes and to distant sites. The latter progression to malignancy is referred to as metastasis.

Cancer can be viewed as a breakdown in the communication between tumor cells and their environment, including their normal neighboring cells. Signals, both growth-stimulatory and growth-inhibitory, are routinely exchanged between cells within a tissue. Normally, cells do not divide in the absence of stimulatory signals, and, likewise, will cease dividing in the presence of inhibitory signals. In a cancerous, or neoplastic, state, a cell acquires the ability to "override" these signals and to proliferate under conditions in which normal cells would not grow.

2686244 03397
Tumor cells must acquire a number of distinct aberrant traits to proliferate. Reflecting this requirement is the fact that the genomes of certain well-studied tumors carry several different independently altered genes, including activated oncogenes and inactivated tumor suppressor genes. Each of these genetic changes appears to be responsible for imparting some of the traits that, in aggregate, represent the full neoplastic phenotype (Land, H. et al., 1983, Science 222:771; Ruley, H.E., 1983, Nature 304:602; Hunter, T., 1991, Cell 64:249).

In addition to unhindered cell proliferation, cells must acquire several traits for tumor progression to occur. For example, early on in tumor progression, cells must evade the host immune system. Further, as tumor mass increases, the tumor must acquire vasculature to supply nourishment and remove metabolic waste. Additionally, cells must acquire an ability to invade adjacent tissue, and, ultimately, cells often acquire the capacity to metastasize to distant sites.

The biochemical basis for immune recognition of tumor cells is unclear. It is possible that the tumorigenicity of cells can increase when the cells' display of Class I histocompatibility antigens is reduced (Schrier, P.I. et al., 1983, Nature 305:771), in that these antigens, in conjunction with tumor-specific antigens are required for the tumor cells to be recognized by cytotoxic T lymphocytes (CTLs). Tumor cells which have lost one or more genes encoding tumor-specific antigens seem to escape recognition by the corresponding reactive CTLs (Van der Bruggen, P. et al., 1991, Science 254:1643).

Once a tumor reaches more than about 1 mm in diameter, it can no longer rely on passive diffusion for nutrition and removal of metabolic waste. At this point, the tumor mass must make intimate contact with the

circulatory system. Thus, cells within more advanced tumors secrete angiogenic factors which promote neovascularization, i.e., the growth of blood vessels from surrounding tissue into the tumor mass (Folkman, J. and Klagsburn, M., 1987, Science 235:442; Liotta, L.A. et al., 1991, Cell 64:327). Among these angiogenic factors are the fibroblast growth factor (FGF) and endothelial cell growth factor (ECGF). Neovascularization can, in fact, be an essential precursor to metastasis. First, the process is required for a large increase in tumor cell number, which in turn, allows the appearance of rare metastatic variants. Further, neovascularization provides a direct portal entry into the circulatory system which can be used by metastasizing cells.

A variety of biochemical factors have been associated with different phases of metastases. Cell surface receptors for collagen, glycoproteins such as laminin, or proteoglycans, facilitate tumor cell attachment, an important step in invasion and metastases. Attachment then triggers the release of degradative enzymes which facilitate the penetration of tumor cells through tissue barriers. Once the tumor cell has entered the target tissue, specific growth factors are required for further proliferation.

It is apparent that the complex process of tumor progression must involve multiple gene products. It is therefore important to define the role of specific genes involved in tumor progression, to identify those gene products involved in the tumor progression process and to further identify those gene products which can serve as therapeutic targets for the diagnosis, prevention and treatment of metastases of various forms of cancers.

Some attempts have been made to study genes which are thought to elicit or augment tumor progression

phenotypes. Mutations may drive a wave of cellular multiplication associated with gradual increases in tumor size, disorganization and malignancy. For example, a mutation in the tumor suppressor gene which is a negative regulator of cellular proliferation, results in a loss of crucial control over tumor growth and progression. Differential expression of the following suppressor genes has been demonstrated in human cancers: the retinoblastoma gene, RB; the Wilms' tumor gene, WT1 (11p); the gene deleted in colon carcinoma, DCC (18q); the neurofibromatosis type 1 gene, NF1 (17q); and the gene involved in familial adenomatous polyposis coli, APC (5q) (Vogelstein, B. and Kinzler, K.W., 1993, Trends Genet. 9:138-141).

Insight into the complex events that lead from normal cellular growth to neoplasia, invasion and metastasis is crucial for the development of effective diagnostic and therapeutic strategies. The foregoing studies are aimed at defining the role of particular gene products presumed to be involved in tumor progression. However, such approaches cannot identify the full panoply of gene products that are involved in the cascade of steps in tumor progression. A great need, therefore, exists for the successful identification of those genes which are differentially expressed in cells involved in or predisposed to a tumor progression phenotype. Such differentially expressed gene and/or gene products can represent useful diagnostic markers and/or therapeutic targets for tumor progression disorders. With respect to diagnostic techniques, such genes and/or gene products could represent useful markers for the diagnosis, especially early diagnosis, given the correlation between early diagnosis and successful cancer treatment. With respect to therapeutic treatments, such differentially expressed genes and/or gene products could represent useful

targets for therapeutic treatment of various forms of tumor progression disorders, including metastatic and non-metastatic neoplastic disorders, and for inhibiting the progression of pre-neoplastic lesions (e.g., hyperplastic lesions or other benign tumors) to malignant tumors.

Differentially expressed genes involved in tumor metastasis have been identified using murine melanoma cell lines of varying metastatic potentials, N-nitroso-methylurea-induced rat mammary carcinomas, mammary carcinoma cell lines, human breast tumors and spontaneous colonic and intestinal tumors in mice (Steeg, P.S., et al., 1988, J. Natl. Cancer Inst. 80:200-204; Qian, F., et al., 1994, Cell 77:335-347; Leone, A., et al., 1991, 65:25-35; Zou, Z., et al., 1994, Science 263:526-529; and Fodde, R., et al., 1994, Proc. Natl. Acad. Sci. USA 91:8969-8973).

3. SUMMARY OF THE INVENTION

The present invention relates to methods and compositions for diagnosis, prevention, and treatment of tumor progression. Specifically, murine and human genes are identified and described which are differentially expressed in tumor cells relative to normal cells and/or to tumor cells at a different stage of tumor progression. For example, genes are identified which are differentially expressed in benign (e.g., non-malignant) tumor cells relative to malignant, metastatic tumor cells. The modulation of the expression of the identified genes and/or the activity of the identified gene products can be utilized therapeutically to treat disorders involving tumor progression, including, for example, metastatic disorders. As such, methods and compositions are described for the identification of novel therapeutic compounds for the

inhibition of tumor progression and the treatment of tumor progression disorders, including metastatic diseases.

Further, the identified genes and/or gene products can be used to identify cells exhibiting or predisposed to a disorder involving a tumor progression phenotype, thereby diagnosing individuals having, or at high risk for developing, such disorders. Additionally, the identified genes and/or gene products can be used to grade or stage identified tumor cells. Still further, the detection of the differential expression of identified genes can be used to devise treatments (for example, chemoprevention) before the benign cells attain a malignant state. Still further, the detection of differential expression of identified genes can be used to design a preventive intervention in pre-neoplastic cells in individuals at high risk.

"Tumor progression," as used herein, refers to any event which, first, promotes the transition of a normal, non-neoplastic cell to a cancerous, neoplastic one. Such events include ones which occur prior to the onset of neoplasia, and which predispose, or act as a step toward, the cell becoming neoplastic. These events can, for example, include ones which cause a normal cell to exhibit a pre-neoplastic phenotype. Second, such events also include ones which bring about the transition from a pre-neoplastic state to a neoplastic one. Such events can, for example, include ones which promote two hallmarks of the neoplastic state, namely unhindered cell proliferation and/or tumor cell invasion of adjacent tissue. Third, tumor progression can include events which promote the transition of a tumor cell to a metastatic state. Within each state, (e.g., pre-neoplastic, neoplastic and metastatic) the term "tumor progression" as used herein can also refer to the disorder

severity or aggressiveness a cell exhibits relative to other cells within the same state.

Because multiple tumor progression events occur as a cell progresses from normal to neoplastic and metastatic states, certain cells will have undergone a different set of such tumor progression events. As such, such cells are referred to herein as belonging to different "tumor progression stages."

A "disorder involving tumor progression" or a "tumor progression disorder," as used herein, refers to the state of a cell or cells which have undergone or are in the process of undergoing a tumor progression event, as defined above.

"Differential expression," as used herein, refers to both quantitative, as well as qualitative, differences in the genes' temporal and/or cellular expression patterns among, for example, normal and neoplastic tumor cells, and/or among tumor cells which have undergone different tumor progression events. Differentially expressed genes can represent "fingerprint genes," and/or "target genes."

"Fingerprint gene," as used herein, refers to a differentially expressed gene whose expression pattern can be utilized as part of a prognostic or diagnostic marker for the evaluation of a disorder involving tumor progression, or which, alternatively, can be used in methods for identifying compounds useful for the treatment of such disorders. For example, the effect of the compound on the fingerprint gene expression normally displayed in connection with disorders involving tumor progression can be used to evaluate the efficacy of the compound as a treatment for such a disorder, or can, additionally, be used to monitor patients undergoing clinical evaluation for the treatment of the disorder.

492250 2442980

"Fingerprint pattern," as used herein, refers to the pattern generated when the expression pattern of a series (which can range from two up to all the fingerprint genes which exist for a given state) of fingerprint genes is determined. A fingerprint pattern can be used in the same diagnostic, prognostic and compound identification methods as the expression of a single fingerprint gene.

"Target gene," as used herein, refers to a differentially expressed gene involved in tumor progression such that modulation of the level of target gene expression or of target gene product activity can act to prevent and/or ameliorate symptoms of the tumor progression. Compounds that modulate the expression of the target gene or the activity of the target gene product can be used in the treatment of neoplastic diseases, including, for example, disorders involving the progression to a metastatic state. Still further, compounds that modulate the expression of the target gene or activity of the target gene product can be used in treatments to prevent benign cells from attaining a malignant state. Still further, compounds that modulate the expression of the target gene or activity of the target gene product can be used to design a preventive intervention in pre-neoplastic cells in individuals at high risk.

Further, "pathway genes" are defined via the ability of their products to interact with other gene products involved in tumor progression disorders. Pathway genes can also exhibit target gene and/or fingerprint gene characteristics.

The present invention includes the products of such fingerprint, target, and pathway genes, as well as antibodies to such gene products. Furthermore, the engineering and use of cell-based and/or animal-based models of tumor progression disorders, including disorders

involving metastasis, to which such gene products can contribute, are described.

5 The present invention also relates to methods for prognostic and diagnostic evaluation of tumor progression conditions, and for the identification of subjects containing cells predisposed to such conditions. Furthermore, the invention provides methods for evaluating the efficacy of therapies for disorders involving tumor progression, and for monitoring the progress of patients
10 participating in clinical trials for the treatment of such diseases.

The tumor progression disorders described herein can include disorders involved in the progression of such human cancers as, for example, human melanomas, breast,
15 gastrointestinal, such as esophageal, stomach, colon, bowel, colorectal and rectal cancers, prostate, bladder, testicular, ovarian, uterine, cervical, brain, lung, bronchial, larynx, pharynx, liver, pancreatic, thyroid, bone, leukemias, lymphomas, and various types of skin
20 cancers.

The invention also provides methods for the identification of compounds that modulate the expression of genes or the activity of gene products involved in tumor progression, including the progression of metastatic
25 neoplastic diseases, as well as methods for the treatment of such diseases. Such methods can, for example, involve the administration of such compounds to individuals exhibiting symptoms or markers of tumor progression, such as markers for metastatic neoplastic diseases.

30 This invention is based, in part on systematic search strategies involving in vivo and in vitro paradigms of tumor progression, including the progression to metastatic disease, coupled with sensitive and high

throughput gene expression assays, to identify genes differentially expressed in tumor cells relative to normal cells and/or relative to tumor cells at a different tumor progression stage. In contrast to approaches that merely
5 evaluate the expression of a given gene product presumed to play a role in one or another of the various stages of tumor progression, such as, for example the progression to a metastatic disease process, the search strategies and assays used herein permit the identification of all genes, whether
10 known or novel, which are differentially expressed in tumor cells relative to normal cells or relative to tumor cells at a different stage of tumor progression.

This comprehensive approach and evaluation permits the discovery of novel genes and gene products, as well as
15 the identification of an array of genes and gene products (whether novel or known) involved in novel pathways that play a major role in the disease pathology. Thus, the present invention makes possible the identification and characterization of targets useful for prognosis, diagnosis,
20 monitoring, rational drug design, and/or other therapeutic intervention of tumor progression disorders, including disorders involving metastasis.

The Example presented in Section 6, below, demonstrates the successful use of tumor progression search
25 strategies of the invention to identify genes which are differentially expressed within tumor cells relative to tumor cells at a different stage of tumor progression. Specifically, the Example identifies a gene which is differentially expressed in metastatic cell populations
30 relative to benign, non-malignant tumor cells.

This gene, referred to herein as the 030 gene (fomy030 in the mouse and fohy030 in humans), is a novel gene which is expressed at a many-fold higher level in non-

metastatic tumor cells relative to its expression in
metastatic tumor cells. The gene appears in mice and has
the cDNA sequence shown in Fig. 3A and 3B (SEQ ID NO:2). A
homologous gene, referred to herein as the fohy030 gene,
5 appears in humans and has the cDNA sequence shown in Fig. 5
(SEQ ID NO:6). An alternative splice form of the human cDNA
has the sequence shown in FIG. 6 (SEQ ID NO:8). Unless
stated expressly otherwise, any general reference to the 030
gene hereinafter refers to both the murine (fomy030) and
10 human (fohy030) homologs of this gene.

The identification of the 030 gene and the
characterization of its expression in particular stages of
metastatic spread provides, therefore, newly identified
targets for the diagnosis, prevention, and treatment of
15 tumor progression disorders, including metastatic neoplastic
diseases.

Its expression pattern indicates that the 030 gene
product acts to inhibit tumor progression. For example, a
reduction in the level of 030 gene expression correlates
20 with an increase in a cell's metastatic potential i.e., a
reduction of 030 gene product in tumor cells can induce or
predispose a cell to progress to a metastatic state.

Hence, any method which can bring about an increase
in the amount of 030 gene product can inhibit or slow the
25 progression to metastasis. In fact, it is possible that the
030 gene product exhibits general tumor inhibition
properties.

A cDNA clone of the murine homolog, designated
fomy030, is described herein in FIGS. 3A and 3B (SEQ ID
30 NO:2) (nucleotide sequence and amino acid sequence), and was
derived from fomy030 mRNA. However, as used herein, fomy030
cDNA refers to any DNA sequence that encodes the amino acid
sequence depicted in FIGS. 3A and 3B (SEQ ID NO:3).

4686550 2442980

A cDNA clone of the human homolog, designated fohy030, is shown in FIG. 5 (SEQ ID NO:6) (nucleotide sequence and amino acid sequence). An alternative splice form of fohy030 is shown in FIG. 6 (SEQ ID NO:8). Both were
5 obtained using the entire mouse fomy030 cDNA as a probe. However, as used herein, fohy030 cDNA refers to any DNA sequence that encodes the amino acid sequences depicted in FIG. 5 (SEQ ID NO:7) and FIG. 6 (SEQ ID NO:9).

3.1. DEFINITIONS

10 "Tumor progression," as used herein, refers to any event which, first, promotes the transition of a normal, non-neoplastic cell to a cancerous, neoplastic one. Such events include ones which occur prior to the onset of neoplasia, and which predispose, or act as a step toward,
15 the cell becoming neoplastic. These events can, for example, include ones which cause a normal cell to exhibit a pre-neoplastic phenotype. Second, such events also include ones which bring about the transition from a pre-neoplastic state to a neoplastic one. Such events can, for example,
20 include ones which promote unhindered cell proliferation and/or tumor cell invasion of adjacent tissue, which are viewed as hallmarks of the neoplastic state. Third, tumor progression can include events which promote the transition of a tumor cell to a metastatic state. Within each state,
25 (e.g., pre-neoplastic, neoplastic and metastatic) the term "tumor progression" as used herein can also refer to the disorder severity or aggressiveness a cell exhibits.

Because multiple tumor progression events occur as a cell progresses from a normal to neoplastic and metastatic
30 states, certain cells will have undergone a different set of such tumor progression events. As such, such cells are

referred to herein as belonging to different "tumor progression stages."

5 A "disorder involving tumor progression" or a "tumor progression disorder," as used herein, refers to the state of a cell or cells which have undergone or are in the process of undergoing a tumor progression event, as defined above.

10 "Differential expression," as used herein, refers to both quantitative, as well as qualitative differences in the genes' temporal and/or cellular expression patterns among, for example, normal and neoplastic tumor cells, and/or among tumor cells which have undergone different tumor progression events. Differentially expressed genes can represent "fingerprint genes," and/or "target genes."

15 "Fingerprint gene," as used herein, refers to a differentially expressed gene whose expression pattern can be utilized as part of a prognostic or diagnostic marker for the evaluation of tumor progression, or which, alternatively, can be used in methods for identifying
20 compounds useful for the treatment of tumor progression. For example, the effect of the compound on the fingerprint gene expression normally displayed in connection with tumor progression can be used to evaluate the efficacy of the compound as a treatment for tumor progression, or can,
25 additionally, be used to monitor patients undergoing clinical evaluation for the treatment of tumor progression.

30 "Fingerprint pattern," as used herein, refers to the pattern generated when the expression pattern of a series (which can range from two up to all the fingerprint genes which exist for a given state) of fingerprint genes is determined. A fingerprint pattern can be used in the same diagnostic, prognostic and compound identification methods as the expression of a single fingerprint gene.

268250" 21429880

"Target gene," as used herein, refers to a differentially expressed gene involved in tumor progression such that modulation of the level of target gene expression or of target gene product activity can act to prevent and/or ameliorate symptoms of the tumor progression. Compounds that modulate target gene expression or activity of the target gene product can be used in the treatment of tumor progression and tumor progression disorders, including, for example, disorders involving the progression to a metastatic state.

Further, "pathway genes" are defined via the ability of their products to interact with other gene products involved in tumor progression. Pathway genes can also exhibit target gene and/or fingerprint gene characteristics.

15 4. DESCRIPTION OF THE FIGURES

FIG. 1 is a Northern blot confirming differential regulation of the 030 gene. Total RNA (12 μ g/lane) obtained from F1 (lanes 1 and 3) and F10 (lanes 2 and 4) melanoma cell cultures was hybridized with a cDNA probe prepared by random priming of reamplified romy030 band. (See materials and methods below in Section 6.1.). The romy030 probe identifies an RNA band of approximately 3 kb, corresponding to a fomy030 mRNA.

FIG. 2 is a nucleotide sequence of romy030 band (SEQ ID NO:1).

FIGS. 3A and 3B are representations of the nucleotide and derived amino acid sequences of cDNA clone fomy030 (SEQ ID NOS:2 [nucleotide sequence] and 3 [amino acid sequence]) derived from fomy030 mRNA.

FIG. 4 is a Northern blot analysis confirming differential regulation of the fomy030 gene. Lane 1 is B16

F1, lane 2 is B16 F10, and lanes 3-6 are B16 H5, B16 H6, B16 H7 and B16 H8.

FIG. 5 is a representation of the nucleotide and deduced amino acid sequences of cDNA clone of fohy030 (SEQ ID NOS:6 [nucleotide sequence] and 7 [amino acid sequence]).

FIG. 6 is a comparison of the nucleotide and deduced amino acid sequences of another cDNA clone of fohy030 (SEQ ID NOS:8 [nucleotide sequence] and 9 [amino acid sequence]).

In FIGS. 3A and 3B, the nucleotide sequence is numbered starting at the first nucleotide, whereas in FIGS. 5 and 6, the nucleotide sequence is numbered starting at the ATG start codon.

5. DETAILED DESCRIPTION OF THE INVENTION

Methods and compositions for the prevention, treatment and diagnosis of tumor progression, including tumor progression involving metastatic disorders, in cells involved in human tumors. Such human tumors may include, for example, human melanomas, breast, gastrointestinal tumors such as esophageal, stomach, duodenal, colon, colorectal and rectal cancers, prostate, bladder, testicular, ovarian, uterine, cervical, brain, lung, bronchial, larynx, pharynx, liver, pancreatic, thyroid, bone, various types of skin cancers and other neoplastic conditions such as leukemias, lymphomas. The invention is based, in part, on the evaluation and expression and role of all genes that are differentially expressed in tumor cells relative to normal cells and/or relative to tumor cells at a different stage of tumor progression. This permits the definition of disease pathways and identification of targets in such pathways that are useful for diagnosis, prevention and treatment of tumor progression, including the tumor

progression disorders involving metastatic neoplastic diseases.

Genes, termed "target genes" and/or "fingerprint genes" are described which are differentially expressed in tumor cells relative to their expression in normal cells or relative to their expression in tumor cells which are at a different stage of tumor progression. Additionally, genes, termed "pathway genes" are described whose gene products exhibit an ability to interact with gene products involved in tumor progression, including tumor progression disorders involving metastatic neoplastic disorders. Pathway genes can additionally have fingerprint and/or target gene characteristics. Methods for the identification of such fingerprint, target, and pathway genes are also described.

Further, the gene products of such fingerprint, target, and pathway genes are described in Section 5.2.2, antibodies to such gene products are described in Section 5.2.3, as are cell-and animal-based models of tumor progression disorders to which such gene products can contribute, in Section 5.2.4.

Methods for the identification of compounds which modulate the expression of genes or the activity of gene products involved in tumor progression are described in Section 5.3. Methods for monitoring the efficacy of compounds during clinical trials are described in Section 5.3.5. Additionally described, below, are methods for treatment of tumor progression disorders, including metastatic diseases.

Also discussed, below, are methods for prognostic and diagnostic evaluation of tumor progression and disorders involving tumor progression, including metastatic disorders, and, further, for the identification of subjects exhibiting a predisposition to such disorders.

5.1. IDENTIFICATION OF DIFFERENTIALLY EXPRESSED GENES

Described herein are methods for the identification of differentially expressed genes which are involved in tumor progression. There exist a number of levels or stages at which the differential expression of such genes can be exhibited. For example, differential expression can occur in tumor cells relative to normal cells, or in tumor cells within different stages of tumor progression. For example, genes can be identified which are differentially expressed in pre-neoplastic versus neoplastic cells. Such genes can include, for example, ones which promote unhindered cell proliferation or tumor cell invasion of adjacent tissue, both of which are viewed as hallmarks of the neoplastic state. Further, differential expression can occur in benign (e.g., non-malignant) tumor cells versus metastatic, malignant tumor cells. Still further, differential expression can occur among cells within any one of these states (e.g., pre-neoplastic, neoplastic and metastatic), and can indicate, for example, a difference in tumor progression severity or aggressiveness of one cell relative to that of another cell within the same state.

Methods for the identification of such differentially expressed genes are described, below, in Section 5.1.1. Methods for the further characterization of such differentially expressed genes, and for their categorization as target and/or fingerprint genes, are presented, below, in Section 5.3.

"Differential expression" as used herein refers to both quantitative, as well as qualitative differences in the genes' temporal and/or tissue expression patterns. Thus, a differentially expressed gene can qualitatively have its expression activated or completely inactivated in, for

example, normal versus tumor progression states, in cells within different tumor progression states or among cells within a single given tumor progression state. Such a qualitatively regulated gene will exhibit an expression pattern within a given state which is detectable by standard techniques in one such state, but is not detectable in both states being compared. "Detectable," as used herein, refers to an RNA expression level which is detectable via the standard techniques of differential display, RT (reverse transcriptase)-coupled PCR, Northern and/or RNase protection analyses.

Alternatively, a differentially expressed gene can exhibit an expression level which differs, i.e., is quantitatively increased or decreased in normal versus tumor progression states, in cells within different tumor progression states or among cells within a single given tumor progression state.

The degree to which expression differs need only be large enough to be visualized via standard characterization techniques, such as, for example, the differential display technique described below. Other standard, well-known characterization techniques by which expression differences can be visualized include, but are not limited to, quantitative RT (reverse transcriptase)-coupled PCR and Northern analyses and RNase protection techniques.

Differentially expressed genes can be further described as target genes and/or fingerprint genes. "Fingerprint gene," as used herein, refers to a differentially expressed gene whose expression pattern can be utilized as part of a prognostic or diagnostic marker in tumor progression evaluation, or which, alternatively, may be used in methods for identifying compounds useful for the prevention or treatment of tumor progression and tumor

progression disorders, including metastatic disorders. A fingerprint gene can also have the characteristics of a target gene or a pathway gene (see below, in Section 5.2).

"Fingerprint pattern," as used herein, refers to the pattern generated when the expression pattern of a series (which can range from two up to all the fingerprint genes which exist for a given state) of fingerprint genes is determined. A fingerprint pattern can be used in the same diagnostic, prognostic and compound identification methods as the expression of a single fingerprint gene.

"Target gene," as used herein, refers to a differentially expressed gene involved in tumor progression in a manner by which modulation of the level of target gene expression or of target gene product activity can act to prevent and/or ameliorate symptoms of disorders involving tumor progression. Tumor progression disorders include, for example, disorders involved in human tumors, including, but not limited to human melanomas, breast, gastrointestinal, such as esophageal, stomach, colon, bowel, colorectal and rectal cancers, prostate, bladder, testicular, ovarian, uterine, cervical, brain, lung, bronchial, larynx, pharynx, liver, pancreatic, thyroid, bone, leukemias, lymphomas and various types of skin cancers. A target gene can also have the characteristics of a fingerprint gene and/or a pathway gene (as described, below, in Section 5.2).

5.1.1. METHODS FOR THE IDENTIFICATION OF DIFFERENTIALLY EXPRESSED GENES

A variety of methods can be utilized for the identification of genes which are involved in tumor progression. Described in Section 5.1.1.1 are experimental paradigms which can be utilized for the generation of samples which can be used for the identification of such

genes. Material generated in paradigm categories can be characterized for the presence of differentially expressed gene sequences as discussed, below, in Section 5.1.1.2.

5.1.1.1. PARADIGMS FOR THE IDENTIFICATION
OF DIFFERENTIALLY EXPRESSED GENES

Paradigms which represent models of tumor progression states are described herein. These paradigms can be utilized for the identification of genes which are differentially expressed in normal cells versus cells in tumor progression states, in cells within different tumor progression states or among cells within a single given tumor progression state.

The paradigms described herein include at least two groups of cells of a given cell type, preferably genetically matched cells (e.g., cells derived from variants of the same cell line, or cells derived from a single individual or biological sample), whose expression patterns are compared and analyzed for differential expression. Methods for the analysis of paradigm material are described, below, in Section 5.1.1.2.

Once a particular gene has been identified through the use of one paradigm, its expression pattern can be further characterized, for example, by studying its expression in a different paradigm. A gene can, for example, be regulated one way, i.e., can exhibit one differential gene expression pattern, in a given paradigm, but can be regulated differently in another paradigm. The use, therefore, of multiple paradigms can be helpful in distinguishing the roles and relative importance of particular genes in tumor progression.

In one embodiment of such a paradigm, referred to herein as the "in vitro" paradigm, cell lines can be used to

463350 2442980

identify genes which are differentially expressed in tumor progression states. Differentially expressed genes are detected, as described herein, by comparing the pattern of gene expression between the experimental and control

5 conditions. In such a paradigm, genetically matched tumor cell lines (e.g., variants of the same cell line) are generally utilized. For example, the gene expression pattern of two variant cell lines can be compared, wherein one variant exhibits characteristics of one tumor progression

10 state while the other variant exhibits characteristics of another tumor progression state. Alternatively, two variant cell lines, both of which exhibit characteristics of the same tumor progression state, but which exhibit differing degrees of tumor progression disorder severity or

15 aggressiveness. Further, genetically matched cell lines can be utilized, one of which exhibits characteristics of a tumor progression state, while the other exhibits a normal cellular phenotype.

The variant cell lines utilized herein can exhibit

20 such tumor progression characteristics as, for example, a high or low metastatic potential, which refers to the likelihood that a cell will give rise to a distant site tumor mass. Alternatively, one or more such variant cell lines can exhibit pre-neoplastic characteristics or can

25 exhibit characteristics generally associated with one or more neoplastic cell phenotypes, such as, for example, cell proliferation or invasion phenotypes.

In accordance with this aspect of the invention, the cell line variants are cultured under appropriate

30 conditions, the cells are harvested, and RNA is isolated and analyzed for differentially expressed genes, as described in detail in Section 5.1.1.2, below.

46550 24429880

Examples of cell lines that can be used as part of such in vitro paradigms include but are not limited to variants of melanoma cell lines, such as, for example, the murine melanoma B16 F1 cell line which exhibits a low metastatic potential and the melanoma B16 F10 cell line which exhibits a high metastatic potential (Fidler, I. J., 1973, Nature New Biol 242:148-149); human colon cell lines, such as, for example KM12c (tumor cell line with low metastatic potential) and the KM20L4 (tumor cell line with high metastatic potential; Morikawa K., et al., 1988, Cancer Research 48:1943-1948); prostatic tumor cell lines, such as, for example, DU 145 (non metastatic tumor cell line) and PC-3-M (high metastatic potential tumor cell line; Karmali, R.A. et al., 1987, Anticancer Res. 7:1173-1180, and Koziowski, J.M. et al., 1984, Cancer Research 44:3522-3529); and breast carcinoma tumor cell lines, such as, for example, MCF-7 (non metastatic tumor cell line) and MDA-MB-435 (high metastatic potential tumor cell line; Watts C.K. et al., 1994, Breast Cancer Res. Treat. 31:95-105 and Rose, D.P. et al., 1993, J. Natl. Cancer Inst. 85:1743-1747).

As presented in the Example presented in Section 6, below, this paradigm has been successfully utilized to identify a gene, referred to herein as the 030 gene, which is differentially expressed in cells exhibiting a high metastatic potential relative to cells exhibiting a low metastatic potential. Specifically, the 030 gene is expressed at a many-fold higher level in low metastatic potential cells relative to cells exhibiting a high metastatic potential.

In a second paradigm, referred to herein as the in vivo paradigm, animal models of tumor progression disorders can be utilized to discover differentially expressed gene sequences. The in vivo nature of such tumor progression

models can prove to be especially predictive of the analogous responses in living patients.

A variety of tumor progression animal models can be used for as part of the in vivo paradigms. For example, animal models of tumor progression may be generated by passaging tumor cells in animals (e.g., mice), leading to the appearance of tumors within these animals.

Additional animal models, some of which may exhibit differing tumor progression characteristics, may be generated from the original animal models described above. For example, the tumors which result in the original animals can be removed and grown in vitro. Cells from these in vitro cultures can then be passaged in animals and tumors resulting from this passage can then be isolated. RNA from pre-passage cells, and cells isolated after one or more rounds of passage can then be isolated and analyzed for differential expression. The differential expression can be compared to the metastatic potential expression of such cells. These cells can now represent cells from different tumor progression states, or cells within a given tumor progression state exhibiting differing degrees of severity or aggressiveness. Such passaging techniques can utilizing any of the variant cell lines described, above, for the in vitro paradigms.

Additionally, animal models for tumor progression which can be utilized for such an in vivo paradigm include any of the animal models described, below, in Section 5.7.1. Other models include transgenic mouse model for melanoma (Mintz, B. and Silvers, W.K., 1993, Proc. Natl. Acad. Sci. USA 90:8817-8812), transgenic mice which carry specific adenomatous polyposis coli (APC) gene mutations (Fodde, R., et al., 1994, Proc. Natl. Acad. Sci. USA 91:8969-8973) and the transgenic mouse in which the mammary tumor virus LTR/c-

myc gene is anomalously expressed (Leder, A., et al., 1986, Cells 45:485-495).

A third paradigm, referred to herein as the "specimen paradigm," utilizes samples from surgical and biopsy specimens. Such specimens can represent normal tissue, primary, secondary or metastasized tumors obtained from patients having undergone surgical treatment for disorders involving tumor progression such as, for example, melanomas, colon carcinomas, lung carcinomas, prostatic cancers and breast cancers.

Surgical specimens can be procured under standard conditions involving freezing and storing in liquid nitrogen (see, for example, Karmali, R.A., et al., 1983, Br. J. cancer 48:689-696.) RNA from specimen cells is isolated by, for example, differential centrifugation of homogenized tissue, and analyzed for differential expression relative to other specimen cells, preferably cells obtained from the same patient.

In paradigms designed to identify genes which are involved in tumor progression, compounds known to have an ameliorative effect on the tumor progression symptoms can also be used in paradigms to detect differentially expressed genes. Such compounds can include known therapeutics, as well as compounds that are not useful as therapeutics due to their harmful side effects. For example, tumor cells that are cultured as explained in this Section, above, can be exposed to one of these compounds and analyzed for differential gene expression with respect to untreated tumor cells, according to the methods described below in Section 5.1.1.2. In principle, however, according to the paradigm, any cell type involved in tumor progression and disorders thereof can be treated by these compounds at any stage of the tumor progression process.

Cells involved in tumor progression can also be compared to unrelated cells (e.g., fibroblasts) which have been treated with the compound, such that any generic effects on gene expression that might not be related to the disease or its treatment may be identified. Such generic effects might be manifest, for example, by changes in gene expression that are common to the test cells and the unrelated cells upon treatment with the compound.

By these methods, the genes and gene products upon which these compounds act can be identified and used in the assays described below to identify novel therapeutic compounds for inhibition of tumor progression and the treatment of tumor progression disorders, including metastatic diseases.

5.1.1.2. ANALYSIS OF PARADIGM MATERIAL

In order to identify differentially expressed genes, RNA, either total or mRNA, can be isolated from cells utilized in paradigms such as those described earlier in Section 5.1.1.1. Any RNA isolation technique which does not select against the isolation of mRNA can be utilized for the purification of such RNA samples. See, for example, Ausubel, F.M. et al., eds., 1987-1993, Current Protocols in Molecular Biology, John Wiley & Sons, Inc. New York, which is incorporated herein by reference in its entirety.

Additionally, large numbers of tissue samples can readily be processed using techniques well known to those of skill in the art, such as, for example, the single-step RNA isolation process of Chomczynski, P. (1989, U.S. Patent No. 4,843,155), which is incorporated herein by reference in its entirety.

Transcripts within the collected RNA samples which represent RNA produced by differentially expressed genes can

be identified by utilizing a variety of methods which are well known to those of skill in the art. For example, differential screening (Tedder, T.F. et al., 1988, Proc. Natl. Acad. Sci. USA 85:208-212), subtractive hybridization
5 (Hedrick, S.M. et al., 1984, Nature 308:149-153; Lee, S.W. et al., 1984, Proc. Natl. Acad. Sci. USA 88:2825), and, preferably, differential display (Liang, P. and Pardee, A.B., 1993, U.S. Patent No. 5,262,311, which is incorporated herein by reference in its entirety), can be utilized to
10 identify nucleic acid sequences derived from genes that are differentially expressed.

Differential screening involves the duplicate screening of a cDNA library in which one copy of the library is screened with a total cell cDNA probe corresponding to
15 the mRNA population of one cell type while a duplicate copy of the cDNA library is screened with a total cDNA probe corresponding to the mRNA population of a second cell type. For example, one cDNA probe can correspond to a total cell cDNA probe of a cell type or tissue derived from a control
20 subject, while the second cDNA probe can correspond to a total cell cDNA probe of the same cell type derived from an experimental subject. Those clones which hybridize to one probe but not to the other potentially represent clones derived from genes differentially expressed in the cell type
25 of interest in control versus experimental subjects.

Subtractive hybridization techniques generally involve the isolation of mRNA taken from two different sources, e.g., control and experimental tissue, the hybridization of the mRNA or single-stranded cDNA reverse-
30 transcribed from the isolated mRNA, and the removal of all hybridized, and therefore double-stranded, sequences. The remaining non-hybridized, single-stranded cDNAs, potentially represent clones derived from genes that are differentially

expressed in the two mRNA sources. Such single-stranded cDNAs are then used as the starting material for the construction of a library comprising clones derived from differentially expressed genes.

5 The differential display technique describes a procedure, utilizing the well-known polymerase chain reaction (PCR; the experimental embodiment set forth in Mullis, K.B., 1987, U.S. Patent No. 4,683,202) which allows for the identification of sequences derived from genes which
10 are differentially expressed. First, isolated RNA is reverse-transcribed into single-stranded cDNA, utilizing standard techniques which are well known to those of skill in the art. Primers for the reverse transcriptase reaction can include, but are not limited to, oligo dT-containing
15 primers, preferably of the 3' primer type of oligonucleotide described below. Next, this technique uses pairs of PCR primers, as described below, which allow for the amplification of clones representing a random subset of the RNA transcripts present within any given cell. Utilizing
20 different pairs of primers allows each of the mRNA transcripts present in a cell to be amplified. Among such amplified transcripts can be identified those which have been produced from differentially expressed genes.

 The 3' oligonucleotide primer of the primer pairs
25 can contain an oligo dT stretch of 10-13 dT nucleotides at its 5' end, preferably 11, which hybridizes to the poly(A) tail of mRNA or to the complement of a cDNA reverse transcribed from an mRNA poly(A) tail. Second, in order to increase the specificity of the 3' primer, the primer can
30 contain one or more, preferably two, additional nucleotides at its 3' end. Because, statistically, only a subset of the mRNA derived sequences present in the sample of interest will hybridize to such primers, the additional nucleotides

allow the primers to amplify only a subset of the mRNA derived sequences present in the sample of interest. This is preferred in that it allows more accurate and complete visualization and characterization of each of the bands representing amplified sequences.

The 5' primer can contain a nucleotide sequence expected, statistically, to have the ability to hybridize to cDNA sequences derived from the tissues of interest. The nucleotide sequence can be an arbitrary one, and the length of the 5' oligonucleotide primer can range from about 9 to about 15 nucleotides, with about 13 nucleotides being preferred.

Additionally, arbitrary primer sequences cause the lengths of the amplified partial cDNAs produced to be variable, thus allowing different clones to be separated by using standard denaturing sequencing gel electrophoresis.

PCR reaction conditions should be chosen which optimize amplified product yield and specificity, and, additionally, produce amplified products of lengths which can be resolved utilizing standard gel electrophoresis techniques. Such reaction conditions are well known to those of skill in the art, and important reaction parameters include, for example, length and nucleotide sequence of oligonucleotide primers as discussed above, and annealing and elongation step temperatures and reaction times.

The pattern of clones resulting from the reverse transcription and amplification of the mRNA of two different cell types is displayed via sequencing gel electrophoresis and compared. Differences in the two banding patterns indicate potentially differentially expressed genes.

Once potentially differentially expressed gene sequences have been identified via bulk techniques such as, for example, those described above, the differential

expression of such putatively differentially expressed genes should be corroborated. Corroboration can be accomplished via, for example, such well-known techniques as Northern analysis, quantitative RT-coupled PCR or RNase protection.

5 Upon corroboration, the differentially expressed genes can be further characterized, and can be identified as target and/or fingerprint genes, as discussed, below, in Section 5.1.4.

10 Also, amplified sequences of differentially expressed genes obtained through differential display can be used to isolate the full length clones of the corresponding gene. The full-length coding portion of the gene can readily be isolated, without undue experimentation, by molecular biological techniques well known in the art. For
15 example, the isolated differentially expressed amplified fragment can be labeled and used to screen a cDNA library. Alternatively, the labeled fragment can be used to screen a genomic library.

20 PCR technology can also be utilized to isolate full-length cDNA sequences. As described in this section above, the isolated amplified gene fragments (of about at least 10 nucleotides, preferably longer, of about 15 nucleotides) obtained through differential display have their 5' terminal end at some random point within the gene and have 3'
25 terminal ends at a position corresponding to the 3' end of the transcribed portion of the gene. Once nucleotide sequence information from an amplified fragment is obtained, the remainder of the gene (i.e., the 5' end of the gene, when utilizing differential display) can be obtained
30 using, for example, RT PCR.

 In one embodiment of such a procedure for the identification and cloning of full length gene sequences,

RNA can be isolated, following standard procedures, from an appropriate tissue or cellular source.

A reverse transcription reaction can then be performed on the RNA using an oligonucleotide primer

5 complementary to the mRNA that corresponds to the amplified cloned fragment, for the priming of first strand synthesis. Because the primer is anti-parallel to the mRNA, extension will proceed toward the 5' end of the mRNA. The resulting RNA/DNA hybrid can then be "tailed" with guanines using a
10 standard terminal transferase reaction, the hybrid can be digested with RNAase H, and second strand synthesis can then be primed with a poly-C primer. Using the two primers, the 5' portion of the gene is then amplified using PCR.

Sequences obtained can then be isolated and recombined with
15 previously isolated sequences to generate a full-length cDNA of the differentially expressed genes of the invention. For a review of cloning strategies and recombinant DNA techniques which can be used, see, e.g., Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, Cold Springs
20 Harbor Press, N.Y.; and Ausubel et al., 1989, Current Protocols in Molecular Biology, (Green Publishing Associates and Wiley Interscience, N.Y.).

5.2. METHODS FOR THE IDENTIFICATION OF PATHWAY GENES

25 Methods are described herein for the identification of pathway genes. "Pathway gene," as used herein, refers to a gene whose gene product exhibits the ability to interact with gene products involved in tumor progression. A pathway gene can be differentially expressed and, therefore, can
30 have the characteristics of a target and/or fingerprint gene.

Any method suitable for detecting protein-protein interactions can be employed for identifying pathway gene products by identifying interactions between gene products and gene products known to be involved in tumor progression and tumor progression disorders, including metastatic disorders. Such known gene products can be cellular or extracellular proteins. Those gene products which interact with such known gene products represent pathway gene products and the genes which encode them represent pathway genes.

Among the traditional methods which can be employed are co-immunoprecipitation, cross-linking and co-purification through gradients or chromatographic columns. Utilizing procedures such as these allows for the identification of pathway gene products. Once identified, a pathway gene product can be used, in conjunction with standard techniques, to identify its corresponding pathway gene. For example, at least a portion of the amino acid sequence of the pathway gene product can be ascertained using techniques well known to those of skill in the art, such as via the Edman degradation technique (see, e.g., Creighton, 1983, Proteins: Structures and Molecular Principles, W.H. Freeman & Co., N.Y., pp.34-49). The amino acid sequence obtained can be used as a guide for the generation of oligonucleotide mixtures that can be used to screen for pathway gene sequences. Screening can be accomplished, for example by standard hybridization or PCR techniques. Techniques for the generation of oligonucleotide mixtures and the screening are well known. (See, e.g., Ausubel, supra., and PCR Protocols: A Guide to Methods and Applications, 1990, Innis, M. et al., eds. Academic Press, Inc., New York).

268050" 24429830
Additionally, methods can be employed which result in the simultaneous identification of pathway genes which encode the protein interacting with a protein involved in tumor progression and tumor progression disorders, including
5 metastatic diseases. These methods include, for example, probing expression libraries with labeled protein known or suggested to be involved in metastatic diseases using this protein in a manner similar to the well known technique of antibody probing of λ gt11 libraries.

10 One method which detects protein interactions in vivo, the yeast two-hybrid system, is described in detail for illustration only and not by way of limitation. One version of this system has been described (Chien et al., 1991, Proc. Natl. Acad. Sci. USA, 88:9578-9582) and is
15 commercially available from Clontech (Palo Alto, CA).

Briefly, utilizing such a system, plasmids are constructed that encode two hybrid proteins: the first hybrid protein consists of the DNA-binding domain of a transcription factor (e.g., activation protein) fused to a
20 known protein, in this case, a protein known to be involved in tumor progression, and the second hybrid protein consists of the transcription factor's activation domain fused to an unknown protein that is encoded by a cDNA which has been recombined into this plasmid as part of a cDNA library. The
25 plasmids are transformed into a strain of the yeast *Saccharomyces cerevisiae* that contains a reporter gene (e.g., lacZ) whose expression is regulated by the transcription factor's binding site. Either hybrid protein alone cannot activate transcription of the reporter gene.
30 The DNA binding hybrid protein cannot activate transcription because it does not provide the activation domain function and the activation domain hybrid protein cannot activate transcription because it lacks the domain required for

binding to its target site (e.g., it cannot localize to the transcription activator protein's binding site).

Interaction between the DNA binding hybrid protein and the library encoded protein reconstitutes the functional
5 transcription factor and results in expression of the reporter gene, which is detected by an assay for the reporter gene product.

The two-hybrid system or related methodology can be used to screen activation domain libraries for proteins that
10 interact with a known "bait" gene product. By way of example, and not by way of limitation, gene products (e.g., 030 gene products) known to be involved in tumor progression and tumor progression disorders, such as metastatic diseases, can be used as the bait gene products. Total
15 genomic or cDNA sequences are fused to the DNA encoding an activation domain. This library and a plasmid encoding a hybrid of the bait gene product fused to the DNA-binding domain are cotransformed into a yeast reporter strain, and the resulting transformants are screened for those that
20 express the reporter gene. For example, and not by way of limitation, the bait gene can be cloned into a vector such that it is translationally fused to the DNA encoding the DNA-binding domain of the GAL4 protein. The colonies are purified and the (library) plasmids responsible for reporter
25 gene expression are isolated. The inserts in the plasmids are sequenced to identify the proteins encoded by the cDNA or genomic DNA.

A cDNA library of a cell or tissue source which expresses proteins predicted to interact with the bait gene
30 product can be made using methods routinely practiced in the art. According to the particular system described herein, the library is generated by inserting the cDNA fragments into a vector such that they are translationally fused to

the activation domain of GAL4. This library can be co-transformed along with the bait gene-GAL4 fusion plasmid into a yeast strain which contains a *lacZ* gene whose expression is controlled by a promoter which contains a GAL4 activation sequence. A cDNA encoded protein, fused to GAL4 activation domain, that interacts with the bait gene product will reconstitute an active GAL4 transcription factor and thereby drive expression of the *lacZ* gene. Colonies which express *lacZ* can be detected by their blue color in the presence of X-gal. cDNA containing plasmids from such a blue colony can then be purified and used to produce and isolate the bait gene product interacting protein using techniques routinely practiced in the art.

Once a pathway gene has been identified and isolated, it can be further characterized as, for example, discussed below, in Section 5.3.

5.3. CHARACTERIZATION OF DIFFERENTIALLY EXPRESSED AND PATHWAY GENES

Differentially expressed genes, such as those identified via the methods discussed, above, in Section 5.1, and pathway genes, such as those identified via the methods discussed, above, in Section 5.2, above, as well as genes identified by alternative means, can be further characterized by utilizing, for example, methods such as those discussed herein. Such genes will be referred to herein as "identified genes."

Analyses such as those described herein, yield information regarding the biological function of the identified genes. An assessment of the biological function of the differentially expressed genes, in addition, will allow for their designation as target and/or fingerprint genes.

Specifically, any of the differentially expressed genes whose further characterization indicates that a modulation of the gene's expression or a modulation of the gene product's activity can inhibit tumor progression will be designated "target genes," as defined, above, in Section 5.1. Such target genes and target gene products, along with those discussed below, will constitute the focus of the compound discovery strategies discussed, below, in Section 5.8. Further, such target genes, target gene products and/or modulating compounds can be used as part of the tumor progression disorder treatment methods described, below, in Section 5.9.

Any of the differentially expressed genes whose further characterization indicates that such modulations does not positively affect tumor progression, but whose expression pattern contributes to a gene expression "fingerprint" pattern correlative of, for example, tumor progression will be designated a "fingerprint gene." "Fingerprint patterns" will be more fully discussed, below, in Section 5.11.1. It should be noted that each of the target genes can also function as fingerprint genes, as can all or a portion of the pathway genes.

It should further be noted that the pathway genes can also be characterized according to techniques such as those described herein. Those pathway genes which yield information indicating that they are differentially expressed and that modulation of the gene's expression or a modulation of the gene product's activity can inhibit tumor progression or ameliorate tumor progression-associated symptoms will also be designated "target genes." Such target genes and target gene products, along with those discussed above, will constitute the focus of the compound discovery strategies discussed, below, in Section 5.8 and

can be used as part of the treatment methods described in Section 5.9, below.

It should be additionally noted that the characterization of one or more of the pathway genes can reveal a lack of differential expression, but evidence that modulation of the gene's activity or expression can, nonetheless, ameliorate symptoms of tumor progression. In such cases, these genes and gene products would also be considered a focus of the compound discovery strategies of Section 5.8, below and can be used as part of the treatment methods described in Section 5.9, below.

In instances wherein a pathway gene's characterization indicates that modulation of gene expression or gene product activity cannot retard the tumor progression diseases of interest, but is differentially expressed and contributes to a gene expression fingerprint pattern correlative of, tumor progression states or disorders, such as metastatic diseases, such pathway genes can additionally be designated as fingerprint genes.

A variety of techniques can be utilized to further characterize the identified genes. First, the nucleotide sequence of the identified genes, which can be obtained by utilizing standard techniques well known to those of skill in the art, can be used to further characterize such genes. For example, the sequence of the identified genes can reveal homologies to one or more known sequence motifs which can yield information regarding the biological function of the identified gene product.

Second, an analysis of the tissue and/or cell type distribution of the mRNA produced by the identified genes can be conducted, utilizing standard techniques well known to those of skill in the art. Such techniques can include, for example, Northern analyses, RT-coupled PCR and RNase

protection techniques. Such analyses provide information as to whether the identified genes are expressed in tissues expected to contribute to tumor progression. Such analyses can also provide quantitative information regarding steady state mRNA regulation, yielding data concerning which of the identified genes exhibits a high level of regulation in, preferably, tissues which can be expected to contribute to tumor progression. Additionally, standard in situ hybridization techniques can be utilized to provide information regarding which cells within a given tissue express the identified gene. Such an analysis can provide information regarding the biological function of an identified gene relative to given tumor progression in instances wherein only a subset of the cells within the tissue is thought to be relevant to the disorder.

Third, the sequences of the identified genes can be used, utilizing standard techniques, to place the genes onto genetic maps, e.g., mouse (Copeland, N.G. and Jenkins, N.A., 1991, Trends in Genetics 7:113-118) and human genetic maps (Cohen, D., et al., 1993, Nature 366:698-701). Such mapping information can yield information regarding the genes' importance to human disease by, for example, identifying genes which map within genetic regions to which known genetic tumor progression disorders map.

Fourth, the biological function of the identified genes can be more directly assessed by utilizing relevant in vivo and in vitro systems. In vivo systems can include, but are not limited to, animal systems which naturally exhibit symptoms of tumor progression, such as metastatic disease, or ones which have been engineered to exhibit such symptoms. For example, tumor progression animal models may be generated by injecting animals, such as mice, with tumor cells, some of which will give rise to tumors within the

injected animals. Among the cells which may be utilized for such a purpose are cells listed, above, in Section 5.1.1.1, such as the B16 cell variants.

5 The role of identified gene products (e.g., 030 gene products) can be determined by transfecting cDNAs encoding these gene products into appropriate cell lines, such as, for example, a B16 cell line variant, and analyzing the effect on tumor progression characteristics. For example, the role/function of genes important in the progression of
10 human colorectal cancers are assessed using the KM12c (low metastatic potential) and KM12L4 (highly metastatic) cells implanted into nude mice spleens and the number of hepatic tumors that develop are determined. The function of genes isolated using human colorectal tumors and their hepatic
15 metastases are assessed by expressing the gene in the appropriate KM12 variant. Additionally, the role/function of genes important in the progression of prostatic and breast cancers are assessed using appropriate cell lines described above in Section 5.1.1.1. Importantly, the
20 role/function of genes important in the progression of melanoma, colon, prostate and breast cancers in humans are assessed using biopsy specimens from patients having undergone surgical treatment, as described in Section 5.1.1.1. above.

25 Further, such systems can include, but are not limited to transgenic animal systems such as those described, above, in Section 5.7.1 below. In vitro systems can include, but are not limited to, cell-based systems comprising cell types known or suspected of contributing to
30 tumor progression. Such cells can be wild type cells, or can be non-wild type cells containing modifications known to or suspected of, contributing to tumor progression. Such systems are discussed in detail, below, in Section 5.7.2.

25 4429930
The procedure to identify and isolate the human homologue of the fomy030 gene is described, below, in Section 5.7.3.

5 In further characterizing the biological function of the identified genes, the expression of these genes can be modulated within the in vivo and/or in vitro systems, i.e., either over- or under-expressed, and the subsequent effect on the system then assayed. Alternatively, the activity of the product of the identified gene can be modulated by either increasing or decreasing the level of activity in the
10 in vivo and/or in vitro system of interest, and its subsequent effect then assayed.

The information obtained through such characterizations can suggest relevant methods for the treatment of tumor progression and tumor progression
15 disorders involving the gene of interest. Further, relevant methods for controlling the spread of tumor cells involving the gene of interest can be suggested by information obtained from such characterization. For example, treatment can include a modulation of gene expression and/or gene
20 product activity. Characterization procedures such as those described herein can indicate where such modulation should involve an increase or a decrease in the expression or activity of the gene or gene product of interest. Such methods of treatment are discussed, below, in Section 5.9.

25 **5.4. DIFFERENTIALLY EXPRESSED
AND PATHWAY GENES**

Differentially expressed genes, such as those identified in Section 5.1.1, above, and pathway genes, such as those identified in Section 5.2, above, are described
30 herein.

The differentially expressed and pathway genes of the invention are listed below, in Table 1. The nucleotide

sequence for the differentially expressed fomy030 gene is shown in FIGS. 2 and 3A and 3B. Specifically, FIG. 2 depicts the nucleotide sequence (SEQ ID NO:1) of the amplified cDNA band initially identified via differential display analysis, which is referred to herein as romy030. FIGS. 3A AND 3B depict the nucleotide sequence (SEQ ID NO:2) of a fomy030 cDNA clone which was isolated using a romy030 probe. The deduced amino acid sequence also is shown in FIGS. 3A and 3B (SEQ ID NO:3). FIG. 5 shows the nucleotide (SEQ ID NO:6) and deduced amino acid sequences (SEQ ID NO:7) of a fohy030 cDNA clone which was isolated using the entire mouse fomy030 cDNA as a probe. FIG. 6 shows an alternative splice form of fohy030 (SEQ ID NOs:8 and 9).

Table 1 summarizes information regarding the further characterization of the differentially expressed fomy030 gene of the invention. Table 2 lists E. coli clones, deposited with the Agricultural Research Service Culture Collection (NRRL), which contain sequences found within the genes of Table 1.

In Table 1, the paradigm used initially to detect the differentially expressed gene is described under the column headed "Paradigm of Original Detection." In this column, "↑" indicates that gene expression is higher (*i.e.*, there is a greater steady state amount of detectable mRNA produced by a given gene) in the indicated cell type relative to the other cell type, while "↓" indicates that gene expression is lower (*i.e.*, there is a lower steady state amount of detectable mRNA, produced by a given gene) in the indicated cell type relative to the other cell type. As indicated under this column, the 030 gene was initially identified via a differential screen between B16 F1 (low metastatic potential cells) and B16 F10 (high metastatic potential cells) in which 030 gene expression is lower in

the high metastatic potential B16 F10 cell line than in the low metastatic potential B16 F1 cell line.

5 The Table 1 column headed "Paradigm Expression Pattern" lists the cell type in which gene expression was initially detected. In the case of the 030 gene, gene expression was first detected in melanoma (i.e., B16) cells. "Detectable" as used herein, refers to levels of mRNA which are detectable, via standard differential display, Northern, RT-coupled PCR and/or RNase protection techniques which are
10 well known to those of skill in the art.

Cell types in which differential expression was detected are summarized in Table 1 under the column headed "Cell Type Detected in." In the case of the 030 gene, expression has additionally been detected within melanocyte
15 cells.

Additionally, in instances wherein the genes contain nucleotide sequences similar or homologous to sequences found in nucleic acid databases, references to such similarities are listed. Because the 030 gene is a novel
20 gene, i.e., no homologous gene sequences are present in the published databases, no such reference is listed.

Finally, nucleotide sequences contained within the differentially expressed genes are listed in the Figures indicated under the heading "Seq." In the case of the
25 fomy030 gene, such sequences are listed in FIGS. 2 and 3A and 3B, and for fohy030, in FIGS. 5 and 6.

The genes listed in Table 1 can be obtained using cloning methods well known to those skilled in the art, including, but not limited to, the use of appropriate probes
30 to detect the genes within an appropriate cDNA or gDNA (genomic DNA) library. (See, for example, Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratories, which is incorporated by reference

herein in its entirety). Probes for the novel sequences reported herein can be obtained directly from the isolated clones deposited with the NRRL, as indicated in Table 2, below. Alternatively, oligonucleotide probes for the novel genes can be synthesized, using techniques well known to those of skill in the art, based on the DNA sequences disclosed herein in FIGS. 2, 3A, 3B, 5, and 6.

The probes can be used to screen cDNA libraries prepared from an appropriate cell or cell line in which the gene is transcribed. For example, the genes described herein that were detected in melanocyte cells can be cloned from a cDNA library prepared from melanocyte cells such as, for example, melan-c (Hodgkinson, C.A., et al., 1993, Cell 74:395-404), the cDNA libraries developed from the human melanoma cell line A2058 (Clontech, Palo Alto, CA) and cDNA libraries developed from the murine melanoma cell line K1735 (Stratagene, La Jolla, CA). Genomic DNA libraries can be prepared from any source.

TABLE 1
Differentially Expressed and Pathway Genes

GENE	Sequence ID	Paradigm of Original Detection (↑/↓)	Paradigm Expression Pattern	Cell Type Detected in	Ref.	Seq.
fomy030	2	B16 ↑ F1 B16 ↓ F10	melanoma cells	melanocyte		Fig. 2, 3A & 3B
fohy030	6 & 8	benign nevi ↑ malignant melanoma ↓	biopsy samples	melanocyte		Fig. 5 & 6

Table 2, below, lists an E. coli strain as deposited with the NRRL, which contains an isolated plasmid fomy030 clone. The clone contains a fomy030 cDNA in a pBlueScript SK- (Stratagene, La Jolla, CA) vector which was isolated from a mouse melanocyte cDNA library screened with a romy030 probe, as described in Section 6.2, below.

TABLE 2

GENE	STRAIN DEPOSITED WITH NRRL	PLASMID CLONE CONTAINED WITHIN DEPOSITED STRAIN
fomy030	FOMY030	pFOMY030
fohy030		

As used herein, "differentially expressed gene" (*i.e.*, target and fingerprint genes) or "pathway gene" refers to (a) a gene containing: at least one of the DNA sequences disclosed herein (as shown in FIGS. 2, 3A, 3B, 5, and 6) or contained in the clones listed in Table 2, as deposited with the NRRL; (b) any DNA sequence that encodes the amino acid sequence encoded by: the DNA sequences disclosed herein (as shown in FIGS. 2, 3A, 3B, 5, and 6), contained in the clones, listed in Table 2, as deposited with the NRRL or contained within the coding region of the gene to which the DNA sequences disclosed herein (as shown in FIGS. 2, 3A, 3B, 5, and 6) or contained in the clones listed in Table 2, as deposited with the NRRL, belong; (c) any DNA sequence that hybridizes to the complement of: the coding sequences disclosed herein (as shown in FIGS 2, 3A, 3B, 5, and 6), contained in clones listed in Table 2, as deposited with the NRRL, or contained within the coding region of the gene to which the DNA sequences disclosed

herein (as shown in FIGS. 2, 3A, 3B, 5, and 6) or contained in the clones listed in Table 2, as deposited with the NRRL, belong under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1xSSC/0.1% SDS at 68°C (Ausubel F.M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at p. 2.10.3) and encodes a gene product functionally equivalent to a gene product encoded by a gene of (a), above and/or (d) any DNA sequence that hybridizes to the complement of: the coding sequences disclosed herein, (as shown in FIGS. 2, 3A, 3B, 5, and 6) contained in the clones listed in Table 2, as deposited with the NRRL or contained within the coding region of the gene to which DNA sequences disclosed herein (as shown in FIGS. 2, 3A, 3B, 5, and 6) or contained in the clones, listed in Table 2, as deposited with the NRRL, belong under less stringent conditions, such as moderately stringent conditions, e.g., washing in 0.2xSSC/0.1% SDS at 42°C (Ausubel et al., 1989, supra), yet which still encodes a gene product functionally equivalent to a gene product encoded by a gene of (a), above.

The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore the complements of, the DNA sequences (a) through (d), in the preceding paragraph. Such hybridization conditions can be highly stringent or less highly stringent, as described above. In instances wherein the nucleic acid molecules are deoxyoligonucleotides ("oligos"), highly stringent conditions can refer, e.g., to washing in 6xSSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base

oligos), and 60°C (for 23-base oligos). These nucleic acid molecules can act as target gene antisense molecules, useful, for example, in target gene regulation and/or as antisense primers in amplification reactions of target, fingerprint, and/or pathway gene nucleic acid sequences. Further, such sequences can be used as part of ribozyme and/or triple helix sequences, also useful for target gene regulation. Still further, such molecules can be used as components of diagnostic methods whereby tumor progression disorders can be detected.

The invention also encompasses (a) DNA vectors that contain any of the foregoing coding sequences and/or their complements (i.e., antisense); (b) DNA expression vectors that contain any of the foregoing coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences; and (c) genetically engineered host cells that contain any of the foregoing coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell. As used herein, regulatory elements include but are not limited to inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression. The invention includes fragments of any of the DNA sequences disclosed herein.

In addition to the gene sequences described above, homologues of these gene sequences as can, for example be present in other species, preferably human in instances wherein the above described gene sequences are not human gene sequences, can be identified and can readily be isolated, without undue experimentation, by molecular biological techniques well known in the art. Further, there can exist genes at other genetic loci within the genome that

encode proteins which have extensive homology to one or more domains of such gene products. These genes can also be identified via similar techniques.

For example, the isolated differentially expressed
5 gene sequence can be labeled and used to screen a cDNA
library constructed from mRNA obtained from the organism of
interest. Hybridization conditions will be of a lower
stringency when the cDNA library was derived from an
10 organism different from the type of organism from which the
labeled sequence was derived. Alternatively, the labeled
fragment can be used to screen a genomic library derived
from the organism of interest, again, using appropriately
stringent conditions. Such low stringency conditions will
be well known to those of skill in the art, and will vary
15 predictably depending on the specific organisms from which
the library and the labeled sequences are derived. For
guidance regarding such conditions see, for example,
Sambrook et al., 1989, Molecular Cloning, A Laboratory
Manual, Cold Springs Harbor Press, N.Y.; and Ausubel et al.,
20 1989, Current Protocols in Molecular Biology, (Green
Publishing Associates and Wiley Interscience, N.Y.).

Further, a previously unknown differentially
expressed or pathway gene-type sequence can be isolated by
performing PCR using two degenerate oligonucleotide primer
25 pools designed on the basis of amino acid sequences within
the gene of interest. The template for the reaction can be
cDNA obtained by reverse transcription of mRNA prepared from
human or non-human cell lines or tissue known or suspected
to express a differentially expressed or pathway gene
30 allele. The PCR product can be subcloned and sequenced to
insure that the amplified sequences represent the sequences
of a differentially expressed or pathway gene-like nucleic
acid sequence.

26350" 24429880

The PCR fragment can then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment can be labeled and used to screen a bacteriophage cDNA library. Alternatively, the labeled
5 fragment can be used to screen a genomic library.

PCR technology can also be utilized to isolate full length cDNA sequences. For example, RNA can be isolated, following standard procedures, from an appropriate cellular or tissue source. A reverse transcription reaction can be
10 performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid can then be "tailed" with guanines using a standard terminal transferase reaction, the hybrid can be
15 digested with RNAase H, and second strand synthesis can then be primed with a poly-C primer. Thus, cDNA sequences upstream of the amplified fragment can easily be isolated. For a review of cloning strategies which can be used, see e.g., Sambrook et al., 1989, Molecular Cloning, A Laboratory
20 Manual, Cold Springs Harbor Press, N.Y.; and Ausubel et al., 1989, Current Protocols in Molecular Biology, (Green Publishing Associates and Wiley Interscience, N.Y.).

In cases where the differentially expressed or pathway gene identified is the normal, or wild type, gene,
25 this gene can be used to isolate mutant alleles of the gene. Such an isolation is preferable in processes and disorders which are known or suspected to have a genetic basis. Mutant alleles can be isolated from individuals either known or suspected to have a genotype which contributes to tumor
30 progression symptoms. Mutant alleles and mutant allele products can then be utilized in the therapeutic and diagnostic assay systems described below.

465243 052397
A cDNA of a mutant gene can be isolated, for example, by using PCR, a technique which is well-known to one skilled in the art. In this case, the first cDNA strand can be synthesized by hybridizing a oligo-dT oligonucleotide to mRNA isolated from tissue known or suspected of being expressed in an individual putatively carrying the mutant allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA can then be synthesized using an oligonucleotide that hybridizes specifically to the 5'- end of the normal gene. Using these two primers, the product is then amplified via PCR, cloned into a suitable vector, and subjected to DNA sequence analysis through methods well-known to one skilled in the art. By comparing the DNA sequence of the mutant gene to that of the normal gene, the mutation(s) responsible for the loss or alteration of function of the mutant gene product can be ascertained.

Alternatively, a genomic or cDNA library can be constructed and screened using DNA or RNA, respectively, from a tissue known to or suspected of expressing the gene of interest in an individual suspected of or known to carry the mutant allele. The normal gene or any suitable fragment thereof can then be labeled and used as a probe to identify the corresponding mutant allele in the library. The clone containing this gene can then be purified through methods routinely practiced in the art, and subjected to sequence analysis as described, above, in this Section.

Additionally, an expression library can be constructed utilizing DNA isolated from or cDNA synthesized from a tissue known to or suspected of expressing the gene of interest in an individual suspected of or known to carry the mutant allele. In this manner, gene products made by the putatively mutant tissue can be expressed and screened

using standard antibody screening techniques in conjunction with antibodies raised against the normal gene product, as described, below, in Section 5.2.3. (For screening techniques, see, for example, Harlow, E. and Lane, eds., 1988, "Antibodies: A Laboratory Manual," Cold Spring Harbor Press, Cold Spring Harbor.) In cases where the mutation results in an expressed gene product with altered function (e.g., as a result of a missense mutation), a polyclonal set of antibodies are likely to cross-react with the mutant gene product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis as described in this Section, above.

Taking the fomy030 gene as an example, the fomy030 human homolog can be isolated by a variety of methods. First, sequences found in a murine fomy030 cDNA can be utilized as hybridization probes to detect human fomy030 sequences. This can be accomplished, for example, by probing Southern blots containing total human genomic DNA with a labelled fomy030 probe. Once it is verified that the probe being utilized detects the human 030 gene, one of skill in the art can employ any of several routine approaches to isolate the human gene without undue experimentation.

In one approach, appropriate human cDNA libraries can be screened. Such cDNA libraries can, for example, include human melanocyte, human retina and fetal human brain cDNA libraries. For example, panels of human melanoma cells (such as, for example, SK-MEL-2, ATCC 68-HTB; SK-MEL-5, ATCC 70-HTB; SK-MEL-28, ATCC 72-HTB; G-361, ATCC 1424-CRL; and/or HT-144 [63-HTB] cells) can be screened for 030 expression by, for example, Northern blot analysis. Upon detection of 030 transcript, cDNA libraries can be constructed from RNA isolated from the appropriate cell line, utilizing standard

techniques well known to those of skill in the art. The human cDNA library can then be screened with a 030 probe in order to isolate a human romy030 cDNA. As described below, this method was used to determine the human fohy030 cDNAs in
5 FIGS. 5 and 6.

Alternatively, a human total genomic DNA library can be screened using 030 probes. 030-positive clones can then be sequenced and, further, the intron/exon structure of the human 030 gene may be elucidated. Once genomic sequence is
10 obtained, oligonucleotide primers can be designed based on the sequence for use in the isolation, via, for example RT-coupled PCR, of human 030 cDNA.

The procedures described in these approaches are routine and have been described in detail in Sections
15 5.1.1.2, 5.3 and 5.7.2.

5.5. DIFFERENTIALLY EXPRESSED AND PATHWAY GENE PRODUCTS

Differentially expressed and pathway gene products include those proteins encoded by the differentially
20 expressed and pathway gene sequences described in Section 5.2.1, above, as for example, the peptide listed in FIG. 3. Specifically, differentially expressed and pathway gene products can include differentially expressed and pathway gene polypeptides encoded by the differentially expressed
25 and pathway gene sequences contained in the clones listed in Table 2, above, as deposited with the NRRL, or contained in the coding regions of the genes to which DNA sequences disclosed herein (in FIGS. 3A, 3B, 5, and 6) or contained in the clones, listed in Table 2, as deposited with the NRRL,
30 belong, for example.

In addition, differentially expressed and pathway gene products can include proteins that represent

functionally equivalent gene products. Such an equivalent differentially expressed or pathway gene product can contain deletions, additions or substitutions of amino acid residues within the amino acid sequence encoded by the differentially
5 expressed or pathway gene sequences described, above, in Section 5.2.1, but which result in a silent change thus producing a functionally equivalent differentially expressed on pathway gene product. Amino acid substitutions can be made on the basis of similarity in polarity, charge,
10 solubility, hydrophobicity, hydrophilicity, and/or the amphipatic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine,
15 serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid. "Functionally equivalent," as utilized herein, refers
20 to either a protein capable exhibiting a substantially similar in vivo activity as the endogenous differentially expressed or pathway gene products encoded by the differentially expressed or pathway gene sequences described in Section 5.2.1, above. Alternatively, when utilized as
25 part of assays such as those described, below, in Section 5.3, "functionally equivalent" can refer to peptides capable of interacting with other cellular or extracellular molecules in a manner substantially similar to the way in which the corresponding portion of the endogenous
30 differentially expressed or pathway gene product would.

The differentially expressed or pathway gene products can be produced by synthetic techniques or via recombinant DNA technology using techniques well known in

the art. Methods for preparing the differentially expressed or pathway gene polypeptides and peptides of the invention by expressing nucleic acid encoding differentially expressed or pathway gene sequences are described herein. Methods
5 which are well known to those skilled in the art can be used to construct expression vectors containing differentially expressed or pathway gene protein coding sequences and appropriate transcriptional/translational control signals. These methods include, for example, in vitro recombinant DNA
10 techniques, synthetic techniques and in vivo recombination/genetic recombination. See, for example, the techniques described in Maniatis et al., 1989, Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory, N.Y. which is incorporated by reference herein in their
15 entirety, and Ausubel, 1989, supra. Alternatively, RNA capable of encoding differentially expressed or pathway gene protein sequences can be chemically synthesized using, for example, synthesizers. See, for example, the techniques described in "Oligonucleotide Synthesis," 1984, Gait, M.J.
20 ed., IRL Press, Oxford, which is incorporated by reference herein in its entirety.

A variety of host-expression vector systems can be utilized to express the differentially expressed or pathway gene coding sequences of the invention. Such host-
25 expression systems represent vehicles by which the coding sequences of interest can be produced and subsequently purified, but also represent cells which can, when transformed or transfected with the appropriate nucleotide coding sequences, exhibit the differentially expressed or
30 pathway gene protein of the invention in situ. These include but are not limited to microorganisms such as bacteria (e.g., E. coli, B. subtilis) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA

expression vectors containing differentially expressed or pathway gene protein coding sequences; yeast (e.g., Saccharomyces, Pichia) transformed with recombinant yeast expression vectors containing the differentially expressed or pathway gene protein coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing the differentially expressed or pathway gene protein coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing differentially expressed or pathway gene protein coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

In bacterial systems, a number of expression vectors can be advantageously selected depending upon the use intended for the differentially expressed or pathway gene protein being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of antibodies or to screen peptide libraries, for example, vectors which direct the expression of high levels of fusion protein products that are readily purified can be desirable. Such vectors include, but are not limited, to the E. coli expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which the differentially expressed or pathway gene protein coding sequence can be ligated individually into the vector in frame with the lacZ coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye,

1985, Nucleic Acids Res. 13:3101-3109; Van Heeke & Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors can also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In
5 general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the
10 cloned target gene protein can be released from the GST moiety.

In an insect system, Autographa californica nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in Spodoptera frugiperda
15 cells. The differentially expressed or pathway gene coding sequence can be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example, the polyhedrin promoter). Successful insertion of
20 differentially expressed or pathway gene coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect
25 Spodoptera frugiperda cells in which the inserted gene is expressed (e.g., see Smith et al., 1983, J. Virol. 46:584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems can be utilized. In cases where an
30 adenovirus is used as an expression vector, the differentially expressed or pathway gene coding sequence of interest can be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and

tripartite leader sequence. This chimeric gene can then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a

5 recombinant virus that is viable and capable of expressing differentially expressed or pathway gene protein in infected hosts (e.g., See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals can also be required for efficient translation of inserted

10 differentially expressed or pathway gene coding sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire identified gene, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no

15 additional translational control signals can be needed. However, in cases where only a portion of the identified coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in

20 phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of

25 appropriate transcription enhancer elements, transcription terminators, etc., (see Bittner et al., 1987, Methods in Enzymol. 153:516-544).

In addition, a host cell strain can be chosen which modulates the expression of the inserted sequences, or

30 modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products can be important for the function of the protein. Different host

cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product can be used. Such mammalian host cells include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, etc.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the differentially expressed or pathway gene protein can be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells can be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method can advantageously be used to engineer cell lines which express the identified gene protein. Such engineered cell lines can be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the differentially expressed or pathway gene protein.

463250" 24429880

A number of selection systems can be used, including, but not limited to, the herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, Cell 22:817) genes in tk⁻, hgp⁻rt⁻ or aprt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for dhfr, which confers resistance to methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA 77:3567; O'Hare, et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, J. Mol. Biol. 150:1); and hyg⁺, which confers resistance to hygromycin (Santerre, et al., 1984, Gene 30:147) genes.

An alternative fusion protein system allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl. Acad. Sci. USA 88:8972-8976). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto ni²⁺ nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

When used as a component in assay systems such as that described herein, the differentially expressed or pathway gene protein can be labeled, either directly or indirectly, to facilitate detection of a complex formed

between the differentially expressed or pathway gene protein and a test substance. Any of a variety of suitable labeling systems can be used including but not limited to radioisotopes such as ^{125}I ; enzyme labelling systems that generate a detectable colorimetric signal or light when exposed to substrate; and fluorescent labels.

Where recombinant DNA technology is used to produce the differentially expressed or pathway gene protein for such assay systems, it can be advantageous to engineer fusion proteins that can facilitate labeling, solubility, immobilization and/or detection.

Indirect labeling involves the use of a third protein, such as a labeled antibody, which specifically binds to either a differentially expressed or pathway gene product. Such antibodies include but are not limited to polyclonal, monoclonal, chimeric, single chain, Fab fragments and fragments produced by a Fab expression library.

5.6. ANTIBODIES SPECIFIC FOR DIFFERENTIALLY EXPRESSED OR PATHWAY GENE PRODUCTS

Described herein are methods for the production of antibodies capable of specifically recognizing one or more differentially expressed or pathway gene epitopes. Such antibodies can include, but are not limited to polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')_2 fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. Such antibodies can be used, for example, in the detection of a fingerprint, target, or pathway gene in a biological sample, or,

alternatively, as a method for the inhibition of abnormal target gene activity. Thus, such antibodies can be utilized as tumor progression treatment methods, and/or can be used as part of diagnostic techniques whereby patients can be tested for abnormal levels of fingerprint, target, or pathway gene proteins, or for the presence of abnormal forms of the such proteins.

For the production of antibodies to a differentially expressed or pathway gene, various host animals can be immunized by injection with a differentially expressed or pathway gene protein, or a portion thereof. Such host animals can include but are not limited to rabbits, mice, and rats, to name but a few. Various adjuvants can be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*.

Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of animals immunized with an antigen, such as target gene product (e.g., protein encoded by 030), or an antigenic functional derivative thereof. For the production of polyclonal antibodies, host animals such as those described above, can be immunized by injection with differentially expressed or pathway gene product (e.g., 030) supplemented with adjuvants as also described above.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, can be obtained by any technique which provides for the production

of antibody molecules by continuous cell lines in culture. These include, but are not limited to the hybridoma technique of Kohler and Milstein, (1975, Nature 256:495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the BV-hybridoma technique (Cole et al., 1985, Monoclonal Antibodies And Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). Such antibodies can be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention can be cultivated in vitro or in vivo. Production of high titers of mAbs in vivo makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, Proc. Natl. Acad. Sci., 81:6851-6855; Neuberger et al., 1984, Nature, 312:604-608; Takeda et al., 1985, Nature, 314:452-454; U.S. Patent No. 4,816,567) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778; Bird, 1988, Science 242:423-426; Huston et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward et al., 1989, Nature 334:544-546) and for making humanized monoclonal antibodies (U.S. Patent No. 5,225,539, which is incorporated herein by reference in its entirety) can be

utilized to produce anti-differentially expressed or anti-pathway gene product antibodies.

Antibody fragments which recognize specific epitopes can be generated by known techniques. For example, such fragments include but are not limited to: the $F(ab')_2$ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the $F(ab')_2$ fragments. Alternatively, Fab expression libraries can be constructed (Huse *et al.*, 1989, Science, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

5.7. CELL- AND ANIMAL-BASED MODEL SYSTEMS

Described herein are cell- and animal-based systems which represent reliable models for tumor progression disorders. These systems can be used in a variety of applications. For example, the cell- and animal-based model systems can be used to identify differentially expressed genes via the paradigms described, above, in Section 5.1.1.1. Such systems can also be used to further characterize differentially expressed and pathway genes, as described, above, in Section 5.3. Such further characterization can, for example, indicate that a differentially expressed gene is a target gene, for example. Additionally, such assays can be utilized as part of screening strategies designed to identify compounds which are capable of preventing and/or ameliorating symptoms of tumor progression disorders, including those associated with metastatic diseases, as described, below. Thus, the animal- and cell-based models can be used to identify drugs, pharmaceuticals, therapies and interventions which can be effective in treating tumor progression disorders, such as,

for example, metastatic diseases. In addition, as described in detail, below, in Section 5.10.1, such animal models can be used to determine the LD₅₀ and the ED₅₀ in animal subjects, and such data can be used to determine the in vivo efficacy of potential anti-tumor progression disorder treatments.

5.7.1. ANIMAL-BASED SYSTEMS

Animal-based model systems of tumor progression disorders can be both non-recombinant animals as well as recombinantly engineered transgenic animals.

Non-recombinant animal models for tumor progression can include, for example, murine models of melanoma, prostate cancer and colon cancer. Such models may be generated, for example, by introducing tumor cells into syngeneic mice using techniques such as subcutaneous injection, tail vein injection, spleen implantation, intraperitoneal implantation, implantation under the renal capsule or orthotopic implantation (e.g., colon cancer cells implanted in colonic tissue or prostatic cancer cells implanted in prostate gland). After an appropriate period of time, the tumors which result from these injections can be counted and analyzed.

Among the cells which may be used for the production of such animal models of tumor progression are cells derived from the cell lines listed, above, in Section 5.1.1.1. For example, B16 melanoma cells (Fidler, I.J., 1973, Nature New Biol. 242:148-149), including cell variants exhibiting high (e.g., B16 F10 cells) and low (e.g., B16 F1 cells) metastatic potential may be utilized. Post-injection, pulmonary tumors generally develop in the mouse models. Thus, these animal serve as models of not only melanoma

268550 2442980
tumor progression but also as models of pulmonary metastases.

For the generation of animal models of colorectal cancers, colon cancer cells such as, for example, KM12c (low metastatic potential) and KM12L4 (highly metastatic) cells (Morikawa, K. et al., 1988, Cancer Research 48:1943-1948) can be implanted into nude mice spleens. In these cases, the animals generally develop hepatic tumors. Thus, such animals serve as models of not only colorectal tumor progression but also as models of hepatic metastases.

For the generation of animal models of prostate cancer tumor progression, cells derived from, for example, the high metastatic potential prostatic cell line PC-3-M or the non-metastatic cell line DU 145 (Karmali, R.A. et al., 1987, Anticancer Res. 7:1173-1180; Koziowski, J.M. et al., 1984, Cancer Research 44:3522-3529) may be implanted into the prostates of animals and the resulting tumors may be analyzed and compared to, for example, normal tissue. In such a manner, genes which are differentially expressed in neoplastic versus normal cells as well as versus metastatic cells may be identified.

The role of identified gene products (e.g., 030 gene products) can be determined by transfecting cDNAs encoding such gene products into the appropriate cell line and analyzing its effect on the cells' ability to induce tumor progression in animal models such as these. The role of the identified gene products may be further analyzed by, for example, culturing cells derived from the tumors which develop in the animal models, introducing these cultured cells into animals, and subsequently measuring the level of identified gene product present in the resulting tumor cells. In this manner, cell line variants are developed which can be useful in analyzing the role of quantitative

and/or qualitative differences in the expression of the identified genes on the cells' ability to induce tumor progression. For example, as demonstrated, below, in the Example presented in Section 6, 030 gene expression is
5 inversely related to the metastatic potential of the tumor cell line used to generate such a tumor progression animal model.

Additionally, recombinant animal models exhibiting tumor progression characteristics and/or symptoms of tumor
10 progression disorders, including metastatic diseases, can be utilized, for example, such well-known animal models as the transgenic mouse model for human melanoma and transgenic mice which carry specific mutations which result in multiple intestinal tumors (Mintz, M. and Silvers W.K., 1993, Proc.
15 Natl. Acad. Sci. USA 90:8817-8821; and Fodde, R., et al., 1994, Proc. Natl. Acad. Sci. USA 91:8969-8973). Further, recombinant animal models for tumor progression can be engineered by utilizing, for example, target gene sequences such as those described, above, in Section 5.4, in
20 conjunction with techniques for producing transgenic animals that are well known to those of skill in the art. For example, target gene sequences can be introduced into, and overexpressed in, the genome of the animal of interest, or, if endogenous target gene sequences are present, they can
25 either be overexpressed or, alternatively, can be disrupted in order to underexpress or inactivate target gene expression.

In order to overexpress a target gene sequence, the coding portion of the target gene sequence can be ligated to
30 a regulatory sequence which is capable of driving gene expression in the animal and cell type of interest. Such regulatory regions will be well known to those of skill in

the art, and can be utilized in the absence of undue experimentation.

In order to obtain underexpression of an endogenous target gene sequence, such a sequence can be introduced into the genome of the animal of interest such that the endogenous target gene alleles will be inactivated. Preferably, an engineered sequence comprising at least part of the target gene sequence is utilized and is introduced, via gene targeting, such that the endogenous target sequence is disrupted upon integration of the engineered target gene sequence into the animal's genome. Gene targeting is discussed, below, in this Section.

Animals of any species, including, but not limited to, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, goats, and non-human primates, e.g., baboons, monkeys, and chimpanzees can be used to generate animal models of tumor progression and tumor progression disorders, such as, for example, metastatic diseases.

Any technique known in the art can be used to introduce a target gene transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to pronuclear microinjection (Hoppe, P.C. and Wagner, T.E., 1989, U.S. Pat. No. 4,873,191); retrovirus mediated gene transfer into germ lines (Van der Putten et al., 1985, Proc. Natl. Acad. Sci., USA 82:6148-6152); gene targeting in embryonic stem cells (Thompson et al., 1989, Cell 56:313-321); electroporation of embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814); and sperm-mediated gene transfer (Lavitrano et al., 1989, Cell 57:717-723); etc. For a review of such techniques, see Gordon, 1989, Transgenic Animals, Intl. Rev. Cytol. 115:171-229, which is incorporated by reference herein in its entirety.

463350 24429830

The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, i.e., mosaic animals. The transgene can be integrated, either as a single transgene or in concatamers, e.g., head-to-head tandems or head-to-tail tandems. The transgene can also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko, M., et al., 1992, Proc. Natl. Acad. Sci. USA 89:6232-6236). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

When it is desired that the target gene transgene be integrated into the chromosomal site of the endogenous target gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous target gene of interest are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of, the nucleotide sequence of the endogenous target gene. The transgene can also be selectively introduced into a particular cell type, thus inactivating the endogenous gene of interest in only that cell type, by following, for example, the teaching of Gu et al. (Gu, H., et al., 1994, Science 265:103-106). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant target gene and protein can be assayed utilizing standard techniques. Initial screening can

be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals can also be assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, in situ hybridization analysis, and RT-coupled PCR. Samples of target gene-expressing tissue, can also be evaluated immunocytochemically using antibodies specific for the transgenic product of interest.

The target gene transgenic animals that express target gene mRNA or target gene transgene peptide (detected immunocytochemically, using antibodies directed against target gene product epitopes) at easily detectable levels should then be further evaluated to identify those animals which display tumor progression state characteristics, including tumor progression disorder symptoms. Such tumor progression disorder characteristics and/or symptoms can include, for example, those associated with such tumor cells as found in human melanoma, breast, gastrointestinal, such as esophageal, stomach, colon, bowel, colorectal and rectal cancers, prostate, bladder, testicular, ovarian, uterine, cervical, brain, lung, bronchial, larynx, pharynx, liver, pancreatic, thyroid, bone, leukemias, lymphomas and various types of skin cancers.

Additionally, specific cell types within the transgenic animals can be analyzed for cellular phenotypes characteristic of tumor progression. Such cellular phenotypes can include, for example, differential gene expression characteristic of cells within a given tumor progression state of interest. Further, such cellular phenotypes can include as assessment of a particular cell type fingerprint pattern of expression and its comparison to

known fingerprint expression profiles of the particular cell type in animals exhibiting tumor progression. Such transgenic animals serve as suitable model systems for tumor progression disorders.

5 Once target gene transgenic founder animals are produced (i.e., those animals which express target gene proteins in cells or tissues of interest, and which, preferably, exhibit tumor progression characteristics), they can be bred, inbred, outbred, or crossbred to produce
10 colonies of the particular animal. Examples of such breeding strategies include but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound
15 target gene transgenics that express the target gene transgene of interest at higher levels because of the effects of additive expression of each target gene transgene; crossing of heterozygous transgenic animals to produce animals homozygous for a given integration site in
20 order to both augment expression and eliminate the possible need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; breeding animals to different inbred genetic backgrounds so as to examine effects of modifying
25 alleles on expression of the target gene transgene and the development of symptoms for tumor progression disorders. One such approach is to cross the target gene transgenic founder animals with a wild type strain to produce an F1 generation that exhibits symptoms for tumor progression
30 disorders. The F1 generation can then be inbred in order to develop a homozygous line, if it is found that homozygous target gene transgenic animals are viable.

5.7.2. CELL-BASED ASSAYS

Cells that contain and express target gene sequences which encode target gene protein, and, further, exhibit cellular phenotypes associated with tumor progression disorders, can be utilized to identify compounds that exhibit an ability to prevent and/or ameliorate tumor progression. Cellular phenotypes which can indicate an ability to ameliorate symptoms of tumor progression disorders can include, for example, tumor cells with low or high metastatic potential.

Further, the fingerprint pattern of gene expression of cells of interest can be analyzed and compared to the normal fingerprint pattern. Those compounds which cause cells exhibiting cellular phenotypes of tumor progression disorders, including metastatic diseases, to produce a fingerprint pattern more closely resembling a normal fingerprint pattern for the cell of interest can be considered candidates for further testing regarding an ability to ameliorate the symptoms of such diseases.

Cells which will be utilized for such assays can, for example, include non-recombinant cell lines, such as, but not limited to, melanoma (e.g., B16 F1 and B16 F10 cell lines), human colon (e.g., KM12c and KM20L4 cell lines), prostate (e.g., DU 145 and PC-3-M cell lines) and breast cancer cell lines (e.g., MCF-7 and MDA-MB-435 cell lines). In addition, purified primary or secondary tumor cells derived from either transgenic or non-transgenic tumor cells can be used.

Further, cells which can be used for such assays can also include recombinant, transgenic cell lines. For example, the metastatic disease animal models of the invention, discussed, above, in Section 5.2.4.1, can be used to generate cell lines, containing one or more cell types

involved in metastatic diseases, that can be used as cell culture models for these disorders. While primary cultures derived from the metastasis in transgenic animals of the invention can be utilized, the generation of continuous cell lines is preferred. For examples of techniques which can be used to derive a continuous cell line from the transgenic animals, see Small et al., 1985, Mol. Cell Biol. 5:642-648.

Alternatively, cells of a cell type known to be involved in metastatic diseases can be transfected with sequences capable of increasing or decreasing the amount of target gene expression within the cell. For example, target gene sequences can be introduced into, and over expressed in, the genome of the cell of interest, or, if endogenous target gene sequences are present, they can either be overexpressed or, alternatively, be disrupted in order to underexpress or inactivate target gene expression.

In order to overexpress a target gene sequence, the coding portion of the target gene sequence can be ligated to a regulatory sequence which is capable of driving gene expression in the cell type of interest. Such regulatory regions will be well known to those of skill in the art, and can be utilized in the absence of undue experimentation.

For under expression of an endogenous target gene sequence, such a sequence can be isolated and engineered such that when reintroduced into the genome of the cell type of interest, the endogenous target gene alleles will be inactivated. Preferably, the engineered target gene sequence is introduced via gene targeting such that the endogenous target sequence is disrupted upon integration of the engineered target gene sequence into the cell's genome. Gene targeting is discussed, above, in Section 5.7.1.

Transfection of target gene sequence nucleic acid can be accomplished by utilizing standard techniques. See,

for example, Ausubel, 1989, supra. Transfected cells should be evaluated for the presence of the recombinant target gene sequences, for expression and accumulation of target gene mRNA, and for the presence of recombinant target gene protein production. In instances wherein a decrease in target gene expression is desired, standard techniques can be used to demonstrate whether a decrease in endogenous target gene expression and/or in target gene product production is achieved.

5.8. SCREENING ASSAYS FOR COMPOUNDS THAT INTERACT WITH THE TARGET GENE PRODUCT

The following assays are designed to identify compounds that bind to target gene products, bind to other cellular proteins that interact with a target gene product, and to compounds that interfere with the interaction of the target gene product with other cellular proteins.

Such compounds can include, but are not limited to, other cellular proteins. Specifically, such compounds can include, but are not limited to, peptides, such as, for example, soluble peptides, including, but not limited to Ig-tailed fusion peptides, comprising extracellular portions of target gene product transmembrane receptors, and members of random peptide libraries (see, e.g., Lam, K.S. et al., 1991, Nature 354:82-84; Houghton, R. et al., 1991, Nature 354:84-86), made of D-and/or L-configuration amino acids, phosphopeptides (including, but not limited to, members of random or partially degenerate phosphopeptide libraries; see, e.g., Songyang, Z. et al., 1993, Cell 72:767-778), antibodies (including, but not limited to, polyclonal, monoclonal, humanized, anti-idiotypic, chimeric or single chain antibodies, and FAb, F(ab')₂ and FAb expression library

fragments, and epitope-binding fragments thereof), and small organic or inorganic molecules.

Compounds identified via assays such as those described herein can be useful, for example, in elaborating the biological function of the target gene product, and for ameliorating symptoms of tumor progression. In instances, for example, whereby a tumor progression state or disorder results from a lower overall level of target gene expression, target gene product, and/or target gene product activity in a cell involved in the tumor progression state or disorder, compounds that interact with the target gene product can include ones which accentuate or amplify the activity of the bound target gene protein. Such compounds would bring about an effective increase in the level of target gene activity, thus ameliorating symptoms of the tumor progression disorder or state. In instances whereby mutations within the target gene cause aberrant target gene proteins to be made which have a deleterious effect that leads to tumor progression, compounds that bind target gene protein can be identified that inhibit the activity of the bound target gene protein. Assays for testing the effectiveness of compounds, identified by, for example, techniques such as those described in Section 5.8.1-5.8.3, are discussed, below, in Section 5.8.4.

5.8.1. IN VITRO SCREENING ASSAYS FOR COMPOUNDS THAT BIND TO A TARGET GENE PRODUCT

In vitro systems can be designed to identify compounds capable of binding the target gene products of the invention. Compounds identified can be useful, for example, in modulating the activity of wild type and/or mutant target gene products, preferably mutant target gene proteins, can be useful in elaborating the biological function of the

target gene product, can be utilized in screens for identifying compounds that disrupt normal target gene interactions, or can in themselves disrupt such interactions.

5 The principle of the assays used to identify compounds that bind to the target gene product involves preparing a reaction mixture of the target gene protein and the test compound under conditions and for a time sufficient to allow the two components to interact and bind, thus
10 forming a complex which can be removed and/or detected in the reaction mixture. These assays can be conducted in a variety of ways. For example, one method to conduct such an assay would involve anchoring target gene product or the
15 product/test compound complexes anchored on the solid phase at the end of the reaction. In one embodiment of such a method, the target gene product can be anchored onto a solid surface, and the test compound, which is not anchored, can be labeled, either directly or indirectly.

20 In practice, microtitre plates can conveniently be utilized as the solid phase. The anchored component can be immobilized by non-covalent or covalent attachments. Non-covalent attachment can be accomplished by simply coating the solid surface with a solution of the protein and drying.
25 Alternatively, an immobilized antibody, preferably a monoclonal antibody, specific for the protein to be immobilized can be used to anchor the protein to the solid surface. The surfaces can be prepared in advance and stored.

30 In order to conduct the assay, the nonimmobilized component is added to the coated surface containing the anchored component. After the reaction is complete, unreacted components are removed (e.g., by washing) under

conditions such that any complexes formed will remain immobilized on the solid surface. The detection of complexes anchored on the solid surface can be accomplished in a number of ways. Where the previously immobilized component is pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed. Where the previously nonimmobilized component is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the immobilized component (the antibody, in turn, can be directly labeled or indirectly labeled with a labeled anti-Ig antibody).

Alternatively, a reaction can be conducted in a liquid phase, the reaction products separated from unreacted components, and complexes detected; e.g., using an immobilized antibody specific for target gene or the test compound to anchor any complexes formed in solution, and a labeled antibody specific for the other component of the possible complex to detect anchored complexes.

5.8.2. ASSAYS FOR CELLULAR PROTEINS THAT INTERACT WITH THE TARGET GENE PRODUCT

Any method suitable for detecting protein-protein interactions can be employed for identifying novel target product-cellular or extracellular protein interactions.

These methods are outlined in Section 5.1.3., supra, for the identification of pathway genes, and can be utilized herein with respect to the identification of proteins which interact with identified target proteins. In such a case, the target gene serves as the known "bait" gene.

**5.8.3. ASSAYS FOR COMPOUNDS THAT
INTERFERE WITH TARGET
GENE/CELLULAR PRODUCT INTERACTION**

The target gene products of the invention can, in vivo, interact with one or more cellular or extracellular macromolecules, such as proteins. Such macromolecules include, but are not limited to, nucleic acid molecules and those products identified via methods such as those described, above, in Section 5.8.2. For the purposes of this discussion, such cellular and extracellular macromolecules are referred to herein as "binding partners." Compounds that disrupt such interactions can be useful in regulating the activity of the target gene product, especially mutant target gene products. Such compounds can include, but are not limited to molecules such as antibodies, peptides, and the like described in Section 5.3.1. above.

The basic principle of the assay systems used to identify compounds that interfere with the interaction between the target gene product and its cellular or extracellular binding partner or partners involves preparing a reaction mixture containing the target gene product, and the binding partner under conditions and for a time sufficient to allow the two products to interact and bind, thus forming a complex. In order to test a compound for inhibitory activity, the reaction mixture is prepared in the presence and absence of the test compound. The test compound can be initially included in the reaction mixture, or can be added at a time subsequent to the addition of target gene and its cellular or extracellular binding partner. Control reaction mixtures are incubated without the test compound or with a placebo. The formation of any complexes between the target gene product and the cellular

or extracellular binding partner is then detected. The formation of a complex in the control reaction, but not in the reaction mixture containing the test compound, indicates that the compound interferes with the interaction of the target gene product and the interactive binding partner. Additionally, complex formation within reaction mixtures containing the test compound and normal target gene product can also be compared to complex formation within reaction mixtures containing the test compound and mutant target gene product. This comparison can be important in those cases wherein it is desirable to identify compounds that disrupt interactions of mutant but not normal target gene products.

The assay for compounds that interfere with the interaction of the target gene products and binding partners can be conducted in a heterogeneous or homogeneous format. Heterogeneous assays involve anchoring either the target gene product or the binding partner onto a solid phase and detecting complexes anchored on the solid phase at the end of the reaction. In homogeneous assays, the entire reaction is carried out in a liquid phase. In either approach, the order of addition of reactants can be varied to obtain different information about the compounds being tested. For example, test compounds that interfere with the interaction between the target gene products and the binding partners, e.g., by competition, can be identified by conducting the reaction in the presence of the test substance; i.e., by adding the test substance to the reaction mixture prior to or simultaneously with the target gene product and interactive cellular or extracellular binding partner. Alternatively, test compounds that disrupt preformed complexes, e.g. compounds with higher binding constants that displace one of the components from the complex, can be tested by adding the test compound to the reaction mixture

after complexes have been formed. The various formats are described briefly below.

In a heterogeneous assay system, either the target gene product or the interactive cellular or extracellular binding partner, is anchored onto a solid surface, while the non-anchored species is labeled, either directly or indirectly. In practice, microtitre plates are conveniently utilized. The anchored species can be immobilized by non-covalent or covalent attachments. Non-covalent attachment can be accomplished simply by coating the solid surface with a solution of the target gene product or binding partner and drying. Alternatively, an immobilized antibody specific for the species to be anchored can be used to anchor the species to the solid surface. The surfaces can be prepared in advance and stored.

In order to conduct the assay, the partner of the immobilized species is exposed to the coated surface with or without the test compound. After the reaction is complete, unreacted components are removed (e.g., by washing) and any complexes formed will remain immobilized on the solid surface. The detection of complexes anchored on the solid surface can be accomplished in a number of ways. Where the non-immobilized species is pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed. Where the non-immobilized species is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the initially non-immobilized species (the antibody, in turn, can be directly labeled or indirectly labeled with a labeled anti-Ig antibody). Depending upon the order of addition of reaction components, test compounds which inhibit complex formation or which disrupt preformed complexes can be detected.

Alternatively, the reaction can be conducted in a liquid phase in the presence or absence of the test compound, the reaction products separated from unreacted components, and complexes detected; e.g., using an immobilized antibody specific for one of the binding components to anchor any complexes formed in solution, and a labeled antibody specific for the other partner to detect anchored complexes. Again, depending upon the order of addition of reactants to the liquid phase, test compounds which inhibit complex or which disrupt preformed complexes can be identified.

In an alternate embodiment of the invention, a homogeneous assay can be used. In this approach, a preformed complex of the target gene product and the interactive cellular or extracellular binding partner product is prepared in which either the target gene products or their binding partners are labeled, but the signal generated by the label is quenched due to complex formation (see, e.g., U.S. Patent No. 4,109,496 by Rubenstein which utilizes this approach for immunoassays). The addition of a test substance that competes with and displaces one of the species from the preformed complex will result in the generation of a signal above background. In this way, test substances which disrupt target gene product-cellular or extracellular binding partner interaction can be identified.

In a particular embodiment, the target gene product can be prepared for immobilization using recombinant DNA techniques described in Section 5.1.2, supra. For example, the target gene coding region can be fused to a glutathione-S-transferase (GST) gene using a fusion vector such as pGEX-5X-1, in such a manner that its binding activity is maintained in the resulting fusion product. The interactive cellular or extracellular product can be purified and used

to raise a monoclonal antibody, using methods routinely practiced in the art and described above, in Section 5.2.4. This antibody can be labeled with the radioactive isotope ^{125}I , for example, by methods routinely practiced in the art. In a heterogeneous assay, e.g., the GST-Target gene fusion product can be anchored to glutathione-agarose beads. The interactive cellular or extracellular binding partner product can then be added in the presence or absence of the test compound in a manner that allows interaction and binding to occur. At the end of the reaction period, unbound material can be washed away, and the labeled monoclonal antibody can be added to the system and allowed to bind to the complexed components. The interaction between the target gene product and the interactive cellular or extracellular binding partner can be detected by measuring the amount of radioactivity that remains associated with the glutathione-agarose beads. A successful inhibition of the interaction by the test compound will result in a decrease in measured radioactivity.

Alternatively, the GST-target gene fusion product and the interactive cellular or extracellular binding partner product can be mixed together in liquid in the absence of the solid glutathione-agarose beads. The test compound can be added either during or after the binding partners are allowed to interact. This mixture can then be added to the glutathione-agarose beads and unbound material is washed away. Again the extent of inhibition of the binding partner interaction can be detected by adding the labeled antibody and measuring the radioactivity associated with the beads.

In another embodiment of the invention, these same techniques can be employed using peptide fragments that correspond to the binding domains of the target gene product

and the interactive cellular or extracellular binding partner (in case where the binding partner is a product), in place of one or both of the full length products. Any number of methods routinely practiced in the art can be used to identify and isolate the protein's binding site. These methods include, but are not limited to, mutagenesis of one of the genes encoding one of the products and screening for disruption of binding in a co-immunoprecipitation assay. Compensating mutations in the gene encoding the second species in the complex can be selected. Sequence analysis of the genes encoding the respective products will reveal the mutations that correspond to the region of the product involved in interactive binding. Alternatively, one product can be anchored to a solid surface using methods described in this Section above, and allowed to interact with and bind to its labeled binding partner, which has been treated with a proteolytic enzyme, such as trypsin. After washing, a short, labeled peptide comprising the binding domain can remain associated with the solid material, which can be isolated and identified by amino acid sequencing. Also, once the gene coding for the cellular or extracellular binding partner product is obtained, short gene segments can be engineered to express peptide fragments of the product, which can then be tested for binding activity and purified or synthesized.

5.8.4. ASSAYS FOR AMELIORATION OF TUMOR PROGRESSION SYMPTOMS

Any of the binding compounds, including but not limited to, compounds such as those identified in the foregoing assay systems, can be tested for the ability to prevent and/or ameliorate symptoms of tumor progression and tumor progression disorders, including metastatic disease.

Cell-based and animal model-based assays for the identification of compounds exhibiting an ability to prevent and/or ameliorate tumor progression symptoms are described below.

5 First, cell-based systems such as those described, above, in Section 5.7.2, can be used to identify compounds which can act to ameliorate symptoms of tumor progression . For example, such cell systems can be exposed to a compound, suspected to exhibiting an ability to ameliorate tumor
10 progression symptoms, at a sufficient concentration and for a time sufficient to elicit such an amelioration in the exposed cells. After exposure, the cells are examined to determine whether one or more tumor progression state or tumor progression disorder phenotypes has been altered to
15 resemble a more normal or more wild-type, non-neoplastic disease phenotype.

Taking, as an example, tumor progression involving metastasis, cell-based systems such as the highly metastatic B16 F10 melanoma cell line can be utilized. Upon exposure
20 to such cell systems, compounds can be assayed for their ability to reduce the metastatic potential of such cells. Further, the level of 030 gene expression within these cells may be assayed. Presumably, an increase in the observed level of 030 gene expression would indicate an amelioration
25 of the metastatic tumor progression state.

In addition, animal-based systems, such as those described, above, in Section 5.7.1, can be used to identify compounds capable of ameliorating symptoms of tumor progression. Such animal models can be used as test
30 substrates for the identification of drugs, pharmaceuticals, therapies, and interventions which can be effective in treating tumor progression disorders. For example, animal models can be exposed to a compound suspected to exhibit an

ability to ameliorate tumor progression symptoms, at a sufficient concentration and for a time sufficient to elicit such an amelioration in the exposed animals. The response of the animals to the exposure can be monitored by assessing the reversal of disorders associated with tumor progression. With regard to intervention, any treatments which reverse any aspect of symptoms of tumor progression, such as, for example, those associated with metastatic disease, should be considered as candidates for human therapeutic intervention in the treatment of tumor progression. Dosages of test agents can be determined by deriving dose-response curves, as discussed in Section 5.10, below.

Further, gene expression patterns can be utilized to assess the ability of a compound to ameliorate symptoms of tumor progression and tumor progression disorders. For example, fingerprint gene expression or a fingerprint pattern can then be used in such an assessment. Fingerprint gene expression and fingerprint patterns are described, below, in Section 5.11.

Fingerprint patterns can be characterized for known states (e.g., normal or known pre-neoplastic, neoplastic or metastatic states) within the cell- and/or animal-based model systems. Subsequently, these known fingerprint patterns can be compared to ascertain the effect a test compound has to modify such fingerprint patterns, and to cause the pattern to more closely resemble that of a more desirable fingerprint pattern.

For example, administration of a compound can cause the fingerprint pattern of a metastatic disease model system to more closely resemble a control, normal system. Administration of a compound can, alternatively, cause the fingerprint pattern of a control system to begin to mimic tumor progression states, such as metastatic disease states.

5.8.5. MONITORING OF EFFECTS DURING CLINICAL TRIALS

Monitoring the influence of compounds on tumor progression can be applied not only in basic drug screening, but also in clinical trials. In such clinical trials, the expression of a panel of genes that have been discovered in any one of the paradigms discovered in Section 5.1.1.1 can be used as a "read out" of the tumor progression state of a particular cell.

For example, and not by way of limitation, the paradigm describing the B16 melanoma cells provides for the identification of fingerprint genes (e.g., 030) that are down-regulated in metastatic tumor cells. For example, in a clinical trial, tumor cells can be isolated from the primary tumors removed by surgery, and RNA prepared and analyzed by differential display as described in Section 6.1. The levels of expression of the fingerprint genes can be quantified by Northern blot analysis or RT-PCR, as described in Section 6.1, or alternatively by measuring the amount of protein produced, by one of the methods described in Section 5.7.2. In this way, the fingerprint profiles can serve as putative biomarkers indicative of the metastatic potential of the tumor cell. Thus, by monitoring the level of expression of romy030, a protocol for suitable chemotherapeutic anticancer drugs can be developed based on the metastatic potential of tumor cells in the primary. In cases of inoperable metastatic disease, patients can have biopsies removed for measurement of romy030 expression so that the drug's efficacy can be measured by monitoring the degree of restored expression of romy030.

5.9. COMPOUNDS AND METHODS FOR TREATMENT OF TUMOR PROGRESSION

Described herein are methods and compositions which can be used ameliorate symptoms of tumor progression and disorders involving tumor progression via, first, target gene modulation, and/or second, via a depletion of the cells involved in tumor progression. Target gene modulation can be of a positive or negative nature, depending on the specific situation involved, but each modulatory event yields a net result in which tumor progression symptoms are ameliorated.

"Negative modulation," as used herein, refers to a reduction in the level and/or activity of target gene product relative to the level and/or activity of the target gene product in the absence of the modulatory treatment.

"Positive modulation," as used herein, refers to an increase in the level and/or activity of target gene product relative to the level and/or activity of target gene product in the absence of modulatory treatment.

It is possible that tumor progression can be brought about, at least in part, by an abnormal level of gene product, or by the presence of a gene product exhibiting abnormal activity. As such, the reduction in the level and/or activity of such gene products would bring about the amelioration of tumor progression symptoms. Negative modulatory techniques for the reduction of target gene expression levels or target gene product activity levels are discussed in Section 5.9.1, below.

Alternatively, it is possible that tumor progression can be brought about, at least in part, by the absence or reduction of the level of gene expression, or a reduction in the level of a gene product's activity. As such, an increase in the level of gene expression and/or the activity

of such gene products would bring about the amelioration of tumor progression symptoms.

For example, as demonstrated in the Example presented in Section 6, below, a reduction in the level of 030 gene expression correlates with a highly metastatic tumor progression state. A 030 positive modulatory technique which increased 030 gene expression in cells within a highly metastatic tumor progression state should, therefore, act to ameliorate the symptoms of such a state. Further, because the 030 gene product may exhibit general tumor suppressor features, it is possible that a 030 positive modulatory technique could ameliorate symptoms of many tumor progression events.

Positive modulatory techniques for increasing the target gene expression levels or target gene product activity levels are discussed in Section 5.9.2, below.

Additionally, tumor progression treatment techniques whereby the concentration of cells involved in tumor progression are depleted are described, below, in Section 5.9.3.

Among the tumor progression events which may be treated are those associated with human tumors. Such human tumors may include, for example, human melanomas, breast, gastrointestinal, such as esophageal, stomach, colon, bowel, colorectal and rectal cancers, prostate, bladder, testicular, ovarian, uterine, cervical, brain, lung, bronchial, larynx, pharynx, liver, pancreatic, thyroid, bone, leukemias, lymphomas and various types of skin cancers.

5.9.1. NEGATIVE MODULATORY TECHNIQUES

As discussed, above, successful treatment of tumor progression symptoms and of disorders involving tumor progression can be brought about by techniques which serve
5 to inhibit the expression or activity of target gene products.

For example, compounds such as those identified through assays described, above, in Section 5.8, which exhibit negative modulatory activity, can be used in
10 accordance with the invention to prevent and/or ameliorate symptoms of tumor progression, including tumor progression involving metastatic disorders. As discussed in Section 5.8., above, such molecules can include, but are not limited to peptides, phosphopeptides, small organic or inorganic
15 molecules, or antibodies (including, for example, polyclonal, monoclonal, humanized, anti-idiotypic, chimeric or single chain antibodies, and FAb, F(ab')₂ and FAb expression library fragments, and epitope-binding fragments thereof). Negative modulatory techniques involving antibody
20 administration are described, below, in Section 5.9.1.2. Techniques for the determination and administration of such compounds are described, below, in Section 5.10.

Further, antisense and ribozyme molecules which inhibit expression of the target gene can also be used in
25 accordance with the invention to reduce the level of target gene expression, thus effectively reducing the level of target gene activity. Still further, triple helix molecules can be utilized in reducing the level of target gene activity. Such techniques are described, below, in Section
30 5.9.1.1.

**5.9.1.1. NEGATIVE MODULATORY ANTISENSE,
RIBOZYME AND TRIPLE HELIX APPROACHES**

Among the compounds which can exhibit the ability to prevent and/or ameliorate symptoms of tumor progression are antisense, ribozyme, and triple helix molecules. Such molecules can be designed to reduce or inhibit either wild type, or if appropriate, mutant target gene activity. Techniques for the production and use of such molecules are well known to those of skill in the art.

Anti-sense RNA and DNA molecules act to directly block the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. With respect to antisense DNA, oligodeoxyribonucleotides derived from the translation initiation site, e.g., between the -10 and +10 regions of the Target gene nucleotide sequence of interest, are preferred.

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. (For a review, see, for example, Rossi, J., 1994, Current Biology 4:469-471). The mechanism of ribozyme action involves sequence specific hybridization of the ribozyme molecule to complementary target RNA, followed by a endonucleolytic cleavage. The composition of ribozyme molecules must include one or more sequences complementary to the target gene mRNA and must include the well-known catalytic sequence responsible for mRNA cleavage. For this sequence, see U.S. Pat. No. 5,093,246, which is incorporated by reference herein in its entirety. As such within the scope of the invention are engineered hammerhead motif ribozyme molecules that specifically and efficiently catalyze endonucleolytic cleavage of RNA sequences encoding target gene proteins.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning

the molecule of interest for ribozyme cleavage sites which include the following sequences, GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site can be evaluated for predicted structural features, such as secondary structure, that can render the oligonucleotide sequence unsuitable. The suitability of candidate sequences can also be evaluated by testing their accessibility to hybridization with complementary oligonucleotides, using ribonuclease protection assays.

Nucleic acid molecules to be used in triplex helix formation for the inhibition of transcription should be single stranded and composed of deoxynucleotides. The base composition of these oligonucleotides must be designed to promote triple helix formation via Hoogsteen base pairing rules, which generally require sizeable stretches of either purines or pyrimidines to be present on one strand of a duplex. Nucleotide sequences can be pyrimidine-based, which will result in TAT and CGC⁺ triplets across the three associated strands of the resulting triple helix. The pyrimidine-rich molecules provide base complementarity to a purine-rich region of a single strand of the duplex in a parallel orientation to that strand. In addition, nucleic acid molecules can be chosen that are purine-rich, for example, contain a stretch of G residues. These molecules will form a triple helix with a DNA duplex that is rich in GC pairs, in which the majority of the purine residues are located on a single strand of the targeted duplex, resulting in GGC triplets across the three strands in the triplex.

Alternatively, the potential sequences that can be targeted for triple helix formation can be increased by creating a so called "switchback" nucleic acid molecule.

26250" attc3880

Switchback molecules are synthesized in an alternating 5'-3', 3'-5' manner, such that they base pair with first one strand of a duplex and then the other, eliminating the necessity for a sizeable stretch of either purines or pyrimidines to be present on one strand of a duplex.

In instances wherein the antisense, ribozyme, and/or triple helix molecules described herein are utilized to reduce or inhibit mutant gene expression, it is possible that the technique utilized can also efficiently reduce or inhibit the transcription (triple helix) and/or translation (antisense, ribozyme) of mRNA produced by normal target gene alleles such that the possibility can arise wherein the concentration of normal target gene product present can be lower than is necessary for a normal phenotype. In such cases, to ensure that substantially normal levels of target gene activity are maintained, nucleic acid molecules that encode and express target gene polypeptides exhibiting normal target gene activity can be introduced into cells via gene therapy methods such as those described, below, in Section 5.9.2 that do not contain sequences susceptible to whatever antisense, ribozyme, or triple helix treatments are being utilized. Alternatively, in instances whereby the target gene encodes an extracellular protein, it can be preferable to coadminister normal target gene protein into the cell or tissue in order to maintain the requisite level of cellular or tissue target gene activity.

Anti-sense RNA and DNA, ribozyme and triple helix molecules of the invention can be prepared by any method known in the art for the synthesis of DNA and RNA molecules. These include techniques for chemically synthesizing oligodeoxyribonucleotides and oligoribonucleotides well known in the art such as, for example, solid phase phosphoramidite chemical synthesis. Alternatively, RNA

26850 " 24429880

molecules can be generated by in vitro and in vivo transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences can be incorporated into a wide variety of vectors which incorporate suitable RNA polymerase promoters such as the T7 or SP6 polymerase promoters. Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly, depending on the promoter used, can be introduced stably into cell lines.

10 Various well-known modifications to the DNA molecules can be introduced as a means of increasing intracellular stability and half-life. Possible modifications include but are not limited to the addition of flanking sequences of ribo- or deoxy- nucleotides to the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the oligodeoxyribonucleotide backbone.

5.9.1.2. NEGATIVE MODULATORY ANTIBODY TECHNIQUES

20 Antibodies can be generated which are both specific for target gene product and which reduce target gene product activity. Such antibodies may, therefore, be administered in instances whereby negative modulatory techniques are appropriate for the treatment of tumor progression.

25 Antibodies can be generated using standard techniques described in Section 5.6, above, against the proteins themselves or against peptides corresponding to portions of the proteins. The antibodies include but are not limited to polyclonal, monoclonal, Fab fragments, single chain antibodies, chimeric antibodies, and the like.

30 In instances where the target gene protein to which the antibody is directed is intracellular and whole

antibodies are used, internalizing antibodies can be preferred. However, lipofectin or liposomes can be used to deliver the antibody or a fragment of the Fab region which binds to the target gene epitope into cells. Where

5 fragments of the antibody are used, the smallest inhibitory fragment which binds to the target protein's binding domain is preferred. For example, peptides having an amino acid sequence corresponding to the domain of the variable region of the antibody that binds to the target gene protein can be

10 used. Such peptides can be synthesized chemically or produced via recombinant DNA technology using methods well known in the art (e.g., see Creighton, 1983, supra; and Sambrook et al., 1989, supra). Alternatively, single chain neutralizing antibodies which bind to intracellular target

15 gene product epitopes can also be administered. Such single chain antibodies can be administered, for example, by expressing nucleotide sequences encoding single-chain antibodies within the target cell population by utilizing, for example, techniques such as those described in Marasco

20 et al. (Marasco, W. et al., 1993, Proc. Natl. Acad. Sci. USA 90:7889-7893).

In instances where the target gene protein is extracellular, or is a transmembrane protein, any of the administration techniques described, below in Section 5.10

25 which are appropriate for peptide administration can be utilized to effectively administer inhibitory target gene antibodies to their site of action.

5.9.2. POSITIVE MODULATORY TECHNIQUES

As discussed above, successful treatment of tumor

30 progression symptoms and of disorders involving tumor progression can be brought about by techniques which serve

to increase the level of target gene expression or to increase the activity of a target gene product.

For example, compounds such as those identified through assays described, above, in Section 5.8, which exhibit positive modulatory activity can be used in accordance with the invention to ameliorate tumor progression symptoms. As discussed in Section 5.8, above, such molecules can include, but are not limited to, peptides, phosphopeptides, small organic or inorganic molecules, or antibodies (including, for example, polyclonal, monoclonal, humanized, anti-idiotypic, chimeric or single chain antibodies, and FAb, F(ab')₂ and FAb expression library fragments, and epitope-binding fragments thereof). Positive modulatory techniques involving antibody administration are described, below, in Section 5.9.2.1.

For example, a target gene protein, at a level sufficient to ameliorate tumor progression symptoms can be administered to a patient exhibiting such symptoms. Any of the techniques discussed, below, in Section 5.10, can be utilized for such administration. One of skill in the art will readily know how to determine the concentration of effective, non-toxic doses of the normal target gene protein, utilizing techniques such as those described, below, in Section 5.10.1.

In instances wherein the compound to be administered is a peptide compound, DNA sequences encoding the peptide compound can, alternatively, be directly administered to a patient exhibiting tumor progression symptoms, at a concentration sufficient to generate the production of an amount of target gene product adequate to ameliorate tumor progression symptoms. Any of the techniques described, below, in Section 5.10, which achieve intracellular administration, can be utilized for the administration of

such DNA molecules. The DNA molecules can be produced, for example, by well-known recombinant techniques.

In the case of peptide compounds which act extracellularly, the DNA molecules encoding such peptides
5 can be taken up and expressed by any cell type, so long as a sufficient circulating concentration of peptide results for the elicitation of a reduction in tumor progression symptoms.

In the case of compounds which act intracellularly,
10 the DNA molecules encoding such peptides must be taken up and expressed by cells involved in the tumor progression at a sufficient level to bring about the reduction of tumor progression symptoms.

Any technique which serves to selectively administer
15 DNA molecules to a cell involved in tumor progression is, therefore, preferred for the DNA molecules encoding intracellularly acting peptides.

Further, patients can be treated for symptoms of tumor progression by gene replacement therapy. One or more
20 copies of a normal target gene or a portion of the gene that directs the production of a normal target gene protein with target gene function can be inserted into cells, using vectors which include, but are not limited to adenovirus, adeno-associated virus, and retrovirus vectors, in addition
25 to other particles that introduce DNA into cells, such as liposomes. Techniques such as those described above can be utilized for the introduction of normal target gene sequences into human cells.

In instances wherein the target gene encodes an
30 extracellular, secreted gene product, such gene replacement techniques may be accomplished either in vivo or in vitro. For such cases, the cell types expressing the target gene is less important than achieving a sufficient circulating

concentration of the extracellular molecules for the amelioration of tumor progression symptoms to occur. In vitro, target gene sequences can be introduced into autologous cells. Those cells expressing the target gene
5 sequence of interest can then be reintroduced, preferably by intravenous administration, into the patient such that there results an amelioration of tumor progression symptoms.

In instances wherein the gene replacement involves a gene which encodes a product which acts intracellularly, it
10 is preferred that gene replacement be accomplished in vivo. Further, because the cell type in which the gene replacement must occur is the cell type involved in tumor progression, such techniques must successfully target such tumor progression cells.

15 Taking the 030 gene as an example, an increase in 030 expression can serve to ameliorate tumor progression symptoms, such as, for example, tumor progression symptoms involving metastatic processes. Therefore, any positive modulatory described herein which increases the 030 gene
20 product or gene product activity to a level which is sufficient to ameliorate tumor progression symptoms represents a successful tumor progression therapeutic treatment.

25 5.9.3. METHODS FOR DEPLETING CELLS INVOLVED IN TUMOR PROGRESSION

Techniques described herein can be utilized to deplete the total number of cells involved in tumor progression, thus effectively decreasing the ratio of the tumor cells to non-cancerous cells. Specifically,
30 separation techniques are described which can be used to deplete the total number of tumor cells present within a cell population, and, further, targeting techniques are

described which can be utilized to deplete specific tumor cell subpopulations.

Depending on the particular application, changing the number of cells belonging to tumor cell population can yield inhibitory responses leading to the amelioration of cancerous disorders.

The separation techniques described herein are based on the presence or absence of specific cell surface, preferably transmembrane, markers. By way of example, and not by way of limitation, the techniques described herein utilize tumor specific cell surface markers or antigens and will describe procedures whereby tumor cells can be separated from other cells, thus allowing for selective depletion of tumor cells.

Separation techniques can be utilized which separate and purify cells, tumor cells, for example, in vitro from a population of cells, such as hematopoietic cells autologous to the patient being treated. For example, an initial tumor cell subpopulation-containing population of cells, such as hematopoietic cells, can be obtained from a leukemia patient using standard procedures well known to those of skill in the art. Peripheral blood can be utilized as one potential starting source for such techniques, and can, for example, be obtained via venipuncture and collection into heparinized tubes.

Once the starting source of autologous cells is obtained, tumor cells can be removed, and thus selectively separated and purified, by various methods which utilize antibodies which bind specific markers present on tumor cells while absent on other cells within the starting source. These techniques can include, for example, flow cytometry using a fluorescence activated cell sorter (FACS) and specific fluorochromes, biotin-avidin or

biotin-streptavidin separations using biotin conjugated to cell surface marker-specific antibodies and avidin or streptavidin bound to a solid support such as affinity column matrix or plastic surfaces or magnetic separations using antibody-coated magnetic beads.

Separation via antibodies for specific markers can be by negative or positive selection procedures. In negative separation, antibodies are used which are specific for markers present on undesired cells, in this case tumor cells, which exhibit, for example, the tumor specific cell surface marker. Cells bound by an antibody to such a cell surface marker can be removed or lysed and the remaining desired mixture retained. In positive separation, antibodies specific for markers present on the desired cells of interest, in this case tumor-like cells, are used. Cells bound by the antibody are separated and retained. It will be understood that positive and negative separations can be used substantially simultaneously or in a sequential manner.

A common technique for antibody based separation is the use of flow cytometry such as by a fluorescence activated cell sorter (FACS). Typically, separation by flow cytometry is performed as follows. The suspended mixture of cells are centrifuged and resuspended in media. Antibodies which are conjugated to fluorochrome are added to allow the binding of the antibodies to specific cell surface markers. The cell mixture is then washed by one or more centrifugation and resuspension steps. The mixture is run through a FACS which separates the cells based on different fluorescence characteristics. FACS systems are available in varying levels of performance and ability, including multi-color analysis. The facilitating cell can be identified by a characteristic profile of forward and side scatter which is

influenced by size and granularity, as well as by positive and/or negative expression of certain cell surface markers.

Other separation techniques besides flow cytometry can also provide fast separations. One such method is
5 biotin-avidin based separation by affinity chromatography. Typically, such a technique is performed by incubating cells with biotin-coupled antibodies to specific markers, such as, for example, the transmembrane protein encoded by the tumor-specific marker, followed by passage through an avidin
10 column. Biotin-antibody-cell complexes bind to the column via the biotin-avidin interaction, while other cells pass through the column. The specificity of the biotin-avidin system is well suited for rapid positive separation. Multiple passages can ensure separation of a sufficient
15 level of the tumor cell subpopulation of interest.

In instances whereby the goal of the separation technique is to deplete the overall number of cells belonging to the tumor cell subpopulation, the cells derived from the starting source of cells which has now been
20 effectively depleted of tumor cells can be reintroduced into the patient. Such a depletion of the tumor cell subpopulation results in the amelioration of cancerous disorders associated with tumor progression.

In instances whereby the goal of the separation
25 technique is to augment or increase the overall number of cells belonging to a non-cancerous cell subpopulation, cells derived from the purified normal cell subpopulation can be reintroduced into the patient, thus resulting in the amelioration of cancerous disorders associated with an under
30 activity of the normal cell subpopulation.

The cells to be reintroduced will be cultured and expanded ex vivo prior to reintroduction. Purified normal cell subpopulation cells can be washed, suspended in, for

example, buffered saline, and reintroduced into the patient via intravenous administration.

Cells to be expanded can be cultured, using standard procedures, in the presence of an appropriate expansion agent which induces proliferation of the purified normal cell subpopulation. Such an expansion agent can, for example, be any appropriate cytokine, antigen, or antibody.

Prior to being reintroduced into a patient, the purified normal cells can be modified by, for example, transformation with gene sequences encoding gene products of interest. Such gene products should represent products which enhance the activity of the purified normal cell subpopulation or, alternatively, represent products which repress the activity of one or more of the other normal cell subpopulations. Cell transformation and gene expression procedures are well known to those of skill in the art, and can be as those described, above, in Section 5.2.

Well-known targeting methods can, additionally, be utilized in instances wherein the goal is to deplete the number of cells belonging to a specific tumor cell subpopulation. Such targeting methods can be in vivo or in vitro, and can involve the introduction of targeting agents into a population of cells such that the targeting agents selectively destroy a specific subset of the cells within the population. In vivo administration techniques which can be followed for such targeting agents are described, below, in Section 5.10.

Targeting agents generally comprise, first, a targeting moiety which, in the current instance, causes the targeting agent to selectively associate with a specific tumor cell subpopulation. The targeting agents generally comprise, second, a moiety capable of destroying a cell with which the targeting agent has become associated.

Targeting moieties can include, but are not limited to, antibodies directed to cell surface markers found specifically on the tumor cell subpopulation being targeted, or, alternatively, to ligands, such as growth factors, which bind receptor-type molecules found exclusively on the targeted tumor cell subpopulation.

Destructive moieties include any moiety capable of inactivating or destroying a cell to which the targeting agent has become bound. For example, a destructive moiety can include, but it is not limited to cytotoxins or radioactive agents. Cytotoxins include, for example, plant-, fungus-, or bacteria-derived toxins, with deglycosylated Ricin A chain toxins being generally preferred due to their potency and lengthy half-lives.

5.10. PHARMACEUTICAL PREPARATIONS AND METHODS OF ADMINISTRATION

The identified compounds that inhibit target gene expression, synthesis and/or activity can be administered to a patient at therapeutically effective doses to prevent, treat or ameliorate tumor progression. A therapeutically effective dose refers to that amount of the compound sufficient to result in amelioration of symptoms of tumor progression.

5.10.1. EFFECTIVE DOSE

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as

the ratio LD_{50}/ED_{50} . Compounds which exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects can be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC_{50} (i.e., the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma can be measured, for example, by high performance liquid chromatography.

25 5.10.2. FORMULATIONS AND USE

Pharmaceutical compositions for use in accordance with the present invention can be formulated in conventional manner using one or more physiologically acceptable carriers or excipients.

30 Thus, the compounds and their physiologically acceptable salts and solvates can be formulated for administration by inhalation or insufflation (either through

the mouth or the nose) or oral, buccal, parenteral or rectal administration.

For oral administration, the pharmaceutical compositions can take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets can be coated by methods well known in the art. Liquid preparations for oral administration can take the form of, for example, solutions, syrups or suspensions, or they can be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations can be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations can also contain buffer salts, flavoring, coloring and sweetening agents as appropriate.

Preparations for oral administration can be suitably formulated to give controlled release of the active compound.

For buccal administration the compositions can take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable
5 propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges
10 of e.g., gelatin for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

The compounds can be formulated for parenteral administration by injection, e.g., by bolus injection or
15 continuous infusion. Formulations for injection can be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions can take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and can contain
20 formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient can be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

The compounds can also be formulated in rectal
25 compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

In addition to the formulations described previously, the compounds can also be formulated as a depot
30 preparation. Such long acting formulations can be administered by implantation (for example, subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds can be formulated with suitable

polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

5 The compositions can, if desired, be presented in a pack or dispenser device which can contain one or more unit dosage forms containing the active ingredient. The pack can for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device can be
10 accompanied by instructions for administration.

5.11. DIAGNOSIS OF TUMOR PROGRESSION

A variety of methods can be employed for the diagnosis of tumor progression and of disorders involving tumor progression, including metastatic diseases. Such
15 methods can, for example, utilize reagents such as fingerprint gene nucleotide sequences described in Sections 5.2.1, and antibodies directed against differentially expressed and pathway gene peptides, as described, above, in Section 5.2.1 (peptides) and 5.2.3 (antibodies).
20 Specifically, such reagents can be used, for example, for the detection of the presence of target gene mutations, or the detection of either over or under expression of target gene in RNA.

The methods described herein can be performed, for
25 example, by utilizing pre-packaged diagnostic kits comprising at least one specific fingerprint gene nucleic acid or anti-fingerprint gene antibody reagent described herein, which can be conveniently used, e.g., in clinical settings, to diagnose patients exhibiting symptoms of
30 metastatic diseases.

Any cell type or tissue, preferably T-cells, in which the fingerprint gene is expressed can be utilized in the diagnostics described below.

5.11.1. DETECTION OF FINGERPRINT GENE NUCLEIC ACIDS

5 DNA or RNA from the cell type or tissue to be analyzed can easily be isolated using procedures which are well known to those in the art. Diagnostic procedures can also be performed "in situ" directly upon tissue sections
10 (fixed and/or frozen) of patient tissue obtained from biopsies or resections, such that no nucleic acid purification is necessary. Nucleic acid reagents such as those described in Section 5.1 can be used as probes and/or primers for such in situ procedures (see, for example,
15 Nuovo, G.J., 1992, PCR in situ hybridization: Protocols and Applications, Raven Press, NY).

Fingerprint gene nucleotide sequences, either RNA or DNA, can, for example, be used in hybridization or amplification assays of biological samples to detect gene
20 structures and expression associated with metastasis. Such assays can include, but are not limited to, Southern or Northern analyses, single stranded conformational polymorphism analyses, in situ hybridization assays, and, polymerase chain reaction analyses. Such analyses can
25 reveal both quantitative aspects of the expression pattern of the fingerprint gene, and qualitative aspects of the fingerprint gene expression and/or gene composition. That is, such techniques can include, for example, point mutations, insertions, deletions, chromosomal
30 rearrangements, and/or activation or inactivation of gene expression.

4686250 "24429880

Preferred diagnostic methods for the detection of fingerprint gene-specific nucleic acid molecules can involve for example, contacting and incubating nucleic acids, derived from the cell type or tissue being analyzed, with one or more labeled nucleic acid reagents as are described in Section 5.1, under conditions favorable for the specific annealing of these reagents to their complementary sequences within the nucleic acid molecule of interest. Preferably, the lengths of these nucleic acid reagents are at least 15 to 30 nucleotides. After incubation, all non-annealed nucleic acids are removed from the nucleic acid:fingerprint RNA molecule hybrid. The presence of nucleic acids from the target tissue which have hybridized, if any such molecules exist, is then detected. Using such a detection scheme, the nucleic acid from the tissue or cell type of interest can be immobilized, for example, to a solid support such as a membrane, or a plastic surface such as that on a microtitre plate or polystyrene beads. In this case, after incubation, non-annealed, labeled fingerprint nucleic acid reagents of the type described in Section 5.1 are easily removed. Detection of the remaining, annealed, labeled nucleic acid reagents is accomplished using standard techniques well-known to those in the art.

Alternative diagnostic methods for the detection of fingerprint gene specific nucleic acid molecules can involve their amplification, e.g., by PCR (the experimental embodiment set forth in Mullis, K.B., 1987, U.S. Patent No. 4,683,202), ligase chain reaction (Barany, F., 1991, Proc. Natl. Acad. Sci. USA 88:189-193), self sustained sequence replication (Guatelli, J.C. et al., 1990, Proc. Natl. Acad. Sci. USA 87:1874-1878), transcriptional amplification system (Kwoh, D.Y et al., 1989, Proc. Natl. Acad. Sci. USA 86:1173-1177), Q-Beta Replicase (Lizardi, P.M. et al., 1988,

268250" 24429880
Bio/Technology 6:1197), or any other nucleic acid
amplification method, followed by the detection of the
amplified molecules using techniques well known to those of
skill in the art. These detection schemes are especially
5 useful for the detection of nucleic acid molecules if such
molecules are present in very low numbers.

In one embodiment of such a detection scheme, a cDNA
molecule is obtained from an RNA molecule of interest (e.g.,
by reverse transcription of the RNA molecule into cDNA).
10 Cell types or tissues from which such RNA can be isolated
include any tissue in which wild type fingerprint gene is
known to be expressed. A sequence within the cDNA is then
used as the template for a nucleic acid amplification
reaction, such as a PCR amplification reaction, or the like.
15 The nucleic acid reagents used as synthesis initiation
reagents (e.g., primers) in the reverse transcription and
nucleic acid amplification steps of this method are chosen
from among the fingerprint gene nucleic acid reagents
described in Section 5.1. The preferred lengths of such
20 nucleic acid reagents are at least 19-30 nucleotides. For
detection of the amplified product, the nucleic acid
amplification can be performed using radioactively or non-
radioactively labeled nucleotides. Alternatively, enough
amplified product can be made such that the product can be
25 visualized by standard ethidium bromide staining or by
utilizing any other suitable nucleic acid staining method.

In addition to methods which focus primarily on the
detection of one nucleic acid sequence, fingerprint
profiles, as discussed in Section 5.3.4., can also be
30 assessed in such detection schemes. Fingerprint profiles
can be generated, for example, by utilizing a differential
display procedure, as discussed above in 5.1.1.2, Northern
analysis and/or RT-PCR. Any of the gene sequences

described, above, in Section 5.2.1 can be used as probes and/or PCR primers for the generation and corroboration of such fingerprint profiles.

5.11.2. DETECTION OF TARGET GENE PEPTIDES

5 Antibodies directed against wild type or mutant fingerprint gene peptides, which are discussed, above, in Section 5.2.3, can also be used in tumor progression diagnostics and prognostics, as described, for example, herein. Such diagnostic methods, can be used to detect
10 abnormalities in the level of fingerprint gene protein expression, or abnormalities in the structure and/or tissue, cellular, or subcellular location of fingerprinting gene protein. Structural differences can include, for example, differences in the size, electronegativity, or antigenicity
15 of the mutant fingerprint gene protein relative to the normal fingerprint gene protein.

Protein from the tissue or cell type to be analyzed can easily be isolated using techniques which are well known to those of skill in the art. The protein isolation methods
20 employed herein can, for example, be such as those described in Harlow and Lane (Harlow, E. and Lane, D., 1988, "Antibodies: A Laboratory Manual," Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York), which is incorporated herein by reference in its entirety.

25 Preferred diagnostic methods for the detection of wild type or mutant fingerprint gene peptide molecules can involve, for example, immunoassays wherein fingerprint gene peptides are detected by their interaction with an anti-fingerprint gene specific peptide antibody.

30 For example, antibodies, or fragments of antibodies, such as those described, above, in Section 5.2.3, useful in the present invention can be used to quantitatively or

qualitatively detect the presence of wild type or mutant fingerprint gene peptides. This can be accomplished, for example, by immunofluorescence techniques employing a fluorescently labeled antibody (see below) coupled with
5 light microscopic, flow cytometric, or fluorimetric detection. Such techniques are especially preferred if the fingerprint gene peptides are expressed on the cell surface.

The antibodies (or fragments thereof) useful in the present invention can, additionally, be employed
10 histologically, as in immunofluorescence or immunoelectron microscopy, for in situ detection of target gene peptides. in situ detection can be accomplished by removing a histological specimen from a patient, and applying thereto a labeled antibody of the present invention. The antibody (or
15 fragment) is preferably applied by overlaying the labeled antibody (or fragment) onto a biological sample. Through the use of such a procedure, it is possible to determine not only the presence of the fingerprint gene peptides, but also their distribution in the examined tissue. Using the
20 present invention, those of ordinary skill will readily perceive that any of a wide variety of histological methods (such as staining procedures) can be modified in order to achieve such in situ detection.

Immunoassays for wild type or mutant fingerprint
25 gene peptides typically comprise incubating a biological sample, such as a biological fluid, a tissue extract, freshly harvested cells, or cells which have been incubated in tissue culture, in the presence of a detectably labeled antibody capable of identifying fingerprint gene peptides,
30 and detecting the bound antibody by any of a number of techniques well-known in the art.

The biological sample can be brought in contact with and immobilized onto a solid phase support or carrier such

as nitrocellulose, or other solid support which is capable of immobilizing cells, cell particles or soluble proteins. The support can then be washed with suitable buffers followed by treatment with the detectably labeled

5 fingerprint gene specific antibody. The solid phase support can then be washed with the buffer a second time to remove unbound antibody. The amount of bound label on solid support can then be detected by conventional means.

By "solid phase support or carrier" is intended any
10 support capable of binding an antigen or an antibody. Well-known supports or carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite. The nature of the carrier can be either
15 soluble to some extent or insoluble for the purposes of the present invention. The support material can have virtually any possible structural configuration so long as the coupled molecule is capable of binding to an antigen or antibody. Thus, the support configuration can be spherical, as in a
20 bead, or cylindrical, as in the inside surface of a test tube, or the external surface of a rod. Alternatively, the surface can be flat such as a sheet, test strip, etc. Preferred supports include polystyrene beads. Those skilled in the art will know many other suitable carriers for
25 binding antibody or antigen, or will be able to ascertain the same by use of routine experimentation.

The binding activity of a given lot of anti-wild type or mutant fingerprint gene peptide antibody can be determined according to well-known methods. Those skilled
30 in the art will be able to determine operative and optimal assay conditions for each determination by employing routine experimentation.

One of the ways in which the fingerprint gene peptide-specific antibody can be detectably labeled is by linking the same to an enzyme and use in an enzyme immunoassay (EIA) (Voller, A., "The Enzyme Linked Immunosorbent Assay (ELISA)," *Diagnostic Horizons* 2:1-7, 1978) (Microbiological Associates Quarterly Publication, Walkersville, MD); Voller, A. et al., *J. Clin. Pathol.* 31:507-520 (1978); Butler, J.E., *Meth. Enzymol.* 73:482-523 (1981); Maggio, E. (ed.), *ENZYME IMMUNOASSAY*, CRC Press, Boca Raton, FL, 1980; Ishikawa, E. et al., (eds.) *ENZYME IMMUNOASSAY*, Kaku Shoin, Tokyo, 1981). The enzyme which is bound to the antibody will react with an appropriate substrate, preferably a chromogenic substrate, in such a manner as to produce a chemical moiety which can be detected, for example, by spectrophotometric, fluorimetric or by visual means. Enzymes which can be used to detectably label the antibody include, but are not limited to, malate dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate, dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase and acetylcholinesterase. The detection can be accomplished by colorimetric methods which employ a chromogenic substrate for the enzyme. Detection can also be accomplished by visual comparison of the extent of enzymatic reaction of a substrate in comparison with similarly prepared standards.

Detection can also be accomplished using any of a variety of other immunoassays. For example, by radioactively labeling the antibodies or antibody fragments, it is possible to detect fingerprint gene wild type or mutant peptides through the use of a radioimmunoassay (RIA)

46550" 24429880
(see, for example, Weintraub, B., *Principles of Radioimmunoassays*, Seventh Training Course on Radioligand Assay Techniques, The Endocrine Society, March, 1986, which is incorporated by reference herein). The radioactive
5 isotope can be detected by such means as the use of a gamma counter or a scintillation counter or by autoradiography.

It is also possible to label the antibody with a fluorescent compound. When the fluorescently labeled antibody is exposed to light of the proper wave length, its
10 presence can then be detected due to fluorescence. Among the most commonly used fluorescent labeling compounds are fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthalaldehyde and fluorescamine.

15 The antibody can also be detectably labeled using fluorescence emitting metals such as ^{152}Eu , or others of the lanthanide series. These metals can be attached to the antibody using such metal chelating groups as diethylenetriaminepentacetic acid (DTPA) or
20 ethylenediaminetetraacetic acid (EDTA).

The antibody also can be detectably labeled by coupling it to a chemiluminescent compound. The presence of the chemiluminescent-tagged antibody is then determined by detecting the presence of luminescence that arises during
25 the course of a chemical reaction. Examples of particularly useful chemiluminescent labeling compounds are luminol, isoluminol, theromatic acridinium ester, imidazole, acridinium salt and oxalate ester.

Likewise, a bioluminescent compound can be used to
30 label the antibody of the present invention. Bioluminescence is a type of chemiluminescence found in biological systems in, which a catalytic protein increases the efficiency of the chemiluminescent reaction. The

presence of a bioluminescent protein is determined by detecting the presence of luminescence. Important bioluminescent compounds for purposes of labeling are luciferin, luciferase and aequorin.

5

6. EXAMPLE: IDENTIFICATION AND CHARACTERIZATION OF A NOVEL GENE THAT INHIBITS TUMOR PROGRESSION

10 In the Example presented in this Section, the in vitro paradigm, described, above, in Section 5.1.1.1, was utilized to identify a gene, designated herein as the 030 gene, which is differentially expressed in cells with a high metastatic potential relative to cells having a low metastatic potential. Specifically, the 030 gene is expressed in high metastatic potential cells at a rate which
15 is many-fold lower than it is expressed in non-metastatic cells. Thus, as discussed below, the 030 gene can encode a product important to a number of neoplastic processes, including, for example, the progression of a cell to a metastatic state, the aggressiveness of a cell's metastatic
20 state, and the ability of a primary tumor cell to invade surrounding tissue. Given the differential 030 gene expression pattern revealed in this Section, the 030 gene product can represent a protein having tumor suppressor or inhibitor function.

25

6.1. MATERIALS AND METHODS

6.1.1. CELL CULTURE

B16 F1 and B16 F10 melanoma cell lines were maintained in culture in Eagle's minimal essential medium (MEM) supplemented with 10% fetal calf serum. Cells were
30 harvested from nonconfluent monolayers by a two minute treatment with 0.25% trypsin and 2mM EDTA.

For further characterization of in vivo activity, each cell line was injected into mice. Cells were washed two times in MEM, and the final cell suspension adjusted to 5×10^5 cells per ml in MEM. Two hundred microliters of this cell suspension (1×10^5 cells) was injected i.v. into the lateral tail vein of C57BL/6J mice. After three weeks, the mice were sacrificed and their lungs autopsied. The number of pulmonary tumors was determined by counting surface nodules using a dissecting microscope.

The differential expression of the 030 gene in B16 F1 relative to B16 F10 cell lines was compared with the extent of pulmonary metastases which developed in B16 F1-injected mice relative to B16 F10-injected mice.

6.1.2. Differential Display

Differential mRNA display was carried out as described, above, in Section 5.1.1.2. Details of the differential display are given, below.

RNA Isolation

RNA was isolated, using RNazol, from nonconfluent monolayers of B16 F1 and B16 F10 cell lines.

Isolated RNA was resuspended in DEPC H₂O and quantitated by spectrophotometry at OD₂₆₀. Approximately half of the RNA samples were then treated with DNase I to remove contaminating chromosomal DNA. Each 50 μ l RNA sample (50 μ g), 5.7 μ l 10x PCR buffer (Perkin-Elmer/Cetus) and 1 μ l RNase inhibitor (40 units/ μ l; Boehringer Mannheim, Germany) were mixed together. Two microliters of DNase I (10 units/ μ l; Boehringer Mannheim) was added to the reaction which was incubated for 30 min. at 37°C. The total volume was brought to 200 μ l with DEPC H₂O, extracted once with phenol/chloroform and precipitated by adding 20 μ l 3M NaOAc, pH 4.8, (DEPC-treated), 500 μ l absolute ETOH and incubated

for 1 hour on dry ice. The precipitated sample was centrifuged for 15 min., and the pellet was washed with 70% ETOH. The sample was re-centrifuged, the remaining liquid was aspirated, and the pellet was resuspended in 50 μ l H₂O.

5 The concentration of RNA was measured by reading the OD₂₆₀.

First strand cDNA synthesis

For each RNA sample, duplicate reverse transcription reactions were carried out in parallel. Four hundred ng RNA plus DEPC H₂O in a total volume of 10 μ l were added to 4 μ l
10 T₁₁CC 3' primer (10 μ M; Operon). The mixture was incubated at 70°C for 5 min. to denature the RNA and then placed at room temperature. Twenty-six μ l of reaction mix containing the following components was added to each denatured RNA/primer sample: 8 μ l 5x First Strand Buffer (Gibco/BRL,
15 Gaithersburg, MD), 4 μ l 0.1M DTT (Gibco/BRL), 2 μ l RNase inhibitor (40 units/ μ l) (Boehringer Mannheim), 4 μ l 200 μ M dNTP mix, 6 μ l H₂O, 2 μ l Superscript reverse transcriptase (200 units/ μ l; Gibco/BRL). The reactions were mixed gently and incubated for 30 min. at 42°C. Sixty μ l of H₂O, for a
20 final volume of 100 μ l, was then added and the samples were denatured for 5 min. at 85°C and stored at -20°C.

PCR reactions

The resulting single stranded cDNA molecules were then amplified by PCR. Specifically, 13 μ l of reaction mix
25 was added to each tube of a 96 well plate on ice. The reaction mix contained 6.4 μ l H₂O, 2 μ l 10x PCR Buffer (Perkin-Elmer), 2 μ l 20 μ M dNTPs, 0.4 μ l ³⁵S dATP (12.5 μ Ci/ μ l; 50 μ Ci total; Dupont/NEN), 2 μ l 5' primer OPE4 (5'GTGACATGCC-3'; 10 μ M; Operon), and 0.2 μ l AmpliTaq™
30 Polymerase (5 units/ μ l; Perkin-Elmer). Next, 2 μ l of 3' primer (T₁₁CC, 10 μ M) were added to the side of each tube, followed by 5 μ l of cDNA, also to the sides of the tubes, which were still on ice. Tubes were capped and mixed, and

brought up to 1000 rpm in a centrifuge, then immediately returned to ice. A Perkin-Elmer 9600 thermal cycler was used, and programmed as follows:

	94°C	2 min.	
5	*94°C	15 sec.	* = x 40
	*40°C	2 min.	
	*ramp 72°C	1 min.	
	*72°C	30 sec.	
	72°C	5 min.	
10	4°C	hold	

When the thermal cycler initially reached 94°C, the 96 well plate was removed from ice and placed directly into the cycler. Following the amplification reaction, 15 µl of loading dye, containing 80% formamide, 10 mM EDTA, 1 mg/ml xylene cyanole, 1 mg/ml bromphenol blue were added. The loading dye and reaction were mixed, incubated at 85°C for 5 min., cooled on ice, centrifuged, and placed on ice. Approximately 4 µl from each tube was loaded onto a pre-run (60V) 6% denaturing acrylamide gel. The gel was run at approximately 80V until top dye front was about 1 inch from bottom. The gel was transferred to 3MM paper (Whatman Paper, England) and dried under vacuum. Bands were visualized by autoradiography.

6.1.3. OTHER TECHNIQUES

25 Amplified cDNA Band Isolation and Amplification

PCR bands determined to be of interest in the differential display analysis were recovered from the gel and reamplified.

Briefly, differentially expressed bands were excised from the dried gel with a razor blade and placed into a

microfuge tube with 100 μ l H₂O and heated at 100°C for 5 min., vortexed, heated again to 100°C for 5 min., and vortexed again. After cooling, 100 μ l H₂O, 20 μ l 3M NaOAc, 1 μ l glycogen (20 mg/ml), and 500 μ l ethanol were added and
5 the sample was precipitated on dry ice. After centrifugation, the pellet was washed and resuspended in 10 μ l H₂O.

DNA isolated from the excised differentially expressed bands were then reamplified by PCR using the
10 following reaction conditions:

	58	μ l	H ₂ O
	10	μ l	10x PCR Buffer (see above)
	10	μ l	200 μ M dNTPs
	10	μ l	10 μ M 3' primer (see above)
15	10	μ l	10 μ M 5' primer (see above)
	1.5	μ l	amplified band
	0.5	μ l	AMPLITAQ® polymerase (5 units/ μ l; (Perkin Elmer)

PCR conditions were the same as the initial
20 conditions used to generate the original amplified band, as described, above. After reamplification, glycerol loading dyes were added and samples were loaded onto a 2% preparative TAE/Biogel (Bio101, La Jolla, CA) agarose gel and eluted. Bands were then excised from the gel with a
25 razor blade and vortexed for 15 min. at r.t., and purified using the MERMAID™ kit from Bio101 by adding 3 volumes of MERMAID™ high salt binding solution and 8 μ l of resuspended glassfog in a microfuge tube. Glassfog was then pelleted, washed 3 times with ethanol wash solution, and then DNA was
30 eluted twice in 10 μ l at 50°C.

Subcloning and Sequencing

The TA cloning kit (Invitrogen, San Diego, CA) was used to subclone the amplified bands. The ligation reaction typically consisted of 4 μ l sterile H₂O, 1 μ l ligation buffer, 2 μ l TA cloning vector, 2 μ l PCR product, and 1 μ l T4 DNA ligase. The volume of PCR product can vary, but the total volume of PCR product plus H₂O was always 6 μ l.

Ligations (including vector alone) were incubated overnight at 12°C before bacterial transformation. TA cloning kit

10 competent bacteria (INV α F': enda1, recA1, hsdR17(r-k, m+k), supE44, λ -, thi-1, gyrA, relA1, ϕ 80lacZ α M15 Δ (lacZYA-argF), deoR+, F') were thawed on ice and 2 μ l of 0.5 M β -mercaptoethanol were added to each tube. Two μ l from each ligation were added to each tube of competent cells (50 μ l),
15 mixed without vortexing, and incubated on ice for 30 min. Tubes were then placed in 42°C bath for exactly 30 sec., before being returned to ice for 2 min. Four hundred-fifty μ l of SOC media (Sambrook et al., 1989, supra) were then added to each tube which were then shaken at 37°C for 1 hr.
20 Bacteria were then pelleted, resuspended in approximately 200 μ l SOC and plated on Luria broth agar plates containing X-gal and 60 μ g/ml ampicillin and incubated overnight at 37°C. White colonies were then picked and screened for inserts using PCR.

25 A master mix containing 2 μ l 10 x PCR buffer, 1.6 μ l 2.5 mM dNTP's, 0.1 μ l 25 mM MgCl₂, 0.2 μ l M13 reverse primer (100 ng/ μ l), 0.2 μ l M13 forward primer (100 ng/ μ l), 0.1 μ l AmpliTaq® (Perkin-Elmer), and 15.8 μ l H₂O was made. Forty μ l of the master mix were aliquoted into tubes of a 96 well
30 plate, and whole bacteria were added with a pipette tip prior to PCR. The thermal cycler was programmed for insert screening as follows:

	94°C	2 min.	
	*94°C	15 sec.	* = x 35
	*47°C	2 min.	
	*ramp 72°C	30 sec.	
5	*72°C	30 sec.	
	72°C	10 min.	
	4°C	hold	

Reaction products were eluted on a 2% agarose gel and compared to vector control. Colonies with vectors containing inserts were purified by streaking onto LB/Amp plates. Vectors were isolated from such strains and subjected to sequence analysis, using an Applied Biosystems Automated Sequencer (Applied Biosystems, Inc. Seattle, WA).

Cloning of Human Gene

15 A human retina cDNA library obtained from Clontech was screened using the entire mouse fomy030 cDNA (FIGS. 3A and 3B) as a probe. During this screen, one million library phage were screened, 53 of which were found to hybridize with the mouse fomy030 probe. The cDNA inserts for eight of
20 these positives were isolated, subcloned, and sequenced.

Comparison of the murine fomy030 and human fohy030 sequences demonstrated a high degree of sequence similarity (86% identical at the nucleotide level and 94.4% identical at the amino acid level) within the 5', 1813 base pairs of
25 their cDNAs. However, beyond this point the sequences diverge and share no significant similarity. The sequence of fomy030 at the point of divergence is GTAG, which corresponds to a consensus splice donor site.

Three independent library isolated cDNAs, as well as
30 a cDNA isolated as a 3' RACE product were found to contain the fomy030 sequence. Thus, the most probable explanation for the divergence of the human and murine sequences is the

existence of alternate splice forms of the fomy030 and
fohy030 transcripts. The fomy030 splice version results in
a protein product of 542 amino acids in length, while the
fohy030 splice variant is predicted to encode a protein of
5 1497 amino acids in length (FIG. 5).

Another splice variant is shown in FIG. 6 (SEQ ID
NO:8), and encodes a protein of 1533 amino acids in length
(SEQ ID NO:9). The cDNA of FIG. 5 (SEQ ID NO:6) is missing
34 nucleotides beginning after 2879 in SEQ ID NO:8, and is
10 missing 74 nucleotides beginning after 2926 in SEQ ID NO:8.
Thus, nucleotides 2880-2892 in SEQ ID NO:6 are identical to
nucleotides 2914-2926 in SEQ ID NO:8, and the sequences are
essentially identical starting at 2893 in SEQ ID NO:6 and
3001 in SEQ ID NO:8. The difference in the respective amino
15 acid sequences is that the amino acids are identical from 1
to 844, and then again from 850 to 1497 in SEQ ID NO:7 and
from 886 to 1533 in SEQ ID NO:9.

Within their common 5' sequences, fohy030 was also
found to have an additional three base pairs (GGA) inserted
20 after position 1394 in the mouse cDNA (at positions 1066-
1068 in FIGS. 5 and 6). These additional three base pairs
fall within the open reading frames of both fohy030 and
fomy030, and result in an additional Glycine residue at
position 356 within the open reading frame of fohy030
25 relative to fomy030.

Northern Analysis

Northern analysis was performed to confirm the
differential expression of the genes corresponding to the
amplified bands, as described below.

30 Twelve micrograms of total RNA sample, 1.5 x RNA
loading dyes (60% formamide, 9% formaldehyde, 1.5 x MOPS,
.075% x C/BPB dyes) at a final concentration of 1 x and H₂O
to a final volume of 40 µl were mixed. The tubes were

heated at 65°C for 5 min. and then cooled on ice. The RNA samples analyzed were loaded onto a denaturing 1% agarose gel. The gel was run overnight at 32V in 1 x MOPS buffer.

A 300 ml denaturing 1% agarose gel was made as follows. Three grams of agarose (SeaKem™ LE, FMC BioProducts, Rockland, ME) and 60 ml of 5 x MOPS buffer (0.1M MOPS [pH 7.0], 40 mM NaOAc, 5mM EDTA [pH 8.0]) were added to 210 ml sterile H₂O. The mixture was heated until melted, then cooled to 50°C, at which time 5 µl ethidium bromide (5 mg/ml) and 30 ml of 37% formaldehyde were added to the melted gel mixture. The gel was swirled quickly to mix, and then poured immediately.

After electrophoresis, the gel was photographed with a fluorescent ruler, then was washed three times in DEPC H₂O, for 20 minutes per wash, at room temperature, with shaking. The RNA was then transferred from the gel to Hybond-N® membrane (Amersham), according to the methods of Sambrook *et al.*, 1989, *supra*, in 20 x SSC overnight.

The probes used to detect mRNA were typically synthesized as follows: 2 µl amplified cDNA band (~30 ng), 7 µl H₂O, and 2 µl 10 x Hexanucleotide mix (Boehringer-Mannheim) were mixed and heated to 95°C for 5 min., and then allowed to cool on ice. The volume of the amplified band can vary, but the total volume of the band plus H₂O was always 9 µl. 3 µl dATP/dGTP/dTTP mix (1:1:1 of 0.5 mM each), 5 µl α³²P dCTP 3000 Ci/mM (50 µCi total; Amersham, Arlington Heights, IL), and 1 µl Klenow (2 units; Boehringer-Mannheim) were mixed and incubated at 37°C. After 1 hr., 30 µl TE were added and the reaction was loaded onto a Biospin-6™ column (Biorad, Hercules, CA), and centrifuged. A 1 µl aliquot of eluate was used to measure incorporation in a scintillation counter with scintillant to ensure that 10⁶ cpm/µl of incorporation was achieved.

For pre-hybridization, the blot was placed into a roller bottle containing 10 ml of rapid-hyb solution (Amersham), and placed into 65°C incubator for at least 1 hr. For hybridization, 1×10^7 cpm of the probe was then heated to 95°C, chilled on ice, and added to 10 ml of rapid-hyb solution. The prehybridization solution was then replaced with probe solution and incubated for 16 hours at 65°C. The following day, the blot was washed once for 20 min. at room temperature in 2 x SSC/0.1% SDS and twice for 15 min. at 65°C in 0.1 x SSC/0.1% SDS before being covered in plastic wrap and put down for exposure.

In Situ Hybridization

10 μ m sections of formalin fixed/paraffin embedded benign nevi (non-metastatic growths of melanocytes) and malignant melanoma were post-fixed with 4% PFA/PBS for 15 minutes. After washing with PBS, sections were digested with 21 μ g/ml proteinase K at 37°C for 15 minutes, and again incubated with 4% PFA/PBS for 10 minutes. Sections were then washed with PBS, incubated with 0.2 N HCl for 10 minutes, washed with PBS, incubated with 0.25% acetic anhydride/1 M triethanolamine for 10 minutes, washed with PBS, and dehydrated with 70% ethanol and 100% ethanol.

Hybridizations were performed with 35 S-radiolabeled (5×10^7 cpm/ml) cRNA probes encoding a 1.1 kB segment of the coding region of the human cDNA (clone fohy030), and a 1 kB segment of the coding region of the human H4 histone gene in the presence of 50% formamide, 10% dextran sulfate, 1 x Denhardt's solution, 600 mM NaCl, 10 mM DTT, 0.25% SDS, and 100 μ g/ml tRNA for 18 hours at 55°C. The H4 histone gene was used as a control to show proper transcription of RNA.

After hybridization, slides were washed with 5 x SSC at 55°C, 50% formamide/2 x SSC at 55°C for 30 minutes, 10 mM Tris-HCl(pH 7.6)/500 mM NaCl/1 mM EDTA (TNE) at 37°C for 10

minutes, washed in TNE at 37°C for 10 minutes, incubated once in 2 x SSC at 50°C for 30 minutes, twice in 0.2 x SSC at 50°C for 30 minutes, and dehydrated with 70% ethanol and 100% ethanol. Localization of mRNA transcripts was detected by dipping slides in Kodak NBT-2 photo-emulsion and exposing for 4 days at 4°C. Controls for the *in situ* hybridization experiments included the use of a sense probe which showed no signal above backgrounds levels.

6.2. RESULTS

An in vitro paradigm, as described, above, in Section 5.1.1.1, was carried out using the melanoma cell lines, B16 F1 and B16 F10. The B16 F1 cell line exhibits a low metastatic potential, while the B16 F10 cell line exhibits a high metastatic potential. Thus, the two cell lines were grown in vitro as described in Section 6.1.1, RNA was isolated from these cells and differential display carried out as described in Section 6.1.

The differential display analysis identified a band, designated romy030, which represents a cDNA derived from RNA produced by a gene which was expressed at a much higher level in the B16 F1 cells, i.e., the low metastatic potential cells, relative to the gene's expression in B16 F10 cells, i.e., high metastatic potential cells. The gene corresponding to the romy030 band is referred to herein as the fomy030 or 030 gene.

The amplified romy030 band was isolated, reamplified, subcloned, and sequenced, as described, above, in Section 6.1.3. The romy030 nucleotide sequence (SEQ ID NO:1) is shown in FIG. 2.

A BLAST (Altschul, S.F. et al., 1990, J. Mol. Biol. 215:403-410) database search with the romy030 nucleotide sequence revealed no sequences within the database which are

similar to that of romy030. Thus, 030, the gene corresponding to romy030, appears to represent a novel, previously unknown gene which is differentially expressed in cells exhibiting a low metastatic potential relative to those cells exhibiting a high metastatic potential.

To confirm this putative differential regulation, amplified romy030 cDNA was used to probe Northern RNA blots containing RNA from B16 F1 and B16 F10 cells. FIG. 1 shows the results of one such Northern blot analysis, in which it is demonstrated that the steady state levels of fomy030 mRNA are significantly higher in the low metastatic potential cells (*i.e.*, the B16 F1 cells) relative to the high metastatic potential cells (*i.e.*, B16 F10 cells). Lanes 1 and 3 represent F1 cells and Lanes 2 and 4 represent F10 cells respectively. Thus, this Northern analysis confirmed the putative differential fomy030 regulation which had been suggested by the differential display results.

Two specific oligonucleotides were generated based on the sequence of romy030, romy030U 5'-GGGGAAGCACATCAAGGAAC-3' (SEQ ID NO:4) and romy030L 5'-GCAACTACACTCGGAAAAGC-3' (SEQ ID NO:5), for use in PCR reactions. cDNA libraries prepared from mRNA isolated from normal melanocytes and a mouse melanoma cell line were analyzed for the presence of fomy030 by PCR, utilizing the above romy030 probes. Fomy030 was detected in the melanocyte library but not in the melanoma library. The melanoma library was generated from a highly metastatic mouse melanoma K-1735 m2. This result is consistent the observation that fomy030 is present at reduced levels in the metastatic B16 F10 melanoma cell line. A radioactive DNA probe was generated from the subcloned romy030 DNA. This probe was used to screen the normal mouse melanocyte cDNA library. Three independent positive clones were identified

and isolated during this screening. These clones were designated fomy030a, fomy030b, and fomy030c. These cDNAs were sequenced and the overlapping portions were found to be identical. The nucleotide sequence of all three fomy030 cDNAs, designated as the fomy030 sequence (SEQ ID NO:2) is depicted in FIGS. 3A and 3B, and contains the sequence of romy030. The findings described herein suggest a novel role for fomy030 in tumor progression. A down-regulation of 030 can be used as a diagnostic marker for tumor progression, especially for the progression to metastasis. Further, 030 gene products can be used in the prevention and treatment of tumor progression disorders.

Fohy030 Expression in Human Tissue Samples

To determine whether the fohy030 gene product is differentially expressed in clinically relevant human disease, fohy030 gene expression was analyzed in biopsy sections of human benign nevi (non-metastatic growths of melanocytes) and malignant melanoma using *in situ* hybridization. Fohy030 expression was detected in small intermittent cells in the basal layer of the epidermis (likely, melanocytes) and in the majority of nevus cells in patients diagnosed with benign nevi. No expression of fohy030 was detected in the majority of melanoma cells in patients diagnosed with metastatic melanoma, though expression was detected in normal melanocytic cells in the same tissue section. These results show that the fohy030 gene product is associated with metastasis suppression.

6.3. 030 GENE EXPRESSION IS INVERSELY CORRELATED WITH METASTATIC POTENTIAL

6.3.1. EXPERIMENTAL PROTOCOLS AND RESULTS

The relationship between 030 gene expression and tumor progression was confirmed as described herein.

Specifically, the metastatic potentials of six variants of the B16 cell line were tested in animals and the metastatic potential was compared to the level of 030 gene expression observed within the cell variants.

5 A single cell suspension of B16 F1 cells (low metastatic potential) was injected intravenously into syngeneic C57BL/6 mice. After three weeks, lung tumors were excised and seeded into tissue culture. The following six cell lines were grown in culture: B16 G1, B16 G2, B16 G3,
10 B16 G4, B16 G9 and B16 G12.

 To test the metastatic ability of the above listed six tumor cell lines, the same number of cells for each of the six cell lines intravenously into different groups of syngeneic C57BL/6 mice. Three weeks later, the mice were
15 killed and the lungs were removed aseptically. Significantly more number of tumors were observed in mice injected with the following three cell lines: B16 G4, B16 G9 and B16 G12. These results demonstrate that the B16 G4, B16 G9 and B16 G12 cell lines have high metastatic potential
20 and the B16 G1, B16 G2 and B16 G3 cell lines have low metastatic potential.

 The lung tumors produced from these three highly metastatic cell lines (B16 G4, B16 G9 and B16 G12) were then excised and seeded into tissue culture to produce the
25 following four cell lines: B16 H5, B16,H6, B16 H7 and B16 H8.

 Northern analysis was performed to determine the expression of 030 gene in the above listed cell lines (i.e., B16 H5, B16,H6, B16 H7 and B16 H8) using procedures
30 described above in Section 6.1.3. Figure 4 shows the results of one such Northern blot analysis, in which it is demonstrated that the steady state levels of 030 mRNA are significantly lower in the highly metastatic cells (i.e.,

0883449 033397
B16 H5, B16,H6, B16 H7 and B16 H8) relative to the B16 F1 low metastatic potential cells. Lane 1 represents the B16 F1 cells, lane 2 is B16 F10 metastatic cells and lanes 3-6 represent B16 H5, B16,H6, B16 H7 and B16 H8.

5 Thus, this Northern analysis confirmed the initial finding in this invention that 030 expression is inversely related to the metastatic potential of tumor cells and supports the theory that the 030 gene product plays a role in inhibiting tumor progression, including the progression
10 to a high metastatic potential state. In this regard, it is important to note that the tumor cell number and homogeneity, and the syngeneic recipient did not change from one cell line to another in the above protocols. Therefore, the differences in metastatic incidence can only be
15 attributed to properties intrinsic to the various cell lines used. The clonal selection of tumors from successive metastases results in cells better capable of survival, formation and progression of tumor foci in the lung. This indicates that the decrease in expression of 030 observed in
20 the highly metastatic four cell lines (i.e., B16 H5, B16,H6, B16 H7 and B16 H8) is an intrinsic property of these cell lines and is related to the development, progression and metastatic potential of the tumor cells.

25 **7. EXAMPLE: USE OF FINGERPRINT GENES
 AS SURROGATE MARKERS IN CLINICAL TRIALS**

 The expression pattern of the fingerprint genes of the invention may be utilized as surrogate markers to monitor clinical human trials of drugs being tested for their efficacy as tumor progression treatments, or may,
30 additionally, be used to monitor patients undergoing clinical evaluation for the treatment of tumor progression. "Fingerprint gene," as used herein is defined as in Section

3, above. Individual fingerprint gene expression patterns may be analyzed or, alternatively, fingerprint patterns may be analyzed. "Fingerprint pattern," as used herein is defined as in Section 3, above.

5 The effect of the compound on the fingerprint gene expression normally displayed in connection with a disorder involving tumor progression can be used to evaluate the efficacy of the compound as a treatment for such a disorder. Additionally, fingerprint gene expression can be used to
10 monitor patients undergoing clinical evaluation for the treatment of the disorder.

 According to the invention, the fingerprint gene expression and fingerprint pattern derived from any of the paradigms described in Section 5.1.1.1 can be used to
15 monitor clinical trials of drugs in human patients. The paradigms described in Section 5.1.1.1, and illustrated in the Example presented in Section 6, above, for example, provide the fingerprint pattern of B16 melanoma cells. This profile gives an indicative reading, therefor, of the
20 metastatic and non-metastatic states of melanoma cells. Accordingly, the influence of anticancer chemotherapeutic agents on the melanoma cells can be measured by performing differential display on melanoma cells of patients undergoing clinical tests.

25 **7.1. TREATMENT OF PATIENTS AND PROCUREMENT
 OF TUMOR CELLS OR BIOPSIES**

 Test patients can be administered compounds suspected of antimetastatic activity. Control patients can be given a placebo.

30 Tumor cells or biopsies can be drawn from each patient after a determined period of treatment and RNA can be isolated as described in Section 6.6.1, above.

7.2. ANALYSIS OF SAMPLES

RNA can be subjected to differential display analysis as described in Section 6.6.1, above. A decrease in the metastatic potential of tumor cells is indicated by an increase in the intensity of the romy030 band, as described in Section 6.2, above.

8. DEPOSIT OF MICROORGANISMS

The following microorganism was deposited with the Agricultural Research Service Culture Collection (NRRL), Peoria, Illinois, on March 3, 1995 and assigned the indicated accession number:

<u>Microorganism</u>	<u>NRRL Accession No.</u>
<u>E. coli</u>	B-21416

OTHER EMBODIMENTS

The present invention is not to be limited in scope by the specific embodiments described which are intended as single illustrations of individual aspects of the invention and functionally equivalent methods and components are within the scope of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

174591.B11

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Shyjan, Andrew W.
- (ii) TITLE OF INVENTION: COMPOSITIONS AND METHODS FOR THE
DIAGNOSIS, PREVENTION AND TREATMENT OF TUMOR
PROGRESSION
- (iii) NUMBER OF SEQUENCES: 9
- (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Fish & Richardson P.C.
(B) STREET: 225 Franklin Street
(C) CITY: Boston
(D) STATE: MA
(E) COUNTRY: USA
(F) ZIP: 02110-2804
- (v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER: US
(B) FILING DATE: 29-MAR-1996
(C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: US 08/412,431
(B) FILING DATE: 29-MAR-1995
- (viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Fasse, J. Peter
(B) REGISTRATION NUMBER: 32,983
(C) REFERENCE/DOCKET NUMBER: 07334/004001
- (ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 617/542-5070
(B) TELEFAX: 617/542-8906
(C) TELEX: 200154

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 186 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GGTGCTGGAG TACCTCATGG GCGGTGCCTA CCGCTGCAAC TACACTCGGA AAAGCTTCOG	60
GACTCTCTAC AACAACTTGT TTGGCCCTAA GAGGGTAGAG CTCAGCAGAC ACACAGTGTC	120

CTGTGCCTCC CAGAGTAACA TGTGGTTCCT TGATGTGCTT CCCC AAAAGC CCACCTGTGC 180
AGAATG 186

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2729 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

AAGGAGGCTA GGCTGCACCC TTCCCGCTTG CTCAGCAGCT GAGGCAGGGT CAGAAAGCAT 60
GGATAGAGAA GACATTTTGC AAAAGGGAAT GCATCTTTGT AATTCCCAGT ACAAAGACC 120
CTAACAGATG TTGCTGTGGT CAGCTCACTA ACCAGCACAT CCCCCCTTG CCGAGTGGGG 180
CTCCCAGCAC AACAGGAGAG GACACCAAGC AGGCAGACAC GCAGTCCGGG AAATGGTCTG 240
TCAGCAAACA CACCCAGAGC TACCCAACAG ACTCCTATGG GATTCTTGAA TTCCAGGGTG 300
GGGGTTACTC CAATAAAGCC ATG TAC ATC CGA GTC TCC TAC GAC ACC AAG 350
Met Tyr Ile Arg Val Ser Tyr Asp Thr Lys
1 5 10
CCA GAT TCC CTG CTC CAC CTC ATG GTG AAG GAC TGG CAG CTG GAG CTC 398
Pro Asp Ser Leu Leu His Leu Met Val Lys Asp Trp Gln Leu Glu Leu
15 20 25
CCG AAG CTC TTG ATA TCT GTG CAC GGA GGC CTC CAA AGC TTC GAG ATG 446
Pro Lys Leu Leu Ile Ser Val His Gly Gly Leu Gln Ser Phe Glu Met
30 35 40
CAG TCC AAA CTG AAG CAG GTG TTT GGG AAA GGT CTG ATC AAG GCT GCC 494
Gln Ser Lys Leu Lys Gln Val Phe Gly Lys Gly Leu Ile Lys Ala Ala
45 50 55
ATG ACC ACG GGG GCG TGG ATC TTC ACC GGG GGT GTG AGC ACT GGT GTC 542
Met Thr Thr Gly Ala Trp Ile Phe Thr Gly Gly Val Ser Thr Gly Val
60 65 70
GTC AGC CAT GTG GGG GAT GCC TTG AAA GAC CAC TCC TCC AAG TCC AGA 590
Val Ser His Val Gly Asp Ala Leu Lys Asp His Ser Ser Lys Ser Arg
75 80 85 90
GGC CGG CTC TGT GCT ATA GGA ATT GCT CCC TGG GGC ATG GTG GAG AAC 638
Gly Arg Leu Cys Ala Ile Gly Ile Ala Pro Trp Gly Met Val Glu Asn
95 100 105
AAG GAA GAC CTG ATT GGA AAA GAT GTA ACA AGA GTC TAT CAG ACC ATG 686
Lys Glu Asp Leu Ile Gly Lys Asp Val Thr Arg Val Tyr Gln Thr Met
110 115 120
TCC AAC CCT CTG AGC AAG CTC TCT GTG CTC AAC AAT TCC CAC ACT CAC 734
Ser Asn Pro Leu Ser Lys Leu Ser Val Leu Asn Asn Ser His Thr His

125					130					135						
TTC	ATC	TTG	GCT	GAC	AAC	GGC	ACC	CTG	GGC	AAG	TAT	GGT	GCT	GAG	GTG	782
Phe	Ile	Leu	Ala	Asp	Asn	Gly	Thr	Leu	Gly	Lys	Tyr	Gly	Ala	Glu	Val	
140						145				150						
AAG	CTT	CGA	AGA	CAG	CTG	GAA	AAA	CAC	ATC	TCC	CTG	CAG	AAG	ATC	AAC	830
Lys	Leu	Arg	Arg	Gln	Leu	Glu	Lys	His	Ile	Ser	Leu	Gln	Lys	Ile	Asn	
155				160						165				170		
ACA	AGG	CTG	GGC	CAG	GGT	GTA	CCT	GTC	GTG	GGC	CTA	GTG	GTA	GAA	GGT	878
Thr	Arg	Leu	Gly	Gln	Gly	Val	Pro	Val	Val	Gly	Leu	Val	Val	Glu	Gly	
				175				180						185		
GGT	CCT	AAC	GTG	GTT	TCT	ATC	GTC	CTG	GAG	TAT	CTC	AAA	GAA	GAC	CCT	926
Gly	Pro	Asn	Val	Val	Ser	Ile	Val	Leu	Glu	Tyr	Leu	Lys	Glu	Asp	Pro	
		190						195				200				
CCT	GTC	CCT	GTG	GTG	GTT	TGC	GAT	GGC	AGT	GGA	CGT	GCC	TCT	GAC	ATT	974
Pro	Val	Pro	Val	Val	Val	Cys	Asp	Gly	Ser	Gly	Arg	Ala	Ser	Asp	Ile	
		205				210						215				
TTG	TCC	TTC	GCA	CAC	AAA	TAC	TGC	GAC	GAA	GGA	GGA	GTC	ATA	AAC	GAG	1022
Leu	Ser	Phe	Ala	His	Lys	Tyr	Cys	Asp	Glu	Gly	Gly	Val	Ile	Asn	Glu	
220						225				230						
TCC	CTG	CGG	GAC	CAG	CTT	CTA	GTT	ACC	ATT	CAG	AAA	ACA	TTT	AAT	TAC	1070
Ser	Leu	Arg	Asp	Gln	Leu	Leu	Val	Thr	Ile	Gln	Lys	Thr	Phe	Asn	Tyr	
235				240						245				250		
AGC	AAG	TCC	CAG	TCG	TAT	CAG	CTG	TTT	GCA	ATT	ATC	ATG	GAG	TGC	ATG	1118
Ser	Lys	Ser	Gln	Ser	Tyr	Gln	Leu	Phe	Ala	Ile	Ile	Met	Glu	Cys	Met	
				255				260						265		
AAG	AAG	AAA	GAA	CTC	GTC	ACT	GTG	TTT	CGG	ATG	GGT	TCC	GAG	GGT	CAG	1166
Lys	Lys	Lys	Glu	Leu	Val	Thr	Val	Phe	Arg	Met	Gly	Ser	Glu	Gly	Gln	
		270				275						280				
CAA	GAT	GTC	GAG	ATG	GCA	ATT	TTA	ACT	GCC	TTG	CTC	AAA	GGA	ACC	AAC	1214
Gln	Asp	Val	Glu	Met	Ala	Ile	Leu	Thr	Ala	Leu	Leu	Lys	Gly	Thr	Asn	
		285				290						295				
GCA	TCA	GCT	CCA	GAT	CAG	CTG	AGC	TTG	GCC	CTG	GCT	TGG	AAC	CGG	GTC	1262
Ala	Ser	Ala	Pro	Asp	Gln	Leu	Ser	Leu	Ala	Leu	Ala	Trp	Asn	Arg	Val	
300						305				310						
GAC	ATA	GCG	CGA	AGC	CAG	ATC	TTC	GTC	TTT	GGC	CCA	CAC	TGG	CCG	CCA	1310
Asp	Ile	Ala	Arg	Ser	Gln	Ile	Phe	Val	Phe	Gly	Pro	His	Trp	Pro	Pro	
315				320						325				330		
CTG	GGA	AGC	CTG	GCC	CCT	CCT	GTG	GAC	ACC	AAA	GCC	GCA	GAG	AAG	GAA	1358
Leu	Gly	Ser	Leu	Ala	Pro	Pro	Val	Asp	Thr	Lys	Ala	Ala	Glu	Lys	Glu	
				335				340						345		
AAG	AAG	CCA	CCC	ACA	GCC	ACC	ACC	AAG	GGG	AGA	GGA	AAA	GGA	AAA	GGC	1406
Lys	Lys	Pro	Pro	Thr	Ala	Thr	Thr	Lys	Gly	Arg	Gly	Lys	Gly	Lys	Gly	
		350				355						360				

AAG AAG AAA GGC AAA GTG AAA GAG GAA GTG GAG GAA GAG ACG GAC CCC Lys Lys Lys Gly Lys Val Lys Glu Glu Val Glu Glu Glu Thr Asp Pro 365 370 375	1454
CGG AAG CTT GAG CTG CTC AAC TGG GTG AAT GCC CTG GAG CAA GCC ATG Arg Lys Leu Glu Leu Leu Asn Trp Val Asn Ala Leu Glu Gln Ala Met 380 385 390	1502
CTG GAT GCT CTT GTC CTA GAT CGG GTG GAC TTT GTA AAG CTC CTG ATT Leu Asp Ala Leu Val Leu Asp Arg Val Asp Phe Val Lys Leu Leu Ile 395 400 405 410	1550
GAA AAC GGA GTG AAC ATG CAG CAT TTC CTC ACC ATC CCG AGG CTG GAG Glu Asn Gly Val Asn Met Gln His Phe Leu Thr Ile Pro Arg Leu Glu 415 420 425	1598
GAG CTA TAC AAC ACC AGA CTG GGC CCA CCA AAC ACC CTT CAT CTG CTG Glu Leu Tyr Asn Thr Arg Leu Gly Pro Pro Asn Thr Leu His Leu Leu 430 435 440	1646
GTG CGG GAT GTA AAG AAG AGC AAC CTT CCA CCT GAT TAC CAC ATC AGC Val Arg Asp Val Lys Lys Ser Asn Leu Pro Pro Asp Tyr His Ile Ser 445 450 455	1694
CTC ATT GAT ATA GGA CTG GTG CTG GAG TAC CTC ATG GGC GGT GCC TAC Leu Ile Asp Ile Gly Leu Val Leu Glu Tyr Leu Met Gly Gly Ala Tyr 460 465 470	1742
CGC TGC AAC TAC ACT CGG AAA AGC TTC CGG ACT CTC TAC AAC AAC TTG Arg Cys Asn Tyr Thr Arg Lys Ser Phe Arg Thr Leu Tyr Asn Asn Leu 475 480 485 490	1790
TTT GGC CCT AAG AGG GTA GAG CTC AGC AGA CAC ACA GTG TCC TGT GCC Phe Gly Pro Lys Arg Val Glu Leu Ser Arg His Thr Val Ser Cys Ala 495 500 505	1838
TCC CAG AGT AAC ATG TGG TTC CTT GAT GTG CTT CCC CAA AAG CCC ACC Ser Gln Ser Asn Met Trp Phe Leu Asp Val Leu Pro Gln Lys Pro Thr 510 515 520	1886
TGT GCA GAA TGC AAC TCT TCA CCT CAC CTG TCC CAA ACT GAC ATC ACC Cys Ala Glu Cys Asn Ser Ser Pro His Leu Ser Gln Thr Asp Ile Thr 525 530 535	1934
CCA CCT CTG CCC T GACACCCAGT GCAGGGCCTC CTAGCTTTCA CATGCAGCCA Pro Pro Leu Pro 540	1987
TTACATCGC CTCTCAAGAC TGGGCCAGGC AGTGCAACCT GTCAAGCATG TCTGTCCTCC	2047
CCTCCTTCCT ACAATAGCCC CCCCTCTGGG CCCCATGCCT CTGCTCTCTC AGCCC GTTCT	2107
CCTCCCCACT GATCACTGGC GCTCCTGTTG TCTTCCAAGG CAAGGAACAA GGAAAAGCAT	2167
CTTTTGTCCC ACAAAGTTT AGGGCTCCCC GCTGTTCAAC CATAGCCAAC CTCACTGTAC	2227
ATCGGAGTCA TCCAGGCCAG CTGCCACACA CAAGCCTTCC CCACCCTATC CCAATAGACC	2287

CTATTCCTCC ATCAAAATCA AAGCTAACTC CTGGCCTGCC ACATTGCTTC TTCTTGCTCC 2347
 AGCCTGTTAA ACCTCCAATA AATGTCAGAT CTGTGGGAAG CCTTCCTCAC TCTCACTCCA 2407
 CAGTTTGTAC AGAGAGCGAG AGCCTCGTTT GGTTCCTACTT ACAAGGAAGG CTTTGTGTCT 2467
 GTCTGTCCTT CCCAACTGAC TTCTGTTGAC AGAAGCAGTT TCCACATGAA AGCGTTGACT 2527
 CACCTGGATG TTGTCATTAA TTAATAGTGA TACAAAATAT TGACACTTCT TTTCCTGCTT 2587
 CTTTGTATG CAGCCGAAAG CACTTAAGCT TCTGGGAATG GAAGTAAGTA GGACATGTTT 2647
 GTGGCAGTTT ATTTACTATA TATACCTTTG TCATTCTGTG GAAGCAAAAA TTGCAATGTT 2707
 TTCCATGAAT AAAGCTCGTG CC 2729

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 542 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met Tyr Ile Arg Val Ser Tyr Asp Thr Lys Pro Asp Ser Leu Leu His
 1 5 10 15
 Leu Met Val Lys Asp Trp Gln Leu Glu Leu Pro Lys Leu Leu Ile Ser
 20 25 30
 Val His Gly Gly Leu Gln Ser Phe Glu Met Gln Ser Lys Leu Lys Gln
 35 40 45
 Val Phe Gly Lys Gly Leu Ile Lys Ala Ala Met Thr Thr Gly Ala Trp
 50 55 60
 Ile Phe Thr Gly Gly Val Ser Thr Gly Val Val Ser His Val Gly Asp
 65 70 75 80
 Ala Leu Lys Asp His Ser Ser Lys Ser Arg Gly Arg Leu Cys Ala Ile
 85 90 95
 Gly Ile Ala Pro Trp Gly Met Val Glu Asn Lys Glu Asp Leu Ile Gly
 100 105 110
 Lys Asp Val Thr Arg Val Tyr Gln Thr Met Ser Asn Pro Leu Ser Lys
 115 120 125
 Leu Ser Val Leu Asn Asn Ser His Thr His Phe Ile Leu Ala Asp Asn
 130 135 140
 Gly Thr Leu Gly Lys Tyr Gly Ala Glu Val Lys Leu Arg Arg Gln Leu
 145 150 155 160
 Glu Lys His Ile Ser Leu Gln Lys Ile Asn Thr Arg Leu Gly Gln Gly
 165 170 175

Val Pro Val Val Gly Leu Val Val Glu Gly Gly Pro Asn Val Val Ser
 180 185 190
 Ile Val Leu Glu Tyr Leu Lys Glu Asp Pro Pro Val Pro Val Val Val
 195 200 205
 Cys Asp Gly Ser Gly Arg Ala Ser Asp Ile Leu Ser Phe Ala His Lys
 210 215 220
 Tyr Cys Asp Glu Gly Gly Val Ile Asn Glu Ser Leu Arg Asp Gln Leu
 225 230 235 240
 Leu Val Thr Ile Gln Lys Thr Phe Asn Tyr Ser Lys Ser Gln Ser Tyr
 245 250 255
 Gln Leu Phe Ala Ile Ile Met Glu Cys Met Lys Lys Lys Glu Leu Val
 260 265 270
 Thr Val Phe Arg Met Gly Ser Glu Gly Gln Gln Asp Val Glu Met Ala
 275 280 285
 Ile Leu Thr Ala Leu Leu Lys Gly Thr Asn Ala Ser Ala Pro Asp Gln
 290 295 300
 Leu Ser Leu Ala Leu Ala Trp Asn Arg Val Asp Ile Ala Arg Ser Gln
 305 310 315 320
 Ile Phe Val Phe Gly Pro His Trp Pro Pro Leu Gly Ser Leu Ala Pro
 325 330 335
 Pro Val Asp Thr Lys Ala Ala Glu Lys Glu Lys Lys Pro Pro Thr Ala
 340 345 350
 Thr Thr Lys Gly Arg Gly Lys Gly Lys Gly Lys Lys Lys Gly Lys Val
 355 360 365
 Lys Glu Glu Val Glu Glu Glu Thr Asp Pro Arg Lys Leu Glu Leu Leu
 370 375 380
 Asn Trp Val Asn Ala Leu Glu Gln Ala Met Leu Asp Ala Leu Val Leu
 385 390 395 400
 Asp Arg Val Asp Phe Val Lys Leu Leu Ile Glu Asn Gly Val Asn Met
 405 410 415
 Gln His Phe Leu Thr Ile Pro Arg Leu Glu Glu Leu Tyr Asn Thr Arg
 420 425 430
 Leu Gly Pro Pro Asn Thr Leu His Leu Leu Val Arg Asp Val Lys Lys
 435 440 445
 Ser Asn Leu Pro Pro Asp Tyr His Ile Ser Leu Ile Asp Ile Gly Leu
 450 455 460
 Val Leu Glu Tyr Leu Met Gly Gly Ala Tyr Arg Cys Asn Tyr Thr Arg
 465 470 475 480
 Lys Ser Phe Arg Thr Leu Tyr Asn Asn Leu Phe Gly Pro Lys Arg Val

485

490

495

Glu Leu Ser Arg His Thr Val Ser Cys Ala Ser Gln Ser Asn Met Trp
500 505 510

Phe Leu Asp Val Leu Pro Gln Lys Pro Thr Cys Ala Glu Cys Asn Ser
515 520 525

Ser Pro His Leu Ser Gln Thr Asp Ile Thr Pro Pro Leu Pro
530 535 540

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GGGGAAGCAC ATCAAGGAAC

20

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GCAACTACTA CACTCGGAAA AGC

23

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 4944 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 346..4837

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

ACTCATTATA GGGNTCGAGC GGCCGCCCGG GCAGGTTTGA GCTGTGCCCT CTCCATTCCA 60

CTGCTGTGGC AGGGTCAGAA ATCTTGATA GAGAAAACCT TTTGCAAACG GGAATGTATC 120

TTTGTAAATC CTAGCACGAA AGACTCTAAC AGGTGTTGCT GTGGCCAGTT CACCAACCAG	180
CATATCCCCC CTCTGCCAAG TGCAACACCC AGCAAAAATG AAGAGGAAAG CAAACAGGTG	240
GAGACTCAGC CTGAGAAATG GTCTGTTGCC AAGCACACCC AGAGCTACCC AACAGATTCC	300
TATGGAGTTC TTGAATTCCA GGGTGGCGGA TATTCCAATA AAGCC ATG TAT ATC	354
Met Tyr Ile	
1	
CGT GTA TCC TAT GAC ACC AAG CCA GAC TCA CTG CTC CAT CTC ATG GTG	402
Arg Val Ser Tyr Asp Thr Lys Pro Asp Ser Leu Leu His Leu Met Val	
5 10 15	
AAA GAT TGG CAG CTG GAA CTC CCC AAG CTC TTA ATA TCT GTG CAT GGA	450
Lys Asp Trp Gln Leu Glu Leu Pro Lys Leu Leu Ile Ser Val His Gly	
20 25 30 35	
GGC CTC CAG AAC TTT GAG ATG CAG CCC AAG CTG AAA CAA GTC TTT GGG	498
Gly Leu Gln Asn Phe Glu Met Gln Pro Lys Leu Lys Gln Val Phe Gly	
40 45 50	
AAA GGC CTG ATC AAG GCT GCT ATG ACC ACC GGG GCC TGG ATC TTC ACC	546
Lys Gly Leu Ile Lys Ala Ala Met Thr Thr Gly Ala Trp Ile Phe Thr	
55 60 65	
GGG GGT GTC AGC ACA GGT GTT ATC AGC CAC GTA GGG GAT GCC TTG AAA	594
Gly Gly Val Ser Thr Gly Val Ile Ser His Val Gly Asp Ala Leu Lys	
70 75 80	
GAC CAC TCC TCC AAG TCC AGA GGC CGG GTT TGT GCT ATA GGA ATT GCT	642
Asp His Ser Ser Lys Ser Arg Gly Arg Val Cys Ala Ile Gly Ile Ala	
85 90 95	
CCA TGG GGC ATC GTG GAG AAT AAG GAA GAC CTG GTT GGA AAG GAT GTA	690
Pro Trp Gly Ile Val Glu Asn Lys Glu Asp Leu Val Gly Lys Asp Val	
100 105 110 115	
ACA AGA GTG TAC CAG ACC ATG TCC AAC CCT CTA AGT AAG CTC TCT GTG	738
Thr Arg Val Tyr Gln Thr Met Ser Asn Pro Leu Ser Lys Leu Ser Val	
120 125 130	
CTC AAC AAC TCC CAC ACC CAC TTC ATC CTG GCT GAC AAT GGC ACC CTG	786
Leu Asn Asn Ser His Thr His Phe Ile Leu Ala Asp Asn Gly Thr Leu	
135 140 145	
GGC AAG TAT GGC GCC GAG GTG AAG CTG CGA AGG CTG CTG GAA AAG CAC	834
Gly Lys Tyr Gly Ala Glu Val Lys Leu Arg Arg Leu Leu Glu Lys His	
150 155 160	
ATC TCC CTC CAG AAG ATC AAC ACA AGA CTG GGG CAG GGC GTG CCC CTC	882
Ile Ser Leu Gln Lys Ile Asn Thr Arg Leu Gly Gln Gly Val Pro Leu	
165 170 175	
GTG GGT CTC GTG GTG GAG GGG GGC CCT AAC GTG GTG TCC ATC GTC TTG	930
Val Gly Leu Val Val Glu Gly Gly Pro Asn Val Val Ser Ile Val Leu	
180 185 190 195	

GAA TAC CTG CAA GAA GAG CCT CCC ATC CCT GTG GTG ATT TGT GAT GGC	978
Glu Tyr Leu Gln Glu Glu Pro Pro Ile Pro Val Val Ile Cys Asp Gly	
200 205 210	
AGC GGA CGT GCC TCG GAC ATC CTG TCC TTT GCG CAC AAG TAC TGT GAA	1026
Ser Gly Arg Ala Ser Asp Ile Leu Ser Phe Ala His Lys Tyr Cys Glu	
215 220 225	
GAA GGC GGA ATA ATA AAT GAG TCC CTC AGG GAG CAG CTT CTA GTT ACC	1074
Glu Gly Gly Ile Ile Asn Glu Ser Leu Arg Glu Gln Leu Leu Val Thr	
230 235 240	
ATT CAG AAA ACA TTT AAT TAT AAT AAG GCA CAA TCA CAT CAG CTG TTT	1122
Ile Gln Lys Thr Phe Asn Tyr Asn Lys Ala Gln Ser His Gln Leu Phe	
245 250 255	
GCA ATT ATA ATG GAG TGC ATG AAG AAG AAA GAA CTC GTC ACT GTG TTC	1170
Ala Ile Ile Met Glu Cys Met Lys Lys Lys Glu Leu Val Thr Val Phe	
260 265 270 275	
AGA ATG GGT TCT GAG GGC CAG CAG GAC ATC GAG ATG GCA ATT TTA ACT	1218
Arg Met Gly Ser Glu Gly Gln Gln Asp Ile Glu Met Ala Ile Leu Thr	
280 285 290	
GCC CTG CTG AAA GGA ACA AAC GTA TCT GCT CCA GAT CAG CTG AGC TTG	1266
Ala Leu Leu Lys Gly Thr Asn Val Ser Ala Pro Asp Gln Leu Ser Leu	
295 300 305	
GCA CTG GCT TGG AAC CGC GTG GAC ATA GCA CGA AGC CAG ATC TTT GTC	1314
Ala Leu Ala Trp Asn Arg Val Asp Ile Ala Arg Ser Gln Ile Phe Val	
310 315 320	
TTT GGG CCC CAC TGG ACG CCC CTG GGA AGC CTG GCA CCC CCG ACG GAC	1362
Phe Gly Pro His Trp Thr Pro Leu Gly Ser Leu Ala Pro Pro Thr Asp	
325 330 335	
AGC AAA GCC ACG GAG AAG GAG AAG AAG CCA CCC ATG GCC ACC ACC AAG	1410
Ser Lys Ala Thr Glu Lys Glu Lys Lys Pro Pro Met Ala Thr Thr Lys	
340 345 350 355	
GGA GGA AGA GGA AAA GGG AAA GGC AAG AAG AAA GGG AAA GTG AAA GAG	1458
Gly Gly Arg Gly Lys Gly Lys Gly Lys Lys Lys Gly Lys Val Lys Glu	
360 365 370	
GAA GTG GAG GAA GAA ACT GAC CCC CGG AAG ATA GAG CTG CTG AAC TGG	1506
Glu Val Glu Glu Glu Thr Asp Pro Arg Lys Ile Glu Leu Leu Asn Trp	
375 380 385	
GTG AAT GCT TTG GAG CAA GCG ATG CTA GAT GCT TTA GTC TTA GAT CGT	1554
Val Asn Ala Leu Glu Gln Ala Met Leu Asp Ala Leu Val Leu Asp Arg	
390 395 400	
GTC GAC TTT GTG AAG CTC CTG ATT GAA AAC GGA GTG AAC ATG CAA CAC	1602
Val Asp Phe Val Lys Leu Leu Ile Glu Asn Gly Val Asn Met Gln His	
405 410 415	
TTT CTG ACC ATT CCG AGG CTG GAG GAG CTT TAT AAC ACA AGA CTG GGT	1650
Phe Leu Thr Ile Pro Arg Leu Glu Glu Leu Tyr Asn Thr Arg Leu Gly	

420	425	430	435	
CCA CCA AAC ACA CTT CAT CTG CTG GTG AGG GAT GTG AAA AAG AGC AAC Pro Pro Asn Thr Leu His Leu Leu Val Arg Asp Val Lys Lys Ser Asn 440 445 450				1698
CTT CCG CCT GAT TAC CAC ATC AGC CTC ATA GAC ATC GGG CTC GTG CTG Leu Pro Pro Asp Tyr His Ile Ser Leu Ile Asp Ile Gly Leu Val Leu 455 460 465				1746
GAG TAC CTC ATG GGA GGA GCC TAC CGC TGC AAC TAC ACT CGG AAA AAC Glu Tyr Leu Met Gly Gly Ala Tyr Arg Cys Asn Tyr Thr Arg Lys Asn 470 475 480				1794
TTT CGG ACC CTT TAC AAC AAC TTG TTT GGA CCA AAG AGG CCT AAA GCT Phe Arg Thr Leu Tyr Asn Asn Leu Phe Gly Pro Lys Arg Pro Lys Ala 485 490 495				1842
CTT AAA CTT CTG GGA ATG GAA GAT GAT GAG CCT CCA GCT AAA GGG AAG Leu Lys Leu Leu Gly Met Glu Asp Asp Glu Pro Pro Ala Lys Gly Lys 500 505 510 515				1890
AAA AAA AAA AAA AAG AAA AAG GAG GAA GAG ATC GAC ATT GAT GTG GAC Lys Lys Lys Lys Lys Lys Lys Glu Glu Glu Ile Asp Ile Asp Val Asp 520 525 530				1938
GAC CCT GCC GTG AGT CGG TTC CAG TAT CCC TTC CAC GAG CTG ATG GTG Asp Pro Ala Val Ser Arg Phe Gln Tyr Pro Phe His Glu Leu Met Val 535 540 545				1986
TGG GCA GTG CTG ATG AAA CGC CAG AAA ATG GCA GTG TTC CTC TGG CAG Trp Ala Val Leu Met Lys Arg Gln Lys Met Ala Val Phe Leu Trp Gln 550 555 560				2034
CGA GGG GAA GAG AGC ATG GCC AAG GCC CTG GTG GCC TGC AAG CTC TAC Arg Gly Glu Glu Ser Met Ala Lys Ala Leu Val Ala Cys Lys Leu Tyr 565 570 575				2082
AAG GCC ATG GCC CAC GAG TCC TCC GAG AGT GAT CTG GTG GAT GAC ATC Lys Ala Met Ala His Glu Ser Ser Glu Ser Asp Leu Val Asp Asp Ile 580 585 590 595				2130
TCC CAG GAC TTG GAT AAC AAT TCC AAA GAC TTC GGC CAG CTT GCT TTG Ser Gln Asp Leu Asp Asn Asn Ser Lys Asp Phe Gly Gln Leu Ala Leu 600 605 610				2178
GAG TTA TTA GAC CAG TCC TAT AAG CAT GAC GAG CAG ATC GCT ATG AAA Glu Leu Leu Asp Gln Ser Tyr Lys His Asp Glu Gln Ile Ala Met Lys 615 620 625				2226
CTC CTG ACC TAC GAG CTG AAA AAC TGG AGC AAC TCG ACC TGC CTC AAA Leu Leu Thr Tyr Glu Leu Lys Asn Trp Ser Asn Ser Thr Cys Leu Lys 630 635 640				2274
CTG GCC GTG GCA GCC AAA CAC CGG GAC TTC ATT GCT CAC ACC TGC AGC Leu Ala Val Ala Ala Lys His Arg Asp Phe Ile Ala His Thr Cys Ser 645 650 655				2322

CAG ATG CTG CTG ACC GAT ATG TGG ATG GGA AGA CTG CGG ATG CGG AAG Gln Met Leu Leu Thr Asp Met Trp Met Gly Arg Leu Arg Met Arg Lys 660 665 670 675	2370
AAC CCC GGC CTG AAG GTT ATC ATG GGG ATT CTT CTA CCC CCC ACC ATC Asn Pro Gly Leu Lys Val Ile Met Gly Ile Leu Leu Pro Pro Thr Ile 680 685 690	2418
TTG TTT TTG GAA TTT CGC ACA TAT GAT GAT TTC TCG TAT CAA ACA TCC Leu Phe Leu Glu Phe Arg Thr Tyr Asp Asp Phe Ser Tyr Gln Thr Ser 695 700 705	2466
AAG GAA AAC GAG GAT GGC AAA GAA AAA GAA GAG GAA AAT ACG GAT GCA Lys Glu Asn Glu Asp Gly Lys Glu Lys Glu Glu Glu Asn Thr Asp Ala 710 715 720	2514
AAT GCA GAT GCT GGC TCA AGA AAG GGG GAT GAG GAG AAC GAG CAT AAA Asn Ala Asp Ala Gly Ser Arg Lys Gly Asp Glu Glu Asn Glu His Lys 725 730 735	2562
AAA CAG AGA ATT ATC CCC ATC GGA ACA AAA ATC TGT AAA TTC TAT AAC Lys Gln Arg Ile Ile Pro Ile Gly Thr Lys Ile Cys Lys Phe Tyr Asn 740 745 750 755	2610
GCG CCC ATT GTC AAG TTC TGG TTT TAC ACA ATA TCA TAC TTG GGC TAC Ala Pro Ile Val Lys Phe Trp Phe Tyr Thr Ile Ser Tyr Leu Gly Tyr 760 765 770	2658
CTG CTG CTG TTT AAC TAC GTC ATC CTG GTG CGG ATG GAT GGC TGG CCG Leu Leu Leu Phe Asn Tyr Val Ile Leu Val Arg Met Asp Gly Trp Pro 775 780 785	2706
TCC CTC CAG GAG TGG ATC GTC ATC TCC TAC ATC GTG AGC CTG GCG TTA Ser Leu Gln Glu Trp Ile Val Ile Ser Tyr Ile Val Ser Leu Ala Leu 790 795 800	2754
GAG AAG ATA CGA GAG ATC CTC ATG TCA GAA CCA GGC AAA CTC AGC CAG Glu Lys Ile Arg Glu Ile Leu Met Ser Glu Pro Gly Lys Leu Ser Gln 805 810 815	2802
AAA ATC AAA GTT TGG CTT CAG GAG TAC TGG AAC ATC ACA GAT CTC GTG Lys Ile Lys Val Trp Leu Gln Glu Tyr Trp Asn Ile Thr Asp Leu Val 820 825 830 835	2850
GCC ATT TCC ACA TTC ATG ATT GGA GCA ATG GCC ACG AGA TCT GTG ATG Ala Ile Ser Thr Phe Met Ile Gly Ala Met Ala Thr Arg Ser Val Met 840 845 850	2898
ATG ATT GGA AAG ATG ATG ATC GAC ATG CTG TAC TTT GTG GTC ATC ATG Met Ile Gly Lys Met Met Ile Asp Met Leu Tyr Phe Val Val Ile Met 855 860 865	2946
CTG GTC GTG CTC ATG AGT TTC GGA GTA GCC CGT CAA GCC ATT CTG CAT Leu Val Val Leu Met Ser Phe Gly Val Ala Arg Gln Ala Ile Leu His 870 875 880	2994
CCA GAG GAG AAG CCC TCT TGG AAA CTG GCC CGA AAC ATC TTC TAC ATG Pro Glu Glu Lys Pro Ser Trp Lys Leu Ala Arg Asn Ile Phe Tyr Met	3042

885	890	895	
CCC TAC TGG ATG ATC TAT GGA GAG GTG TTT GCA GAC CAG ATA GAC CTC Pro Tyr Trp Met Ile Tyr Gly Glu Val Phe Ala Asp Gln Ile Asp Leu 900 905 910 915			3090
TAC GCC ATG GAA ATT AAT CCT CCT TGT GGT GAG AAC CTA TAT GAT GAG Tyr Ala Met Glu Ile Asn Pro Pro Cys Gly Glu Asn Leu Tyr Asp Glu 920 925 930			3138
GAG GGC AAG CGG CTT CCT CCC TGT ATC CCC GGC GCC TGG CTC ACT CCA Glu Gly Lys Arg Leu Pro Pro Cys Ile Pro Gly Ala Trp Leu Thr Pro 935 940 945			3186
GCA CTC ATG GCG TGC TAT CTA CTG GTC GCC AAC ATC CTG CTG GTG AAC Ala Leu Met Ala Cys Tyr Leu Leu Val Ala Asn Ile Leu Leu Val Asn 950 955 960			3234
CTG CTG ATT GCT GTG TTC AAC AAT ACC TTC TTT GAA GTA AAA TCA ATA Leu Leu Ile Ala Val Phe Asn Asn Thr Phe Phe Glu Val Lys Ser Ile 965 970 975			3282
TCC AAC CAG GTG TGG AAG TTC CAG CGA TAT CAG CTG ATT ATG ACA TTT Ser Asn Gln Val Trp Lys Phe Gln Arg Tyr Gln Leu Ile Met Thr Phe 980 985 990 995			3330
CAT GAC AGG CCA GTC CTG CCC CCA CCG ATG ATC ATT TTA AGC CAC ATC His Asp Arg Pro Val Leu Pro Pro Pro Met Ile Ile Leu Ser His Ile 1000 1005 1010			3378
TAC ATC ATC ATT ATG CGT CTC AGC GGC CGC TGC AGG AAA AAG AGA GAA Tyr Ile Ile Ile Met Arg Leu Ser Gly Arg Cys Arg Lys Lys Arg Glu 1015 1020 1025			3426
GGG GAC CAA GAG GAA CGG GAT CGT GGA TTG AAG CTC TTC CTT AGC GAC Gly Asp Gln Glu Glu Arg Asp Arg Gly Leu Lys Leu Phe Leu Ser Asp 1030 1035 1040			3474
GAG GAG CTA AAG AGG CTG CAT GAG TTC GAG GAG CAG TGC GTG CAG GAG Glu Glu Leu Lys Arg Leu His Glu Phe Glu Glu Gln Cys Val Gln Glu 1045 1050 1055			3522
CAC TTC CGG GAG AAG GAG GAT GAG CAG CAG TCG TCC AGC GAC GAG CGC His Phe Arg Glu Lys Glu Asp Glu Gln Gln Ser Ser Ser Asp Glu Arg 1060 1065 1070 1075			3570
ATC CGG GTC ACT TCT GAA AGA GTT GAA AAT ATG TCA ATG AGG TTG GAA Ile Arg Val Thr Ser Glu Arg Val Glu Asn Met Ser Met Arg Leu Glu 1080 1085 1090			3618
GAA ATC AAT GAA AGA GAA ACT TTT ATG AAA ACT TCC CTG CAG ACT GTT Glu Ile Asn Glu Arg Glu Thr Phe Met Lys Thr Ser Leu Gln Thr Val 1095 1100 1105			3666
GAC CTT CGA CTT GCT CAG CTA GAA GAA TTA TCT AAC AGA ATG GTG AAT Asp Leu Arg Leu Ala Gln Leu Glu Glu Leu Ser Asn Arg Met Val Asn 1110 1115 1120			3714

GCT CTT GAA AAT CTT GCG GGA ATC GAC AGG TCT GAC CTG ATC CAG GCA Ala Leu Glu Asn Leu Ala Gly Ile Asp Arg Ser Asp Leu Ile Gln Ala 1125 1130 1135	3762
CGG TCC CGG GCT TCT TCT GAA TGT GAG GCA ACG TAT CTT CTC CGG CAA Arg Ser Arg Ala Ser Ser Glu Cys Glu Ala Thr Tyr Leu Leu Arg Gln 1140 1145 1150 1155	3810
AGC AGC ATC AAT AGC GCT GAT GGC TAC AGC TTG TAT CGA TAT CAT TTT Ser Ser Ile Asn Ser Ala Asp Gly Tyr Ser Leu Tyr Arg Tyr His Phe 1160 1165 1170	3858
AAC GGA GAA GAG TTA TTA TTT GAG GAT ACA TCT CTC TCC ACG TCA CCA Asn Gly Glu Glu Leu Leu Phe Glu Asp Thr Ser Leu Ser Thr Ser Pro 1175 1180 1185	3906
GGG ACA GGA GTC AGG AAA AAA ACC TGT TCC TTC CGT ATA AAG GAA GAG Gly Thr Gly Val Arg Lys Lys Thr Cys Ser Phe Arg Ile Lys Glu Glu 1190 1195 1200	3954
AAG GAC GTG AAA ACG CAC CTA GTC CCA GAA TGT CAG AAC AGT CTT CAC Lys Asp Val Lys Thr His Leu Val Pro Glu Cys Gln Asn Ser Leu His 1205 1210 1215	4002
CTT TCA CTG GGC ACA AGC ACA TCA GCA ACC CCA GAT GGC AGT CAC CTT Leu Ser Leu Gly Thr Ser Thr Ser Ala Thr Pro Asp Gly Ser His Leu 1220 1225 1230 1235	4050
GCA GTA GAT GAC TTA AAG AAC GCT GAA GAG TCA AAA TTA GGT CCA GAT Ala Val Asp Asp Leu Lys Asn Ala Glu Glu Ser Lys Leu Gly Pro Asp 1240 1245 1250	4098
ATT GGG ATT TCA AAG GAA GAT GAT GAA AGA CAG ACA GAC TCT AAA AAA Ile Gly Ile Ser Lys Glu Asp Asp Glu Arg Gln Thr Asp Ser Lys Lys 1255 1260 1265	4146
GAA GAA ACT ATT TCC CCA AGT TTA AAT AAA ACA GAT GTG ATA CAT GGA Glu Glu Thr Ile Ser Pro Ser Leu Asn Lys Thr Asp Val Ile His Gly 1270 1275 1280	4194
CAG GAC AAA TCA GAT GTT CAA AAC ACT CAG CTA ACA GTG GAA ACG ACA Gln Asp Lys Ser Asp Val Gln Asn Thr Gln Leu Thr Val Glu Thr Thr 1285 1290 1295	4242
AAT ATA GAA GGC ACT ATT TCC TAT CCC CTG GAA GAA ACC AAA ATT ACA Asn Ile Glu Gly Thr Ile Ser Tyr Pro Leu Glu Glu Thr Lys Ile Thr 1300 1305 1310 1315	4290
CGC TAT TTC CCC GAT GAA ACG ATC AAT GCT TGT AAA ACA ATG AAG TCC Arg Tyr Phe Pro Asp Glu Thr Ile Asn Ala Cys Lys Thr Met Lys Ser 1320 1325 1330	4338
AGA AGC TTC GTC TAT TCC CGG GGA AGA AAG CTG GTC GGT GGG GTT AAC Arg Ser Phe Val Tyr Ser Arg Gly Arg Lys Leu Val Gly Gly Val Asn 1335 1340 1345	4386
CAG GAT GTA GAG TAC AGT TCA ATC ACG GAC CAG CAA TTG ACG ACG GAA Gln Asp Val Glu Tyr Ser Ser Ile Thr Asp Gln Gln Leu Thr Thr Glu 1350 1355 1360	4434

1350	1355	1360	
TGG CAA TGC CAA GTT CAA AAG ATC ACG CGC TCT CAT AGC ACA GAT ATT Trp Gln Cys Gln Val Gln Lys Ile Thr Arg Ser His Ser Thr Asp Ile 1365 1370 1375			4482
CCT TAC ATT GTG TCG GAA GCT GCA GTG CAA GCT GAG CAA AAA GAG CAG Pro Tyr Ile Val Ser Glu Ala Ala Val Gln Ala Glu Gln Lys Glu Gln 1380 1385 1390 1395			4530
TTT GCA GAT ATG CAA GAT GAA CAC CAT GTC GCT GAA GCA ATT CCT CGA Phe Ala Asp Met Gln Asp Glu His His Val Ala Glu Ala Ile Pro Arg 1400 1405 1410			4578
ATC CCT CGC TTG TCC CTA ACC ATT ACT GAC AGA AAT GGG ATG GAA AAC Ile Pro Arg Leu Ser Leu Thr Ile Thr Asp Arg Asn Gly Met Glu Asn 1415 1420 1425			4626
TTA CTG TCT GTG AAG CCA GAT CAA ACT TTG GGA TTC CCA TCT CTC AGG Leu Leu Ser Val Lys Pro Asp Gln Thr Leu Gly Phe Pro Ser Leu Arg 1430 1435 1440			4674
TCA AAA AGT TTA CAT GGA CAT CCT AGG AAT GTG AAA TCC ATT CAG GGA Ser Lys Ser Leu His Gly His Pro Arg Asn Val Lys Ser Ile Gln Gly 1445 1450 1455			4722
AAG TTA GAC AGA TCT GGA CAT GCC AGT AGT GTA AGC AGC TTA GTA ATT Lys Leu Asp Arg Ser Gly His Ala Ser Ser Val Ser Ser Leu Val Ile 1460 1465 1470 1475			4770
GTG TCT GGA ATG ACA GCA GAA GAA AAA AAG GTT AAG AAA GAG AAA GCT Val Ser Gly Met Thr Ala Glu Glu Lys Lys Val Lys Lys Glu Lys Ala 1480 1485 1490			4818
TCC ACA GAA ACT GAA TGC T AGTCTGTTTT GTTCTTTAA TTTTTTTTTT Ser Thr Glu Thr Glu Cys 1495			4867
TAACAGTCAG AACCACTAAT GGGTGTGATC TTGGCCATCC TAAACATCCA TCCAATTTC			4927
TAAAAACATT TTCCCTT			4944

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1497 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met	Tyr	Ile	Arg	Val	Ser	Tyr	Asp	Thr	Lys	Pro	Asp	Ser	Leu	Leu	His
1					5				10					15	
Leu	Met	Val	Lys	Asp	Trp	Gln	Leu	Glu	Leu	Pro	Lys	Leu	Leu	Ile	Ser
		20						25					30		
Val	His	Gly	Gly	Leu	Gln	Asn	Phe	Glu	Met	Gln	Pro	Lys	Leu	Lys	Gln

35					40					45					
Val	Phe	Gly	Lys	Gly	Leu	Ile	Lys	Ala	Ala	Met	Thr	Thr	Gly	Ala	Trp
50						55					60				
Ile	Phe	Thr	Gly	Gly	Val	Ser	Thr	Gly	Val	Ile	Ser	His	Val	Gly	Asp
65					70					75					80
Ala	Leu	Lys	Asp	His	Ser	Ser	Lys	Ser	Arg	Gly	Arg	Val	Cys	Ala	Ile
				85					90					95	
Gly	Ile	Ala	Pro	Trp	Gly	Ile	Val	Glu	Asn	Lys	Glu	Asp	Leu	Val	Gly
			100					105					110		
Lys	Asp	Val	Thr	Arg	Val	Tyr	Gln	Thr	Met	Ser	Asn	Pro	Leu	Ser	Lys
		115				120						125			
Leu	Ser	Val	Leu	Asn	Asn	Ser	His	Thr	His	Phe	Ile	Leu	Ala	Asp	Asn
	130					135					140				
Gly	Thr	Leu	Gly	Lys	Tyr	Gly	Ala	Glu	Val	Lys	Leu	Arg	Arg	Leu	Leu
145				150						155					160
Glu	Lys	His	Ile	Ser	Leu	Gln	Lys	Ile	Asn	Thr	Arg	Leu	Gly	Gln	Gly
			165						170					175	
Val	Pro	Leu	Val	Gly	Leu	Val	Val	Glu	Gly	Gly	Pro	Asn	Val	Val	Ser
			180					185					190		
Ile	Val	Leu	Glu	Tyr	Leu	Gln	Glu	Glu	Pro	Pro	Ile	Pro	Val	Val	Ile
		195					200					205			
Cys	Asp	Gly	Ser	Gly	Arg	Ala	Ser	Asp	Ile	Leu	Ser	Phe	Ala	His	Lys
	210					215					220				
Tyr	Cys	Glu	Glu	Gly	Gly	Ile	Ile	Asn	Glu	Ser	Leu	Arg	Glu	Gln	Leu
225				230						235					240
Leu	Val	Thr	Ile	Gln	Lys	Thr	Phe	Asn	Tyr	Asn	Lys	Ala	Gln	Ser	His
			245						250					255	
Gln	Leu	Phe	Ala	Ile	Ile	Met	Glu	Cys	Met	Lys	Lys	Lys	Glu	Leu	Val
			260					265					270		
Thr	Val	Phe	Arg	Met	Gly	Ser	Glu	Gly	Gln	Gln	Asp	Ile	Glu	Met	Ala
		275					280					285			
Ile	Leu	Thr	Ala	Leu	Leu	Lys	Gly	Thr	Asn	Val	Ser	Ala	Pro	Asp	Gln
	290					295					300				
Leu	Ser	Leu	Ala	Leu	Ala	Trp	Asn	Arg	Val	Asp	Ile	Ala	Arg	Ser	Gln
305					310					315					320
Ile	Phe	Val	Phe	Gly	Pro	His	Trp	Thr	Pro	Leu	Gly	Ser	Leu	Ala	Pro
			325						330					335	
Pro	Thr	Asp	Ser	Lys	Ala	Thr	Glu	Lys	Glu	Lys	Lys	Pro	Pro	Met	Ala
			340					345					350		

Thr Thr Lys Gly Gly Arg Gly Lys Gly Lys Gly Lys Lys Lys Gly Lys
 355 360 365
 Val Lys Glu Glu Val Glu Glu Glu Thr Asp Pro Arg Lys Ile Glu Leu
 370 375 380
 Leu Asn Trp Val Asn Ala Leu Glu Gln Ala Met Leu Asp Ala Leu Val
 385 390 395 400
 Leu Asp Arg Val Asp Phe Val Lys Leu Leu Ile Glu Asn Gly Val Asn
 405 410 415
 Met Gln His Phe Leu Thr Ile Pro Arg Leu Glu Glu Leu Tyr Asn Thr
 420 425 430
 Arg Leu Gly Pro Pro Asn Thr Leu His Leu Leu Val Arg Asp Val Lys
 435 440 445
 Lys Ser Asn Leu Pro Pro Asp Tyr His Ile Ser Leu Ile Asp Ile Gly
 450 455 460
 Leu Val Leu Glu Tyr Leu Met Gly Gly Ala Tyr Arg Cys Asn Tyr Thr
 465 470 475 480
 Arg Lys Asn Phe Arg Thr Leu Tyr Asn Asn Leu Phe Gly Pro Lys Arg
 485 490 495
 Pro Lys Ala Leu Lys Leu Leu Gly Met Glu Asp Asp Glu Pro Pro Ala
 500 505 510
 Lys Gly Lys Lys Lys Lys Lys Lys Lys Lys Glu Glu Glu Ile Asp Ile
 515 520 525
 Asp Val Asp Asp Pro Ala Val Ser Arg Phe Gln Tyr Pro Phe His Glu
 530 535 540
 Leu Met Val Trp Ala Val Leu Met Lys Arg Gln Lys Met Ala Val Phe
 545 550 555 560
 Leu Trp Gln Arg Gly Glu Glu Ser Met Ala Lys Ala Leu Val Ala Cys
 565 570 575
 Lys Leu Tyr Lys Ala Met Ala His Glu Ser Ser Glu Ser Asp Leu Val
 580 585 590
 Asp Asp Ile Ser Gln Asp Leu Asp Asn Asn Ser Lys Asp Phe Gly Gln
 595 600 605
 Leu Ala Leu Glu Leu Leu Asp Gln Ser Tyr Lys His Asp Glu Gln Ile
 610 615 620
 Ala Met Lys Leu Leu Thr Tyr Glu Leu Lys Asn Trp Ser Asn Ser Thr
 625 630 635 640
 Cys Leu Lys Leu Ala Val Ala Ala Lys His Arg Asp Phe Ile Ala His
 645 650 655

Thr Cys Ser Gln Met Leu Leu Thr Asp Met Trp Met Gly Arg Leu Arg
 660 665 670
 Met Arg Lys Asn Pro Gly Leu Lys Val Ile Met Gly Ile Leu Leu Pro
 675 680 685
 Pro Thr Ile Leu Phe Leu Glu Phe Arg Thr Tyr Asp Asp Phe Ser Tyr
 690 695 700
 Gln Thr Ser Lys Glu Asn Glu Asp Gly Lys Glu Lys Glu Glu Glu Asn
 705 710 715 720
 Thr Asp Ala Asn Ala Asp Ala Gly Ser Arg Lys Gly Asp Glu Glu Asn
 725 730 735
 Glu His Lys Lys Gln Arg Ile Ile Pro Ile Gly Thr Lys Ile Cys Lys
 740 745 750
 Phe Tyr Asn Ala Pro Ile Val Lys Phe Trp Phe Tyr Thr Ile Ser Tyr
 755 760 765
 Leu Gly Tyr Leu Leu Leu Phe Asn Tyr Val Ile Leu Val Arg Met Asp
 770 775 780
 Gly Trp Pro Ser Leu Gln Glu Trp Ile Val Ile Ser Tyr Ile Val Ser
 785 790 795 800
 Leu Ala Leu Glu Lys Ile Arg Glu Ile Leu Met Ser Glu Pro Gly Lys
 805 810 815
 Leu Ser Gln Lys Ile Lys Val Trp Leu Gln Glu Tyr Trp Asn Ile Thr
 820 825 830
 Asp Leu Val Ala Ile Ser Thr Phe Met Ile Gly Ala Met Ala Thr Arg
 835 840 845
 Ser Val Met Met Ile Gly Lys Met Met Ile Asp Met Leu Tyr Phe Val
 850 855 860
 Val Ile Met Leu Val Val Leu Met Ser Phe Gly Val Ala Arg Gln Ala
 865 870 875 880
 Ile Leu His Pro Glu Glu Lys Pro Ser Trp Lys Leu Ala Arg Asn Ile
 885 890 895
 Phe Tyr Met Pro Tyr Trp Met Ile Tyr Gly Glu Val Phe Ala Asp Gln
 900 905 910
 Ile Asp Leu Tyr Ala Met Glu Ile Asn Pro Pro Cys Gly Glu Asn Leu
 915 920 925
 Tyr Asp Glu Glu Gly Lys Arg Leu Pro Pro Cys Ile Pro Gly Ala Trp
 930 935 940
 Leu Thr Pro Ala Leu Met Ala Cys Tyr Leu Leu Val Ala Asn Ile Leu
 945 950 955 960

Leu Val Asn Leu Leu Ile Ala Val Phe Asn Asn Thr Phe Phe Glu Val
 965 970 975
 Lys Ser Ile Ser Asn Gln Val Trp Lys Phe Gln Arg Tyr Gln Leu Ile
 980 985 990
 Met Thr Phe His Asp Arg Pro Val Leu Pro Pro Pro Met Ile Ile Leu
 995 1000 1005
 Ser His Ile Tyr Ile Ile Ile Met Arg Leu Ser Gly Arg Cys Arg Lys
 1010 1015 1020
 Lys Arg Glu Gly Asp Gln Glu Glu Arg Asp Arg Gly Leu Lys Leu Phe
 1025 1030 1035 1040
 Leu Ser Asp Glu Glu Leu Lys Arg Leu His Glu Phe Glu Glu Gln Cys
 1045 1050 1055
 Val Gln Glu His Phe Arg Glu Lys Glu Asp Glu Gln Gln Ser Ser Ser
 1060 1065 1070
 Asp Glu Arg Ile Arg Val Thr Ser Glu Arg Val Glu Asn Met Ser Met
 1075 1080 1085
 Arg Leu Glu Glu Ile Asn Glu Arg Glu Thr Phe Met Lys Thr Ser Leu
 1090 1095 1100
 Gln Thr Val Asp Leu Arg Leu Ala Gln Leu Glu Glu Leu Ser Asn Arg
 1105 1110 1115 1120
 Met Val Asn Ala Leu Glu Asn Leu Ala Gly Ile Asp Arg Ser Asp Leu
 1125 1130 1135
 Ile Gln Ala Arg Ser Arg Ala Ser Ser Glu Cys Glu Ala Thr Tyr Leu
 1140 1145 1150
 Leu Arg Gln Ser Ser Ile Asn Ser Ala Asp Gly Tyr Ser Leu Tyr Arg
 1155 1160 1165
 Tyr His Phe Asn Gly Glu Glu Leu Leu Phe Glu Asp Thr Ser Leu Ser
 1170 1175 1180
 Thr Ser Pro Gly Thr Gly Val Arg Lys Lys Thr Cys Ser Phe Arg Ile
 1185 1190 1195 1200
 Lys Glu Glu Lys Asp Val Lys Thr His Leu Val Pro Glu Cys Gln Asn
 1205 1210 1215
 Ser Leu His Leu Ser Leu Gly Thr Ser Thr Ser Ala Thr Pro Asp Gly
 1220 1225 1230
 Ser His Leu Ala Val Asp Asp Leu Lys Asn Ala Glu Glu Ser Lys Leu
 1235 1240 1245
 Gly Pro Asp Ile Gly Ile Ser Lys Glu Asp Asp Glu Arg Gln Thr Asp
 1250 1255 1260

Ser Lys Lys Glu Glu Thr Ile Ser Pro Ser Leu Asn Lys Thr Asp Val
 1265 1270 1275 1280
 Ile His Gly Gln Asp Lys Ser Asp Val Gln Asn Thr Gln Leu Thr Val
 1285 1290 1295
 Glu Thr Thr Asn Ile Glu Gly Thr Ile Ser Tyr Pro Leu Glu Glu Thr
 1300 1305 1310
 Lys Ile Thr Arg Tyr Phe Pro Asp Glu Thr Ile Asn Ala Cys Lys Thr
 1315 1320 1325
 Met Lys Ser Arg Ser Phe Val Tyr Ser Arg Gly Arg Lys Leu Val Gly
 1330 1335 1340
 Gly Val Asn Gln Asp Val Glu Tyr Ser Ser Ile Thr Asp Gln Gln Leu
 1345 1350 1355 1360
 Thr Thr Glu Trp Gln Cys Gln Val Gln Lys Ile Thr Arg Ser His Ser
 1365 1370 1375
 Thr Asp Ile Pro Tyr Ile Val Ser Glu Ala Ala Val Gln Ala Glu Gln
 1380 1385 1390
 Lys Glu Gln Phe Ala Asp Met Gln Asp Glu His His Val Ala Glu Ala
 1395 1400 1405
 Ile Pro Arg Ile Pro Arg Leu Ser Leu Thr Ile Thr Asp Arg Asn Gly
 1410 1415 1420
 Met Glu Asn Leu Leu Ser Val Lys Pro Asp Gln Thr Leu Gly Phe Pro
 1425 1430 1435 1440
 Ser Leu Arg Ser Lys Ser Leu His Gly His Pro Arg Asn Val Lys Ser
 1445 1450 1455
 Ile Gln Gly Lys Leu Asp Arg Ser Gly His Ala Ser Ser Val Ser Ser
 1460 1465 1470
 Leu Val Ile Val Ser Gly Met Thr Ala Glu Glu Lys Lys Val Lys Lys
 1475 1480 1485
 Glu Lys Ala Ser Thr Glu Thr Glu Cys
 1490 1495

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5055 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 346..4945

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

ACTCATTATA GGGNTCGAGC GGCCGCCCGG GCAGGTTTGA GCTGTGCCCT CTCCATTCCA	60
CTGCTGTGGC AGGGTCAGAA ATCTTGATA GAGAAAACCT TTTGCAAACG GGAATGTATC	120
TTTGTAATTC CTAGCACGAA AGACTCTAAC AGGTGTTGCT GTGGCCAGTT CACCAACCAG	180
CATATCCCCC CTCTGCCAAG TGCAACACCC AGCAAAAATG AAGAGGAAAG CAAACAGGTG	240
GAGACTCAGC CTGAGAAATG GTCTGTTGCC AAGCACACCC AGAGCTACCC AACAGATTCC	300
TATGGAGTTC TTGAATTCCA GGGTGGCGGA TATTCCAATA AAGCC ATG TAT ATC	354
Met Tyr Ile	
1	
CGT GTA TCC TAT GAC ACC AAG CCA GAC TCA CTG CTC CAT CTC ATG GTG	402
Arg Val Ser Tyr Asp Thr Lys Pro Asp Ser Leu Leu His Leu Met Val	
5 10 15	
AAA GAT TGG CAG CTG GAA CTC CCC AAG CTC TTA ATA TCT GTG CAT GGA	450
Lys Asp Trp Gln Leu Glu Leu Pro Lys Leu Leu Ile Ser Val His Gly	
20 25 30 35	
GGC CTC CAG AAC TTT GAG ATG CAG CCC AAG CTG AAA CAA GTC TTT GGG	498
Gly Leu Gln Asn Phe Glu Met Gln Pro Lys Leu Lys Gln Val Phe Gly	
40 45 50	
AAA GGC CTG ATC AAG GCT GCT ATG ACC ACC GGG GCC TGG ATC TTC ACC	546
Lys Gly Leu Ile Lys Ala Ala Met Thr Thr Gly Ala Trp Ile Phe Thr	
55 60 65	
GGG GGT GTC AGC ACA GGT GTT ATC AGC CAC GTA GGG GAT GCC TTG AAA	594
Gly Gly Val Ser Thr Gly Val Ile Ser His Val Gly Asp Ala Leu Lys	
70 75 80	
GAC CAC TCC TCC AAG TCC AGA GGC CGG GTT TGT GCT ATA GGA ATT GCT	642
Asp His Ser Ser Lys Ser Arg Gly Arg Val Cys Ala Ile Gly Ile Ala	
85 90 95	
CCA TGG GGC ATC GTG GAG AAT AAG GAA GAC CTG GTT GGA AAG GAT GTA	690
Pro Trp Gly Ile Val Glu Asn Lys Glu Asp Leu Val Gly Lys Asp Val	
100 105 110 115	
ACA AGA GTG TAC CAG ACC ATG TCC AAC CCT CTA AGT AAG CTC TCT GTG	738
Thr Arg Val Tyr Gln Thr Met Ser Asn Pro Leu Ser Lys Leu Ser Val	
120 125 130	
CTC AAC AAC TCC CAC ACC CAC TTC ATC CTG GCT GAC AAT GGC ACC CTG	786
Leu Asn Asn Ser His Thr His Phe Ile Leu Ala Asp Asn Gly Thr Leu	
135 140 145	
GGC AAG TAT GGC GCC GAG GTG AAG CTG CGA AGG CTG CTG GAA AAG CAC	834

Gly	Lys	Tyr	Gly	Ala	Glu	Val	Lys	Leu	Arg	Arg	Leu	Leu	Glu	Lys	His	
	150						155				160					
ATC	TCC	CTC	CAG	AAG	ATC	AAC	ACA	AGA	CTG	GGG	CAG	GGC	GTG	CCC	CTC	882
Ile	Ser	Leu	Gln	Lys	Ile	Asn	Thr	Arg	Leu	Gly	Gln	Gly	Val	Pro	Leu	
	165					170				175						
GTG	GGT	CTC	GTG	GTG	GAG	GGG	GGC	CCT	AAC	GTG	GTG	TCC	ATC	GTC	TTG	930
Val	Gly	Leu	Val	Val	Glu	Gly	Gly	Pro	Asn	Val	Val	Ser	Ile	Val	Leu	
180					185					190					195	
GAA	TAC	CTG	CAA	GAA	GAG	CCT	CCC	ATC	CCT	GTG	GTG	ATT	TGT	GAT	GGC	978
Glu	Tyr	Leu	Gln	Glu	Glu	Pro	Pro	Ile	Pro	Val	Val	Ile	Cys	Asp	Gly	
			200					205						210		
AGC	GGA	CGT	GCC	TCG	GAC	ATC	CTG	TCC	TTT	GCG	CAC	AAG	TAC	TGT	GAA	1026
Ser	Gly	Arg	Ala	Ser	Asp	Ile	Leu	Ser	Phe	Ala	His	Lys	Tyr	Cys	Glu	
			215					220					225			
GAA	GGC	GGA	ATA	ATA	AAT	GAG	TCC	CTC	AGG	GAG	CAG	CTT	CTA	GTT	ACC	1074
Glu	Gly	Gly	Ile	Ile	Asn	Glu	Ser	Leu	Arg	Glu	Gln	Leu	Leu	Val	Thr	
		230					235					240				
ATT	CAG	AAA	ACA	TTT	AAT	TAT	AAT	AAG	GCA	CAA	TCA	CAT	CAG	CTG	TTT	1122
Ile	Gln	Lys	Thr	Phe	Asn	Tyr	Asn	Lys	Ala	Gln	Ser	His	Gln	Leu	Phe	
	245					250					255					
GCA	ATT	ATA	ATG	GAG	TGC	ATG	AAG	AAG	AAA	GAA	CTC	GTC	ACT	GTG	TTC	1170
Ala	Ile	Ile	Met	Glu	Cys	Met	Lys	Lys	Lys	Glu	Leu	Val	Thr	Val	Phe	
260					265					270					275	
AGA	ATG	GGT	TCT	GAG	GGC	CAG	CAG	GAC	ATC	GAG	ATG	GCA	ATT	TTA	ACT	1218
Arg	Met	Gly	Ser	Glu	Gly	Gln	Gln	Asp	Ile	Glu	Met	Ala	Ile	Leu	Thr	
				280					285					290		
GCC	CTG	CTG	AAA	GGA	ACA	AAC	GTA	TCT	GCT	CCA	GAT	CAG	CTG	AGC	TTG	1266
Ala	Leu	Leu	Lys	Gly	Thr	Asn	Val	Ser	Ala	Pro	Asp	Gln	Leu	Ser	Leu	
			295					300					305			
GCA	CTG	GCT	TGG	AAC	CGC	GTG	GAC	ATA	GCA	CGA	AGC	CAG	ATC	TTT	GTC	1314
Ala	Leu	Ala	Trp	Asn	Arg	Val	Asp	Ile	Ala	Arg	Ser	Gln	Ile	Phe	Val	
		310					315					320				
TTT	GGG	CCC	CAC	TGG	ACG	CCC	CTG	GGA	AGC	CTG	GCA	CCC	CCG	ACG	GAC	1362
Phe	Gly	Pro	His	Trp	Thr	Pro	Leu	Gly	Ser	Leu	Ala	Pro	Pro	Thr	Asp	
	325					330					335					
AGC	AAA	GCC	ACG	GAG	AAG	GAG	AAG	AAG	CCA	CCC	ATG	GCC	ACC	ACC	AAG	1410
Ser	Lys	Ala	Thr	Glu	Lys	Glu	Lys	Lys	Pro	Pro	Met	Ala	Thr	Thr	Lys	
					345					350					355	
GGA	GGA	AGA	GGA	AAA	GGG	AAA	GGC	AAG	AAG	AAA	GGG	AAA	GTG	AAA	GAG	1458
Gly	Gly	Arg	Gly	Lys	Gly	Lys	Gly	Lys	Lys	Lys	Gly	Lys	Val	Lys	Glu	
				360				365					370			
GAA	GTG	GAG	GAA	GAA	ACT	GAC	CCC	CGG	AAG	ATA	GAG	CTG	CTG	AAC	TGG	1506
Glu	Val	Glu	Glu	Glu	Thr	Asp	Pro	Arg	Lys	Ile	Glu	Leu	Leu	Asn	Trp	
				375				380					385			

GTG AAT GCT TTG GAG CAA GCG ATG CTA GAT GCT TTA GTC TTA GAT CGT	1554
Val Asn Ala Leu Glu Gln Ala Met Leu Asp Ala Leu Val Leu Asp Arg	
390 395 400	
GTC GAC TTT GTG AAG CTC CTG ATT GAA AAC GGA GTG AAC ATG CAA CAC	1602
Val Asp Phe Val Lys Leu Leu Ile Glu Asn Gly Val Asn Met Gln His	
405 410 415	
TTT CTG ACC ATT CCG AGG CTG GAG GAG CTT TAT AAC ACA AGA CTG GGT	1650
Phe Leu Thr Ile Pro Arg Leu Glu Glu Leu Tyr Asn Thr Arg Leu Gly	
420 425 430 435	
CCA CCA AAC ACA CTT CAT CTG CTG GTG AGG GAT GTG AAA AAG AGC AAC	1698
Pro Pro Asn Thr Leu His Leu Leu Val Arg Asp Val Lys Lys Ser Asn	
440 445 450	
CTT CCG CCT GAT TAC CAC ATC AGC CTC ATA GAC ATC GGG CTC GTG CTG	1746
Leu Pro Pro Asp Tyr His Ile Ser Leu Ile Asp Ile Gly Leu Val Leu	
455 460 465	
GAG TAC CTC ATG GGA GGA GCC TAC CGC TGC AAC TAC ACT CGG AAA AAC	1794
Glu Tyr Leu Met Gly Gly Ala Tyr Arg Cys Asn Tyr Thr Arg Lys Asn	
470 475 480	
TTT CGG ACC CTT TAC AAC AAC TTG TTT GGA CCA AAG AGG CCT AAA GCT	1842
Phe Arg Thr Leu Tyr Asn Asn Leu Phe Gly Pro Lys Arg Pro Lys Ala	
485 490 495	
CTT AAA CTT CTG GGA ATG GAA GAT GAT GAG CCT CCA GCT AAA GGG AAG	1890
Leu Lys Leu Leu Gly Met Glu Asp Asp Glu Pro Pro Ala Lys Gly Lys	
500 505 510 515	
AAA AAA AAA AAA AAG AAA AAG GAG GAA GAG ATC GAC ATT GAT GTG GAC	1938
Lys Lys Lys Lys Lys Lys Lys Glu Glu Glu Ile Asp Ile Asp Val Asp	
520 525 530	
GAC CCT GCC GTG AGT CGG TTC CAG TAT CCC TTC CAC GAG CTG ATG GTG	1986
Asp Pro Ala Val Ser Arg Phe Gln Tyr Pro Phe His Glu Leu Met Val	
535 540 545	
TGG GCA GTG CTG ATG AAA CGC CAG AAA ATG GCA GTG TTC CTC TGG CAG	2034
Trp Ala Val Leu Met Lys Arg Gln Lys Met Ala Val Phe Leu Trp Gln	
550 555 560	
CGA GGG GAA GAG AGC ATG GCC AAG GCC CTG GTG GCC TGC AAG CTC TAC	2082
Arg Gly Glu Glu Ser Met Ala Lys Ala Leu Val Ala Cys Lys Leu Tyr	
565 570 575	
AAG GCC ATG GCC CAC GAG TCC TCC GAG AGT GAT CTG GTG GAT GAC ATC	2130
Lys Ala Met Ala His Glu Ser Ser Glu Ser Asp Leu Val Asp Asp Ile	
580 585 590 595	
TCC CAG GAC TTG GAT AAC AAT TCC AAA GAC TTC GGC CAG CTT GCT TTG	2178
Ser Gln Asp Leu Asp Asn Asn Ser Lys Asp Phe Gly Gln Leu Ala Leu	
600 605 610	
GAG TTA TTA GAC CAG TCC TAT AAG CAT GAC GAG CAG ATC GCT ATG AAA	2226
Glu Leu Leu Asp Gln Ser Tyr Lys His Asp Glu Gln Ile Ala Met Lys	

615					620					625						
CTC	CTG	ACC	TAC	GAG	CTG	AAA	AAC	TGG	AGC	AAC	TCG	ACC	TGC	CTC	AAA	2274
Leu	Leu	Thr	Tyr	Glu	Leu	Lys	Asn	Trp	Ser	Asn	Ser	Thr	Cys	Leu	Lys	
		630					635					640				
CTG	GCC	GTG	GCA	GCC	AAA	CAC	CGG	GAC	TTC	ATT	GCT	CAC	ACC	TGC	AGC	2322
Leu	Ala	Val	Ala	Ala	Lys	His	Arg	Asp	Phe	Ile	Ala	His	Thr	Cys	Ser	
	645					650					655					
CAG	ATG	CTG	CTG	ACC	GAT	ATG	TGG	ATG	GGA	AGA	CTG	CGG	ATG	CGG	AAG	2370
Gln	Met	Leu	Leu	Thr	Asp	Met	Trp	Met	Gly	Arg	Leu	Arg	Met	Arg	Lys	
660					665					670					675	
AAC	CCC	GGC	CTG	AAG	GTT	ATC	ATG	GGG	ATT	CTT	CTA	CCC	CCC	ACC	ATC	2418
Asn	Pro	Gly	Leu	Lys	Val	Ile	Met	Gly	Ile	Leu	Leu	Pro	Pro	Thr	Ile	
				680					685					690		
TTG	TTT	TTG	GAA	TTT	CGC	ACA	TAT	GAT	GAT	TTC	TCG	TAT	CAA	ACA	TCC	2466
Leu	Phe	Leu	Glu	Phe	Arg	Thr	Tyr	Asp	Asp	Phe	Ser	Tyr	Gln	Thr	Ser	
			695					700					705			
AAG	GAA	AAC	GAG	GAT	GGC	AAA	GAA	AAA	GAA	GAG	GAA	AAT	ACG	GAT	GCA	2514
Lys	Glu	Asn	Glu	Asp	Gly	Lys	Glu	Lys	Glu	Glu	Glu	Asn	Thr	Asp	Ala	
		710					715					720				
AAT	GCA	GAT	GCT	GGC	TCA	AGA	AAG	GGG	GAT	GAG	GAG	AAC	GAG	CAT	AAA	2562
Asn	Ala	Asp	Ala	Gly	Ser	Arg	Lys	Gly	Asp	Glu	Glu	Asn	Glu	His	Lys	
	725					730						735				
AAA	CAG	AGA	AGT	ATT	CCC	ATC	GGA	ACA	AAG	ATC	TGT	GAA	TTC	TAT	AAC	2610
Lys	Gln	Arg	Ser	Ile	Pro	Ile	Gly	Thr	Lys	Ile	Cys	Glu	Phe	Tyr	Asn	
740					745					750					755	
GCG	CCC	ATT	GTC	AAG	TTC	TGG	TTT	TAC	ACA	ATA	TCA	TAC	TTG	GGC	TAC	2658
Ala	Pro	Ile	Val	Lys	Phe	Trp	Phe	Tyr	Thr	Ile	Ser	Tyr	Leu	Gly	Tyr	
				760					765					770		
CTG	CTG	CTG	TTT	AAC	TAC	GTC	ATC	CTG	GTG	CGG	ATG	GAT	GGC	TGG	CCG	2706
Leu	Leu	Leu	Phe	Asn	Tyr	Val	Ile	Leu	Val	Arg	Met	Asp	Gly	Trp	Pro	
			775					780					785			
TCC	CTC	CAG	GAG	TGG	ATC	GTC	ATC	TCC	TAC	ATC	GTG	AGC	CTG	GCG	TTA	2754
Ser	Leu	Gln	Glu	Trp	Ile	Val	Ile	Ser	Tyr	Ile	Val	Ser	Leu	Ala	Leu	
		790					795					800				
GAG	AAG	ATA	CGA	GAG	ATC	CTC	ATG	TCA	GAA	CCA	GGC	AAA	CTC	AGC	CAG	2802
Glu	Lys	Ile	Arg	Glu	Ile	Leu	Met	Ser	Glu	Pro	Gly	Lys	Leu	Ser	Gln	
	805					810					815					
AAA	ATC	AAA	GTT	TGG	CTT	CAG	GAG	TAC	TGG	AAC	ATC	ACA	GAT	CTC	GTG	2850
Lys	Ile	Lys	Val	Trp	Leu	Gln	Glu	Tyr	Trp	Asn	Ile	Thr	Asp	Leu	Val	
820					825					830					835	
GCC	ATT	TCC	ACA	TTC	ATG	ATT	GGA	GCA	ATT	CTT	CGC	CTA	CAG	AAC	CAG	2898
Ala	Ile	Ser	Thr	Phe	Met	Ile	Gly	Ala	Ile	Leu	Arg	Leu	Gln	Asn	Gln	
				840					845					850		

CCC	TAC	ATG	GGC	TAT	GGC	CGG	GTG	ATC	TAC	TGT	GTG	GAT	ATC	ATC	TTC	2946
Pro	Tyr	Met	Gly	Tyr	Gly	Arg	Val	Ile	Tyr	Cys	Val	Asp	Ile	Ile	Phe	
			855					860					865			
TGG	TAC	ATC	CGT	GTC	CTG	GAC	ATC	TTT	GGT	GTC	AAC	AAG	TAT	CTG	GGG	2994
Trp	Tyr	Ile	Arg	Val	Leu	Asp	Ile	Phe	Gly	Val	Asn	Lys	Tyr	Leu	Gly	
		870					875					880				
CCA	TAC	GTG	ATG	ATG	ATT	GGA	AAG	ATG	ATG	ATC	GAC	ATG	CTG	TAC	TTT	3042
Pro	Tyr	Val	Met	Met	Ile	Gly	Lys	Met	Met	Ile	Asp	Met	Leu	Tyr	Phe	
		885				890					895					
GTG	GTC	ATC	ATG	CTG	GTC	GTG	CTC	ATG	AGT	TTC	GGA	GTA	GCC	CGT	CAA	3090
Val	Val	Ile	Met	Leu	Val	Val	Leu	Met	Ser	Phe	Gly	Val	Ala	Arg	Gln	
900					905					910					915	
GCC	ATT	CTG	CAT	CCA	GAG	GAG	AAG	CCC	TCT	TGG	AAA	CTG	GCC	CGA	AAC	3138
Ala	Ile	Leu	His	Pro	Glu	Glu	Lys	Pro	Ser	Trp	Lys	Leu	Ala	Arg	Asn	
				920					925					930		
ATC	TTC	TAC	ATG	CCC	TAC	TGG	ATG	ATC	TAT	GGA	GAG	GTG	TTT	GCA	GAC	3186
Ile	Phe	Tyr	Met	Pro	Tyr	Trp	Met	Ile	Tyr	Gly	Glu	Val	Phe	Ala	Asp	
			935					940					945			
CAG	ATA	GAC	CTC	TAC	GCC	ATG	GAA	ATT	AAT	CCT	CCT	TGT	GGT	GAG	AAC	3234
Gln	Ile	Asp	Leu	Tyr	Ala	Met	Glu	Ile	Asn	Pro	Pro	Cys	Gly	Glu	Asn	
		950				955						960				
CTA	TAT	GAT	GAG	GAG	GGC	AAG	CGG	CTT	CCT	CCC	TGT	ATC	CCC	GGC	GCC	3282
Leu	Tyr	Asp	Glu	Glu	Gly	Lys	Arg	Leu	Pro	Pro	Cys	Ile	Pro	Gly	Ala	
		965				970					975					
TGG	CTC	ACT	CCA	GCA	CTC	ATG	GCG	TGC	TAT	CTA	CTG	GTC	GCC	AAC	ATC	3330
Trp	Leu	Thr	Pro	Ala	Leu	Met	Ala	Cys	Tyr	Leu	Leu	Val	Ala	Asn	Ile	
980					985					990					995	
CTG	CTG	GTG	AAC	CTG	CTG	ATT	GCT	GTG	TTC	AAC	AAT	ACT	TTC	TTT	GAA	3378
Leu	Leu	Val	Asn	Leu	Leu	Ile	Ala	Val	Phe	Asn	Asn	Thr	Phe	Phe	Glu	
				1000				1005					1010			
GTA	AAA	TCA	ATA	TCC	AAC	CAG	GTG	TGG	AAG	TTC	CAG	CGA	TAT	CAG	CTG	3426
Val	Lys	Ser	Ile	Ser	Asn	Gln	Val	Trp	Lys	Phe	Gln	Arg	Tyr	Gln	Leu	
			1015				1020					1025				
ATT	ATG	ACA	TTT	CAT	GAC	AGG	CCA	GTC	CTG	CCC	CCA	CCG	ATG	ATC	ATT	3474
Ile	Met	Thr	Phe	His	Asp	Arg	Pro	Val	Leu	Pro	Pro	Pro	Met	Ile	Ile	
			1030				1035					1040				
TTA	AGC	CAC	ATC	TAC	ATC	ATC	ATT	ATG	CGT	CTC	AGC	GGC	CGC	TGC	AGG	3522
Leu	Ser	His	Ile	Tyr	Ile	Ile	Ile	Met	Arg	Leu	Ser	Gly	Arg	Cys	Arg	
			1045			1050					1055					
AAA	AAG	AGA	GAA	GGG	GAC	CAA	GAG	GAA	CGG	GAT	CGT	GGA	TTG	AAG	CTC	3570
Lys	Lys	Arg	Glu	Gly	Asp	Gln	Glu	Glu	Arg	Asp	Arg	Gly	Leu	Lys	Leu	
1060				1065					1070						1075	
TTC	CTT	AGC	GAC	GAG	GAG	CTA	AAG	AGG	CTG	CAT	GAG	TTC	GAG	GAG	CAG	3618
Phe	Leu	Ser	Asp	Glu	Glu	Leu	Lys	Arg	Leu	His	Glu	Phe	Glu	Glu	Gln	

1080										1085					1090					
TGC	GTG	CAG	GAG	CAC	TTC	CGG	GAG	AAG	GAG	GAT	GAG	CAG	CAG	TCG	TCC					3666
Cys	Val	Gln	Glu	His	Phe	Arg	Glu	Lys	Glu	Asp	Glu	Gln	Gln	Ser	Ser					
			1095					1100					1105							
AGC	GAC	GAG	CGC	ATC	CGG	GTC	ACT	TCT	GAA	AGA	GTT	GAA	AAT	ATG	TCA					3714
Ser	Asp	Glu	Arg	Ile	Arg	Val	Thr	Ser	Glu	Arg	Val	Glu	Asn	Met	Ser					
		1110					1115					1120								
ATG	AGG	TTG	GAA	GAA	ATC	AAT	GAA	AGA	GAA	ACT	TTT	ATG	AAA	ACT	TCC					3762
Met	Arg	Leu	Glu	Glu	Ile	Asn	Glu	Arg	Glu	Thr	Phe	Met	Lys	Thr	Ser					
		1125				1130					1135									
CTG	CAG	ACT	GTT	GAC	CTT	CGA	CTT	GCT	CAG	CTA	GAA	GAA	TTA	TCT	AAC					3810
Leu	Gln	Thr	Val	Asp	Leu	Arg	Leu	Ala	Gln	Leu	Glu	Glu	Leu	Ser	Asn					
		1140			1145				1150						1155					
AGA	ATG	GTG	AAT	GCT	CTT	GAA	AAT	CTT	GCG	GGA	ATC	GAC	AGG	TCT	GAC					3858
Arg	Met	Val	Asn	Ala	Leu	Glu	Asn	Leu	Ala	Gly	Ile	Asp	Arg	Ser	Asp					
				1160					1165						1170					
CTG	ATC	CAG	GCA	CGG	TCC	CGG	GCT	TCT	TCT	GAA	TGT	GAG	GCA	ACG	TAT					3906
Leu	Ile	Gln	Ala	Arg	Ser	Arg	Ala	Ser	Ser	Glu	Cys	Glu	Ala	Thr	Tyr					
			1175					1180						1185						
CTT	CTC	CGG	CAA	AGC	AGC	ATC	AAT	AGC	GCT	GAT	GGC	TAC	AGC	TTG	TAT					3954
Leu	Leu	Arg	Gln	Ser	Ser	Ile	Asn	Ser	Ala	Asp	Gly	Tyr	Ser	Leu	Tyr					
		1190					1195					1200								
CGA	TAT	CAT	TTT	AAC	GGA	GAA	GAG	TTA	TTA	TTT	GAG	GAT	ACA	TCT	CTC					4002
Arg	Tyr	His	Phe	Asn	Gly	Glu	Glu	Leu	Leu	Phe	Glu	Asp	Thr	Ser	Leu					
		1205				1210					1215									
TCC	ACG	TCA	CCA	GGG	ACA	GGA	GTC	AGG	AAA	AAA	ACC	TGT	TCC	TTC	CGT					4050
Ser	Thr	Ser	Pro	Gly	Thr	Gly	Val	Arg	Lys	Lys	Thr	Cys	Ser	Phe	Arg					
				1225					1230						1235					
ATA	AAG	GAA	GAG	AAG	GAC	GTG	AAA	ACG	CAC	CTA	GTC	CCA	GAA	TGT	CAG					4098
Ile	Lys	Glu	Glu	Lys	Asp	Val	Lys	Thr	His	Leu	Val	Pro	Glu	Cys	Gln					
				1240					1245					1250						
AAC	AGT	CTT	CAC	CTT	TCA	CTG	GGC	ACA	AGC	ACA	TCA	GCA	ACC	CCA	GAT					4146
Asn	Ser	Leu	His	Leu	Ser	Leu	Gly	Thr	Ser	Thr	Ser	Ala	Thr	Pro	Asp					
				1255				1260					1265							
GGC	AGT	CAC	CTT	GCA	GTA	GAT	GAC	TTA	AAG	AAC	GCT	GAA	GAG	TCA	AAA					4194
Gly	Ser	His	Leu	Ala	Val	Asp	Asp	Leu	Lys	Asn	Ala	Glu	Glu	Ser	Lys					
			1270				1275					1280								
TTA	GGT	CCA	GAT	ATT	GGG	ATT	TCA	AAG	GAA	GAT	GAT	GAA	AGA	CAG	ACA					4242
Leu	Gly	Pro	Asp	Ile	Gly	Ile	Ser	Lys	Glu	Asp	Asp	Glu	Arg	Gln	Thr					
		1285				1290					1295									
GAC	TCT	AAA	AAA	GAA	GAA	ACT	ATT	TCC	CCA	AGT	TTA	AAT	AAA	ACA	GAT					4290
Asp	Ser	Lys	Lys	Glu	Glu	Thr	Ile	Ser	Pro	Ser	Leu	Asn	Lys	Thr	Asp					
				1300		1305				1310					1315					

GTG ATA CAT GGA CAG GAC AAA TCA GAT GTT CAA AAC ACT CAG CTA ACA Val Ile His Gly Gln Asp Lys Ser Asp Val Gln Asn Thr Gln Leu Thr 1320 1325 1330	4338
GTG GAA ACG ACA AAT ATA GAA GGC ACT ATT TCC TAT CCC CTG GAA GAA Val Glu Thr Thr Asn Ile Glu Gly Thr Ile Ser Tyr Pro Leu Glu Glu 1335 1340 1345	4386
ACC AAA ATT ACA CGC TAT TTC CCC GAT GAA ACG ATC AAT GCT TGT AAA Thr Lys Ile Thr Arg Tyr Phe Pro Asp Glu Thr Ile Asn Ala Cys Lys 1350 1355 1360	4434
ACA ATG AAG TCC AGA AGC TTC GTC TAT TCC CGG GGA AGA AAG CTG GTC Thr Met Lys Ser Arg Ser Phe Val Tyr Ser Arg Gly Arg Lys Leu Val 1365 1370 1375	4482
GGT GGG GTT AAC CAG GAT GTA GAG TAC AGT TCA ATC ACG GAC CAG CAA Gly Gly Val Asn Gln Asp Val Glu Tyr Ser Ser Ile Thr Asp Gln Gln 1380 1385 1390 1395	4530
TTG ACG ACG GAA TGG CAA TGC CAA GTT CAA AAG ATC ACG CGC TCT CAT Leu Thr Thr Glu Trp Gln Cys Gln Val Gln Lys Ile Thr Arg Ser His 1400 1405 1410	4578
AGC ACA GAT ATT CCT TAC ATT GTG TCG GAA GCT GCA GTG CAA GCT GAG Ser Thr Asp Ile Pro Tyr Ile Val Ser Glu Ala Ala Val Gln Ala Glu 1415 1420 1425	4626
CAA AAA GAG CAG TTT GCA GAT ATG CAA GAT GAA CAC CAT GTC GCT GAA Gln Lys Glu Gln Phe Ala Asp Met Gln Asp Glu His His Val Ala Glu 1430 1435 1440	4674
GCA ATT CCT CGA ATC CCT CGC TTG TCC CTA ACC ATT ACT GAC AGA AAT Ala Ile Pro Arg Ile Pro Arg Leu Ser Leu Thr Ile Thr Asp Arg Asn 1445 1450 1455	4722
GGG ATG GAA AAC TTA CTG TCT GTG AAG CCA GAT CAA ACT TTG GGA TTC Gly Met Glu Asn Leu Ser Val Lys Pro Asp Gln Thr Leu Gly Phe 1460 1465 1470 1475	4770
CCA TCT CTC AGG TCA AAA AGT TTA CAT GGA CAT CCT AGG AAT GTG AAA Pro Ser Leu Arg Ser Lys Ser Leu His Gly His Pro Arg Asn Val Lys 1480 1485 1490	4818
TCC ATT CAG GGA AAG TTA GAC AGA TCT GGA CAT GCC AGT AGT GTA AGC Ser Ile Gln Gly Lys Leu Asp Arg Ser Gly His Ala Ser Ser Val Ser 1495 1500 1505	4866
AGC TTA GTA ATT GTG TCT GGA ATG ACA GCA GAA GAA AAA AAG GTT AAG Ser Leu Val Ile Val Ser Gly Met Thr Ala Glu Glu Lys Lys Val Lys 1510 1515 1520	4914
AAA GAG AAA GCT TCC ACA GAA ACT GAA TGC T AGTCTGTTTT GTTCTTTAA Lys Glu Lys Ala Ser Thr Glu Thr Glu Cys 1525 1530	4965
TTTTTTTTTT TAACAGTCAG AAACCCACTA ATGGGTGTCA TCTTGCCCA TCCTAAACAC	5025

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1533 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

```

Met Tyr Ile Arg Val Ser Tyr Asp Thr Lys Pro Asp Ser Leu Leu His
 1           5           10           15
Leu Met Val Lys Asp Trp Gln Leu Glu Leu Pro Lys Leu Leu Ile Ser
          20           25           30
Val His Gly Gly Leu Gln Asn Phe Glu Met Gln Pro Lys Leu Lys Gln
          35           40           45
Val Phe Gly Lys Gly Leu Ile Lys Ala Ala Met Thr Thr Gly Ala Trp
          50           55           60
Ile Phe Thr Gly Gly Val Ser Thr Gly Val Ile Ser His Val Gly Asp
          65           70           75           80
Ala Leu Lys Asp His Ser Ser Lys Ser Arg Gly Arg Val Cys Ala Ile
          85           90           95
Gly Ile Ala Pro Trp Gly Ile Val Glu Asn Lys Glu Asp Leu Val Gly
          100          105          110
Lys Asp Val Thr Arg Val Tyr Gln Thr Met Ser Asn Pro Leu Ser Lys
          115          120          125
Leu Ser Val Leu Asn Asn Ser His Thr His Phe Ile Leu Ala Asp Asn
          130          135          140
Gly Thr Leu Gly Lys Tyr Gly Ala Glu Val Lys Leu Arg Arg Leu Leu
          145          150          155          160
Glu Lys His Ile Ser Leu Gln Lys Ile Asn Thr Arg Leu Gly Gln Gly
          165          170          175
Val Pro Leu Val Gly Leu Val Val Glu Gly Gly Pro Asn Val Val Ser
          180          185          190
Ile Val Leu Glu Tyr Leu Gln Glu Glu Pro Pro Ile Pro Val Val Ile
          195          200          205
Cys Asp Gly Ser Gly Arg Ala Ser Asp Ile Leu Ser Phe Ala His Lys
          210          215          220
Tyr Cys Glu Glu Gly Gly Ile Ile Asn Glu Ser Leu Arg Glu Gln Leu
          225          230          235          240

```

Leu Val Thr Ile Gln Lys Thr Phe Asn Tyr Asn Lys Ala Gln Ser His
 245 250 255
 Gln Leu Phe Ala Ile Ile Met Glu Cys Met Lys Lys Lys Glu Leu Val
 260 265 270
 Thr Val Phe Arg Met Gly Ser Glu Gly Gln Gln Asp Ile Glu Met Ala
 275 280 285
 Ile Leu Thr Ala Leu Leu Lys Gly Thr Asn Val Ser Ala Pro Asp Gln
 290 295 300
 Leu Ser Leu Ala Leu Ala Trp Asn Arg Val Asp Ile Ala Arg Ser Gln
 305 310 315 320
 Ile Phe Val Phe Gly Pro His Trp Thr Pro Leu Gly Ser Leu Ala Pro
 325 330 335
 Pro Thr Asp Ser Lys Ala Thr Glu Lys Glu Lys Lys Pro Pro Met Ala
 340 345 350
 Thr Thr Lys Gly Gly Arg Gly Lys Gly Lys Gly Lys Lys Lys Gly Lys
 355 360 365
 Val Lys Glu Glu Val Glu Glu Glu Thr Asp Pro Arg Lys Ile Glu Leu
 370 375 380
 Leu Asn Trp Val Asn Ala Leu Glu Gln Ala Met Leu Asp Ala Leu Val
 385 390 395 400
 Leu Asp Arg Val Asp Phe Val Lys Leu Leu Ile Glu Asn Gly Val Asn
 405 410 415
 Met Gln His Phe Leu Thr Ile Pro Arg Leu Glu Glu Leu Tyr Asn Thr
 420 425 430
 Arg Leu Gly Pro Pro Asn Thr Leu His Leu Leu Val Arg Asp Val Lys
 435 440 445
 Lys Ser Asn Leu Pro Pro Asp Tyr His Ile Ser Leu Ile Asp Ile Gly
 450 455 460
 Leu Val Leu Glu Tyr Leu Met Gly Gly Ala Tyr Arg Cys Asn Tyr Thr
 465 470 475 480
 Arg Lys Asn Phe Arg Thr Leu Tyr Asn Asn Leu Phe Gly Pro Lys Arg
 485 490 495
 Pro Lys Ala Leu Lys Leu Leu Gly Met Glu Asp Asp Glu Pro Pro Ala
 500 505 510
 Lys Gly Lys Lys Lys Lys Lys Lys Lys Lys Glu Glu Glu Ile Asp Ile
 515 520 525
 Asp Val Asp Asp Pro Ala Val Ser Arg Phe Gln Tyr Pro Phe His Glu
 530 535 540

Leu Met Val Trp Ala Val Leu Met Lys Arg Gln Lys Met Ala Val Phe
 545 550 555 560
 Leu Trp Gln Arg Gly Glu Glu Ser Met Ala Lys Ala Leu Val Ala Cys
 565 570 575
 Lys Leu Tyr Lys Ala Met Ala His Glu Ser Ser Glu Ser Asp Leu Val
 580 585 590
 Asp Asp Ile Ser Gln Asp Leu Asp Asn Asn Ser Lys Asp Phe Gly Gln
 595 600 605
 Leu Ala Leu Glu Leu Leu Asp Gln Ser Tyr Lys His Asp Glu Gln Ile
 610 615 620
 Ala Met Lys Leu Leu Thr Tyr Glu Leu Lys Asn Trp Ser Asn Ser Thr
 625 630 635 640
 Cys Leu Lys Leu Ala Val Ala Ala Lys His Arg Asp Phe Ile Ala His
 645 650 655
 Thr Cys Ser Gln Met Leu Leu Thr Asp Met Trp Met Gly Arg Leu Arg
 660 665 670
 Met Arg Lys Asn Pro Gly Leu Lys Val Ile Met Gly Ile Leu Leu Pro
 675 680 685
 Pro Thr Ile Leu Phe Leu Glu Phe Arg Thr Tyr Asp Asp Phe Ser Tyr
 690 695 700
 Gln Thr Ser Lys Glu Asn Glu Asp Gly Lys Glu Lys Glu Glu Glu Asn
 705 710 715 720
 Thr Asp Ala Asn Ala Asp Ala Gly Ser Arg Lys Gly Asp Glu Glu Asn
 725 730 735
 Glu His Lys Lys Gln Arg Ser Ile Pro Ile Gly Thr Lys Ile Cys Glu
 740 745 750
 Phe Tyr Asn Ala Pro Ile Val Lys Phe Trp Phe Tyr Thr Ile Ser Tyr
 755 760 765
 Leu Gly Tyr Leu Leu Leu Phe Asn Tyr Val Ile Leu Val Arg Met Asp
 770 775 780
 Gly Trp Pro Ser Leu Gln Glu Trp Ile Val Ile Ser Tyr Ile Val Ser
 785 790 795 800
 Leu Ala Leu Glu Lys Ile Arg Glu Ile Leu Met Ser Glu Pro Gly Lys
 805 810 815
 Leu Ser Gln Lys Ile Lys Val Trp Leu Gln Glu Tyr Trp Asn Ile Thr
 820 825 830
 Asp Leu Val Ala Ile Ser Thr Phe Met Ile Gly Ala Ile Leu Arg Leu
 835 840 845

Gln Asn Gln Pro Tyr Met Gly Tyr Gly Arg Val Ile Tyr Cys Val Asp
 850 855 860
 Ile Ile Phe Trp Tyr Ile Arg Val Leu Asp Ile Phe Gly Val Asn Lys
 865 870 875 880
 Tyr Leu Gly Pro Tyr Val Met Met Ile Gly Lys Met Met Ile Asp Met
 885 890 895
 Leu Tyr Phe Val Val Ile Met Leu Val Val Leu Met Ser Phe Gly Val
 900 905 910
 Ala Arg Gln Ala Ile Leu His Pro Glu Glu Lys Pro Ser Trp Lys Leu
 915 920 925
 Ala Arg Asn Ile Phe Tyr Met Pro Tyr Trp Met Ile Tyr Gly Glu Val
 930 935 940
 Phe Ala Asp Gln Ile Asp Leu Tyr Ala Met Glu Ile Asn Pro Pro Cys
 945 950 955 960
 Gly Glu Asn Leu Tyr Asp Glu Glu Gly Lys Arg Leu Pro Pro Cys Ile
 965 970 975
 Pro Gly Ala Trp Leu Thr Pro Ala Leu Met Ala Cys Tyr Leu Leu Val
 980 985 990
 Ala Asn Ile Leu Leu Val Asn Leu Leu Ile Ala Val Phe Asn Asn Thr
 995 1000 1005
 Phe Phe Glu Val Lys Ser Ile Ser Asn Gln Val Trp Lys Phe Gln Arg
 1010 1015 1020
 Tyr Gln Leu Ile Met Thr Phe His Asp Arg Pro Val Leu Pro Pro Pro
 1025 1030 1035 1040
 Met Ile Ile Leu Ser His Ile Tyr Ile Ile Ile Met Arg Leu Ser Gly
 1045 1050 1055
 Arg Cys Arg Lys Lys Arg Glu Gly Asp Gln Glu Glu Arg Asp Arg Gly
 1060 1065 1070
 Leu Lys Leu Phe Leu Ser Asp Glu Glu Leu Lys Arg Leu His Glu Phe
 1075 1080 1085
 Glu Glu Gln Cys Val Gln Glu His Phe Arg Glu Lys Glu Asp Glu Gln
 1090 1095 1100
 Gln Ser Ser Ser Asp Glu Arg Ile Arg Val Thr Ser Glu Arg Val Glu
 1105 1110 1115 1120
 Asn Met Ser Met Arg Leu Glu Glu Ile Asn Glu Arg Glu Thr Phe Met
 1125 1130 1135
 Lys Thr Ser Leu Gln Thr Val Asp Leu Arg Leu Ala Gln Leu Glu Glu
 1140 1145 1150

Leu Ser Asn Arg Met Val Asn Ala Leu Glu Asn Leu Ala Gly Ile Asp
 1155 1160 1165
 Arg Ser Asp Leu Ile Gln Ala Arg Ser Arg Ala Ser Ser Glu Cys Glu
 1170 1175 1180
 Ala Thr Tyr Leu Leu Arg Gln Ser Ser Ile Asn Ser Ala Asp Gly Tyr
 1185 1190 1195 1200
 Ser Leu Tyr Arg Tyr His Phe Asn Gly Glu Glu Leu Leu Phe Glu Asp
 1205 1210 1215
 Thr Ser Leu Ser Thr Ser Pro Gly Thr Gly Val Arg Lys Lys Thr Cys
 1220 1225 1230
 Ser Phe Arg Ile Lys Glu Glu Lys Asp Val Lys Thr His Leu Val Pro
 1235 1240 1245
 Glu Cys Gln Asn Ser Leu His Leu Ser Leu Gly Thr Ser Thr Ser Ala
 1250 1255 1260
 Thr Pro Asp Gly Ser His Leu Ala Val Asp Asp Leu Lys Asn Ala Glu
 1265 1270 1275 1280
 Glu Ser Lys Leu Gly Pro Asp Ile Gly Ile Ser Lys Glu Asp Asp Glu
 1285 1290 1295
 Arg Gln Thr Asp Ser Lys Lys Glu Glu Thr Ile Ser Pro Ser Leu Asn
 1300 1305 1310
 Lys Thr Asp Val Ile His Gly Gln Asp Lys Ser Asp Val Gln Asn Thr
 1315 1320 1325
 Gln Leu Thr Val Glu Thr Thr Asn Ile Glu Gly Thr Ile Ser Tyr Pro
 1330 1335 1340
 Leu Glu Glu Thr Lys Ile Thr Arg Tyr Phe Pro Asp Glu Thr Ile Asn
 1345 1350 1355 1360
 Ala Cys Lys Thr Met Lys Ser Arg Ser Phe Val Tyr Ser Arg Gly Arg
 1365 1370 1375
 Lys Leu Val Gly Gly Val Asn Gln Asp Val Glu Tyr Ser Ser Ile Thr
 1380 1385 1390
 Asp Gln Gln Leu Thr Thr Glu Trp Gln Cys Gln Val Gln Lys Ile Thr
 1395 1400 1405
 Arg Ser His Ser Thr Asp Ile Pro Tyr Ile Val Ser Glu Ala Ala Val
 1410 1415 1420
 Gln Ala Glu Gln Lys Glu Gln Phe Ala Asp Met Gln Asp Glu His His
 1425 1430 1435 1440
 Val Ala Glu Ala Ile Pro Arg Ile Pro Arg Leu Ser Leu Thr Ile Thr
 1445 1450 1455

Asp Arg Asn Gly Met Glu Asn Leu Leu Ser Val Lys Pro Asp Gln Thr
1460 1465 1470

Leu Gly Phe Pro Ser Leu Arg Ser Lys Ser Leu His Gly His Pro Arg
1475 1480 1485

Asn Val Lys Ser Ile Gln Gly Lys Leu Asp Arg Ser Gly His Ala Ser
1490 1495 1500

Ser Val Ser Ser Leu Val Ile Val Ser Gly Met Thr Ala Glu Glu Lys
1505 1510 1515 1520

Lys Val Lys Lys Glu Lys Ala Ser Thr Glu Thr Glu Cys
1525 1530

174591.B11

WHAT IS CLAIMED IS:

1 1. An isolated nucleic acid comprising the
2 nucleotide sequence SEQ ID NO:2 as depicted in FIG. 3, or
3 the nucleotide sequence of a gene or gene fragment contained
4 in the following clone as deposited with the NRRL: pFOMY030
5 (NRRL accession No. B-21416).

1 2. An isolated nucleic acid which encodes the
2 amino acid sequence depicted in Fig. 3 from amino acid
3 residue number 1 to 542, or its complement.

1 3. An isolated nucleic acid which hybridizes to an
2 isolated nucleic acid which encodes the amino acid sequence
3 depicted in Fig. 3 from amino acid residue number 1 to 542,
4 or its complement.

1 4. A genetically engineered host cell comprising
2 the nucleotide sequence of Claim 2.

1 5. An expression vector comprising the nucleotide
2 sequence of Claim 2 in operative association with a
3 nucleotide sequence regulatory element that controls
4 expression of the nucleotide sequence in a host cell.

1 6. A substantially pure gene product encoded by the
2 nucleic acid of Claim 3.

1 7. An isolated nucleic acid comprising the
2 nucleotide sequence SEQ ID NO:6 as depicted in FIG. 5, or
3 SEQ ID NO:8 as depicted in FIG. 6.

1 8. An isolated nucleic acid which encodes the amino
2 acid sequence depicted in Fig. 5 from amino acid residue
3 number 1 to 1497, or its complement. ✓

1 9. An isolated nucleic acid which encodes the amino ✓
2 acid sequence depicted in Fig. 6 from amino acid residue
3 number 1 to 1533, or its complement.

1 10. An isolated nucleic acid which hybridizes to a ✓
2 nucleic acid that encodes the amino acid sequence depicted
3 in Fig. 5 from amino acid residue number 1 to 1497 (SEQ ID
4 NO:7), or its complement, or the amino acid sequence
5 depicted in Fig. 6 from amino acid residue number 1 to 1533
6 (SEQ ID NO:9), or its complement.

1 11. A nucleotide vector containing the nucleotide
2 sequence of Claim 10.

1 12. A genetically engineered host cell containing
2 the nucleotide sequence of Claim 10.

1 13. An expression vector containing the nucleotide
2 sequence of Claim 10 in operative association with a
3 nucleotide sequence regulatory element that controls
4 expression of the nucleotide sequence in a host cell.

1 14. A substantially pure gene product encoded by
2 the nucleic acid of Claim 10.

1 15. An antibody that immunospecifically binds the
2 gene product of Claim 14.

1 16. A method of diagnosing tumor progression in a
2 mammal, said method comprising:
3 obtaining a test sample of tissue cells from the
4 mammal;
5 obtaining a control sample of known normal cells
6 from the same type of tissue; and
7 detecting in both the test sample and the control
8 sample the level of expression of a gene transcript or gene
9 product of gene 030, wherein a level of expression lower in
10 the test sample than in the control sample indicates a tumor
11 progression state in the test sample.

1 17. A method for treating tumor progression in a
2 mammal, said method comprising administering to the mammal a
3 compound in an amount effective to increase the level of
4 expression or activity of a gene transcript or gene product
5 of gene 030 in cells exhibiting a tumor progression state,
6 to a level effective to ameliorate symptoms of the tumor
7 progression state.

1 18. A method of Claim 17, wherein the compound
2 comprises a nucleic acid whose administration results in an
3 increase in the level of the differentially expressed gene
4 transcript and gene product in the cells in the tumor
5 progression state, thereby ameliorating symptoms of the
6 tumor progression state.

1 19. A method of Claim 18, wherein the nucleic acid
2 comprises a nucleic acid of gene 030.

1 20. A method of Claim 17, wherein the compound is a
2 030 gene product.

1 21. A method of Claim 17, wherein the tumor
2 progression state is neoplasia.

1 22. A method of Claim 17, wherein the tumor
2 progression state is metastasis.

1 23. A method for inhibiting tumor progression in a
2 mammal, said method comprising administering to the mammal a
3 normal allele of a 030 gene so that the normal gene product
4 is expressed, thereby inhibiting tumor progression.

1 24. A method of Claim 23, wherein the tumor
2 progression is metastasis.

1 25. A method for treating tumor progression in a
2 mammal, said method comprising administering to the mammal
3 an effective amount of a 030 gene product.

1 26. A method of Claim 25, wherein the tumor
2 progression is metastasis.

1 27. A method of monitoring the efficacy of a
2 compound in clinical trials for inhibition of tumor
3 progression in a patient, said method comprising
4 obtaining a first sample of tumor tissue cells from
5 the patient;
6 administering the compound to the patient;
7 after a time sufficient for the compound to inhibit
8 tumor progression, obtaining a second sample of tumor tissue
9 cells from the patient; and
10 detecting in the first and second samples the level
11 of expression of a 030 gene transcript or product, wherein a
12 level of expression higher in the second sample than in the
13 first sample indicates that the compound is effective to
14 inhibit tumor progression in the patient.

1 28. A method of Claim 27, wherein the 030 gene
2 transcript or gene product is differentially expressed in
3 individuals predisposed to a metastatic neoplastic disease.

174591.B11

**COMPOSITIONS AND METHODS FOR THE DIAGNOSIS,
PREVENTION AND TREATMENT OF TUMOR PROGRESSION**

ABSTRACT

The present invention relates to methods and compositions for the diagnosis, prevention, and treatment of tumor progression in cells involved in human tumors such as melanomas, breast, gastrointestinal, lung, and bone tumors, various types of skin cancers, and other neoplastic conditions such as leukemias and lymphomas. Genes are identified that are differentially expressed in benign (e.g., non-malignant) tumor cells relative to malignant tumor cells exhibiting a high metastatic potential. Genes are also identified via the ability of their gene products to interact with gene products involved in the progression to, and/or aggressiveness of, neoplastic tumor disease states. The genes and gene products identified can be used diagnostically or for therapeutic intervention.

174591.B11

FIGURE 1




FIGURE 2

romy030

GGTGCTGGAGTACCTCATGGGCGGTGCCTACCGCTGCAACTACACTCGGAAAAGCTTCCGGACTCT
CTACAACAACCTTGTTTGGCCCTAAGAGGGTAGAGCTCAGCAGACACACAGTGTCTGTGCCTCCC
AGAGTAACATGTGGTTCCTTGATGTGCTTCCCCAAAAGCCACCTGTGCAGAATG
(SEQ ID NO:1)

08644-0537
468250-2442580

FIGURE 3A

AAGGAGGCTAGGCTGCACCCCTTCCCGCTTGCTCAGCAGCTGAGGCAGGGTCAGAAAGCATGGATAGAGAAGACATTTTGC	30
AAAAGGGAATGCATCTTTGTAAATCCAGTACAAAAGACCTAACAGATGTTGCTGTGTCAGCTCACTAACCAGCAGAT	160
CCCCCTTTGCGAGTGGGGCTCCCGCACAACAGGAGAGGACACCAAGCAGGCAGACACGCAGTCCCGGAATGGTCTG	240
TCAGCAAACACACCCAGAGCTACCCAACAGACTCCTATGGGATTCTTGAATTCAGGGTGGGGTTACTCCAATAAAGCC	320
ATG TAC ATC CGA GTC TCC TAC GAC ACC AAG CCA GAT TCC CTG CTC CAC CTC ATG GTG AAG	380
M Y I R V S Y D T K P D S L L H L M V K	20
GAC TGG CAG CTG GAG CTC CCG AAG CTC TTG ATA TCT GTG CAC GGA GGC CTC CAA AGC TTC	440
D W Q L E L P K L L I S V H G G L Q S F	40
GAG ATG CAG TCC AAA CTG AAG CAG GTG TTT GGG AAA GGT CTG ATC AAG GCT GCC ATG ACC	500
E M Q S K L K Q V F G K G L I K A A M T	60
ACG GCG GCG TGG ATC TTC ACC GGG GGT GTG AGC ACT GGT GTC GTC ACC GAT CTG CAG GAT	560
T G A W I F T G G V S T G V V S H V G D	80
CCC TTG AAA GAC CAC TCC TCC AAG TCC AGA GGC CGG CTC TGT GCT ATA GGA ATT GCT CCC	620
A L K D H S S K S R G R L C A I G I A P	100
TGG GGC ATG GTG GAG AAC AAG GAA GAC CTG ATT GGA AAA GAT GTA ACA AGA GTC TAT CAG	680
W G M V E N K E D L I G K D V T R V Y Q	120
ACC ATG TCC AAC CCT CTG AGC AAG CTC TCT GTG CTC AAC AAT TCC CAC ACT CAC TTC ATC	740
T M S N P L S K L S V L N N S H T H F I	140
TTG GCT GAC AAC GGC ACC CTG GGC AAG TAT GGT GCT GAG GTG AAG CTT CGA AGA CAG CTG	800
L A D N G T L G K Y G A E V K L R R Q L	160
GAA AAA CAC ATC TCC CTG CAG AAG ATC AAC ACA AGG CTG GGC CAG GGT GTA CCT GTC GTG	860
E K H I S L Q K I N T R L G Q G V P V V	180
GGC CTA GTG GTA GAA GGT GGT CCT AAC GTG GTT TCT ATC GTC CTG GAG TAT CTC AAA GAA	920
G L V V E G G P N V V S I V L E Y L K E	200
GAC CCT CCT GTC CCT GTG GTG GTT TGC GAT GGC AGT GGA CGT GCC TCT GAC ATT TTG TCC	980
D P P V P V V V C D G S G R A S D I L S	220
TTC GCA CAC AAA TAC TGC GAC GAA GGA GGA GTC ATA AAC GAG TCC CTG CGG GAC CAG CTT	1040
F A H K Y C D E G G V I N E S L P D Q L	240
CTA GTT ACC ATT CAG AAA ACA TTT AAT TAC AGC AAG TCC CAG TCG TAT CAG CTG TTT GCA	1100
L V T I Q K T F N Y S K S Q S Y Q L F A	260
ATT ATC ATG GAG TGC ATG AAG AAG AAA GAA CTC GTC ACT GTG TTT CGG ATG GGT TCC GAG	1160
I I M E C M K K K E L V T V F R M G S E	280
GGT CAG CAA GAT GTC GAG ATG GCA ATT TTA ACT GCC TTG CTC AAA GGA ACC AAC GCA TCA	1220
G Q Q D V E M A I L T A L L K G T N A S	300
GCT CCA GAT CAG CTG AGC TTG GCC CTG GCT TGG AAC CGG GTC GAC ATA GCG CGA AGC CAG	1280
A P D Q L S L A L A W N R V D I A R S Q	320
ATC TTC GTC TTT GGC CCA CAC TGG CCG CCA CTG GGA AGC CTG GCC CCT CCT GTG GAC ACC	1340
I F V F G P H W P P L G S L A P P V D T	340
AAA GCC GCA GAG AAG GAA AAG AAG CCA CCC ACA GCC ACC ACC AAG GGG AGA GGA AAA GGA	1400
K A A E K E K K P P T A T T K G R G K G	360
AAA GGC AAG AAG AAA GGC AAA GTG AAA GAG GAA GTG GAG GAA GAG ACG GAC CCC CGG AAG	1460
K G K K K G K V K E E V E E E T D P R K	380

FIGURE 3B

CTT GAG CTG CTC AAC TGG GTG AAT GCC CTG GAG CAA GCC ATG CTG GAT GCT CTT GTC CTA 1520
L E L L N W V N A L E Q A M L D A L V L 400

GAT CGG GTG GAC TTT GTA AAG CTC CTG ATT GAA AAC GGA GTG AAC ATG CAG CAT TTC CTC 1580
D R V D F V K L L I E N G V M M Q H F L 420

ACC ATC CCG AGG CTG GAG GAG CTA TAC AAC ACC AGA CTG GGC CCA CCA AAC ACC CTT CAT 1640
T I P R L E E L Y N T R L G P P M T L H 440

CTG CTG GTG CGG GAT GTA AAG AAG AGC AAC CTT CCA CCT GAT TAC CAC ATC AGC CTC ATT 1700
L L V R D V K K S N L P P D Y H I S L I 460

GAT ATA GGA CTG GTG CTG GAG TAC CTC ATG GGC GGT GCC TAC CGC TGC AAC TAC ACT CGG 1760
D I G L V L E Y L M G G A Y R C N Y T R 480

AAA AGC TTC CGG ACT CTC TAC AAC AAC TTG TTT GGC CCT AAG AAG GTA GAG CTC ACC AGA 1820
K S F R T L Y N N L F G P K R V E L S R 500

CAC ACA GTG TCC TGT GCC TCC CAG AGT AAC ATG TGG TTC CTT GAT GTG CTT CCC CAA AAG 1880
H T V S C A S Q S N M W F L D V L P Q K 520


CCC ACC TGT GCA GAA TGC AAC TCT TCA CCT CAC CTG TCC CAA ACT GAC ATC ACC CCA CCT 1940
P T C A E C N S S P H L S Q T D I T P P 540

CTG CCC TGA CACCCAGTGCAGGGCCTCCTAGCTTTCACATGCAGCCATTCACATCGCCTCTCAGACTGGGCCAGGC 2018
L P * (SER 18 NO:3)

AGTGCAACCTGTCAAGCATGTCTGTCTCCCTCCCTCCTTCTACAAATAGCCCCCCTCTGGGCCCCATGCCCTCTGCTCTCTC 2098
AGCCCGTTCTCTCCCACTGATCACTGGCGCTCCTGTTGTCTTCCAAGGCAAGGAACAAGGAAAAGCATCTTTTGGCCC 2178
ACAAAAGTTTAGGGCTCCCCGCTGTTCACCATAGCCAACCTCACTGTACATCGGAGTCATCCAGGCCAGCTGCCACACA 2258
CAAGCCTTCCCCACCCTATCCCAATAGACCCTATTCTCCATCAAAATCAAAGCTAACTCCTGGCCTGCCACATTGCTTC 2338
TTCTTGCTCCAGCCTGTTAAACCTCCAATAAATGTCAGATCTGTGGGAAGCCTTCTCACTCTCACTCCACAGTTTGTAC 2418
AGAGAGCGAGAGCCTCGTTTGGTTCTACTTACAAGGAAGGCTTTGTGTCTGTCTGTCTTCCCACTGACTTCTGTGAC 2498
AGAAGCAGTTTCCACATGAAAGCGTTGACTCACCTGGATGTTGTCTTAATTAATAGTGATACAAATATTGACACTTCT 2578
TTTCTGCTTCTTTGTTATGCAGCCGAAAGCACTTAAGCTTCTGGGAATGGAAGTAGTAGGACATGTTTGTGGCAGTTT 2658
ATTTACTATATACCTTTGTCTATTCTGTGGAAGCAAAAATTGCAATGTTTCCATGAATAAGCTCTGTGCC 2730
(SER 18 NO:2)

466350 "2442980"

FIGURE 4



0882449 0882449 0882449

FIGURE 5

ACTCATTATAGGGNTCGAGCGGCCCGCCCGGGCAGGTTTGAGCTGTGCCCTCTCCATTCCACTGCTGTGGCAGGGTCAGA
 AATCTTGGATAGAGAAAACCTTTTGCAAACGGGAATGTATCTTTGTAATTCCTAGCACGAAAGACTCTAACAGGTGTTG
 CTGTGGCCAGTTACCAACCAGCATATCCCCCTCTGCCAAGTGCAACACCCAGCAAAAATGAAGAGGAAAGCAAACAG
 GTGGAGACTCAGCCTGAGAAATGGTCTGTTGCAAGCACACCCAGAGCTACCCAACAGATTCTTATGGAGTTCTTGAATT
 CCAGGGTGGCGATATTCCAATAAAGCC

M	Y	I	R	V	S	Y	D	T	K	P	D	S	L	L	H	L	M	V	K		20
ATG	TAT	ATC	CGT	GTA	TCC	TAT	GAC	ACC	AAG	CCA	GAC	TCA	CTG	CTC	CAT	CTC	ATG	GTG	AAA		60
D	W	Q	L	E	L	P	K	L	L	I	S	V	H	G	G	L	Q	N	F		40
GAT	TGG	CAG	CTG	GAA	CTC	CCC	AAG	CTC	TTA	ATA	TCT	GTG	CAT	GGA	GGC	CTC	CAG	AAC	TTT		120
E	M	Q	P	K	L	K	Q	V	F	G	K	G	L	I	K	A	A	M	T		60
GAG	ATG	CAG	CCC	AAG	CTG	AAA	CAA	GTC	TTT	GGG	AAA	GGC	CTG	ATC	AAG	GCT	GCT	ATG	ACC		180
T	G	A	W	I	F	T	G	G	V	S	T	G	V	I	S	H	V	G	D		80
ACC	GGG	GCC	TGG	ATC	TTC	ACC	GGG	GGT	GTC	AGC	ACA	GGT	GTT	ATC	AGC	CAC	GTA	GGG	GAT		240
A	L	K	D	H	S	S	K	S	R	G	R	V	C	A	I	G	I	A	P		100
GCC	TTG	AAA	GAC	CAC	TCC	TCC	AAG	TCC	AGA	GGC	CGG	GTT	TGT	GCT	ATA	GGA	ATT	GCT	CCA		300
W	G	I	V	E	N	K	E	D	L	V	G	K	D	V	T	R	V	Y	Q		120
TGG	GGC	ATC	GTG	GAG	AAT	AAG	GAA	GAC	CTG	GTT	GGA	AAG	GAT	GTA	ACA	AGA	GTG	TAC	CAG		360
T	M	S	N	P	L	S	K	L	S	V	L	N	N	S	H	T	H	F	I		140
ACC	ATG	TCC	AAC	CCT	CTA	AGT	AAG	CTC	TCT	GTG	CTC	AAC	AAC	TCC	CAC	ACC	CAC	TTC	ATC		420
L	A	D	N	G	T	L	G	K	Y	G	A	E	V	K	L	R	R	L	L		160
CTG	GCT	GAC	AAT	GGC	ACC	CTG	GGC	AAG	TAT	GGC	GCC	GAG	GTG	AAG	CTG	CGA	AGG	CTG	CTG		480
E	K	H	I	S	L	Q	K	I	N	T	R	L	G	Q	G	V	P	L	V		180
GAA	AAG	CAC	ATC	TCC	CTC	CAG	AAG	ATC	AAC	ACA	AGA	CTG	GGG	CAG	GGC	GTG	CCC	CTC	GTG		540
G	L	V	V	E	G	G	P	N	V	V	S	I	V	L	E	Y	L	Q	E		200
GGT	CTC	GTG	GTG	GAG	GGG	GGC	CCT	AAC	GTG	GTG	TCC	ATC	GTC	TTG	GAA	TAC	CTG	CAA	GAA		600
E	P	P	I	P	V	V	I	C	D	G	S	G	R	A	S	D	I	L	S		220
GAG	CCT	CCC	ATC	CCT	GTG	GTG	ATT	TGT	GAT	GGC	AGC	GGA	CGT	GCC	TCG	GAC	ATC	CTG	TCC		660
F	A	H	K	Y	C	E	E	G	G	I	I	N	E	S	L	R	E	Q	L		240
TTT	GCG	CAC	AAG	TAC	TGT	GAA	GAA	GGC	GGA	ATA	ATA	AAT	GAG	TCC	CTC	AGG	GAG	CAG	CTT		720
L	V	T	I	Q	K	T	F	N	Y	N	K	A	Q	S	H	Q	L	F	A		260
CTA	GTT	ACC	ATT	CAG	AAA	ACA	TTT	AAT	TAT	AAT	AAG	GCA	CAA	TCA	CAT	CAG	CTG	TTT	GCA		780
I	I	M	E	C	M	K	K	K	E	L	V	T	V	F	R	M	G	S	E		280
ATT	ATA	ATG	GAG	TGC	ATG	AAG	AAG	AAA	GAA	CTC	GTC	ACT	GTG	TTC	AGA	ATG	GGT	TCT	GAG		840
G	Q	Q	D	I	E	M	A	I	L	T	A	L	L	K	G	T	N	V	S		300
GGC	CAG	CAG	GAC	ATC	GAG	ATG	GCA	ATT	TTA	ACT	GCC	CTG	CTG	AAA	GGA	ACA	AAC	GTA	TCT		900
A	P	D	Q	L	S	L	A	L	A	W	N	R	V	D	I	A	R	S	Q		320
GCT	CCA	GAT	CAG	CTG	AGC	TTG	GCA	CTG	GCT	TGG	AAC	CGC	GTG	GAC	ATA	GCA	CGA	AGC	CAG		960
I	F	V	F	G	P	H	W	T	P	L	G	S	L	A	P	P	T	D	S		340
ATC	TTT	GTC	TTT	GGG	CCC	CAC	TGG	ACG	CCC	CTG	GGA	AGC	CTG	GCA	CCC	CCG	ACG	GAC	AGC		1020
K	A	T	E	K	E	K	K	P	P	M	A	T	T	K	G	G	R	G	K		360
AAA	GCC	ACG	GAG	AAG	GAG	AAG	AAG	CCA	CCC	ATG	GCC	ACC	ACC	AAG	GGA	GGA	AGA	GGA	AAA		1080
G	K	G	K	K	K	G	K	V	K	E	E	V	E	E	E	T	D	P	R		380
GGG	AAA	GGC	AAG	AAG	AAA	GGG	AAA	GTG	AAA	GAG	GAA	GTG	GAG	GAA	GAA	ACT	GAC	CCC	CGG		1140
K	I	E	L	L	N	W	V	N	A	L	E	Q	A	M	L	D	A	L	V		400

AAG ATA GAG CTG CTG AAC TGG GTG AAT GCT TTG GAG CAA GCG ATG CTA GAT GCT TTA GTC	1200
L D R V D F V K L L I E N G V N M Q H F	420
TTA GAT CGT GTC GAC TTT GTG AAG CTC CTG ATT GAA AAC GGA GTG AAC ATG CAA CAC TTT	1260
L T I P R L E E L Y N T R L G P P N T L	440
CTG ACC ATT CCG AGG CTG GAG GAG CTT TAT AAC ACA AGA CTG GGT CCA CCA AAC ACA CTT	1320
H L L V R D V K K S N L P P D Y H I S L	460
CAT CTG CTG GTG AGG GAT GTG AAA AAG AGC AAC CTT CCG CCT GAT TAC CAC ATC AGC CTC	1380
I D I G L V L E Y L M G G A Y R C N Y T	480
ATA GAC ATC GGG CTC GTG CTG GAG TAC CTC ATG GGA GGA GCC TAC CGC TGC AAC TAC ACT	1440
R K N F R T L Y N N L F G P K R P K A L	500
CGG AAA AAC TTT CGG ACC CTT TAC AAC AAC TTG TTT GGA CCA AAG AGG CCT AAA GCT CTT	1500
K L L G M E D D E P P A K G K K K K K K	520
AAA CTT CTG GGA ATG GAA GAT GAT GAG CCT CCA GCT AAA GGG AAG AAA AAA AAA AAA AAG	1560
K K E E E I D I D V D D P A V S R F Q Y	540
AAA AAG GAG GAA GAG ATC GAC ATT GAT GTG GAC GAC CCT GCC GTG AGT CGG TTC CAG TAT	1620
P F H E L M V W A V L M K R Q K M A V F	560
CCC TTC CAC GAG CTG ATG GTG TGG GCA GTG CTG ATG AAA CGC CAG AAA ATG GCA GTG TTC	1680
L W Q R G E E S M A K A L V A C K L Y K	580
CTC TGG CAG CGA GGG GAA GAG AGC ATG GCC AAG GCC CTG GTG GCC TGC AAG CTC TAC AAG	1740
A M A H E S S E S D L V D D I S Q D L D	600
GCC ATG GCC CAC GAG TCC TCC GAG AGT GAT CTG GTG GAT GAC ATC TCC CAG GAC TTG GAT	1800
N N S K D F G Q L A L E L L D Q S Y K H	620
AAC AAT TCC AAA GAC TTC GGC CAG CTT GCT TTG GAG TTA TTA GAC CAG TCC TAT AAG CAT	1860
D E Q I A M K L L T Y E L K N W S N S T	640
GAC GAG CAG ATC GCT ATG AAA CTC CTG ACC TAC GAG CTG AAA AAC TGG AGC AAC TCG ACC	1920
C L K L A V A A K H R D F I A H T C S Q	660
TGC CTC AAA CTG GCC GTG GCA GCC AAA CAC CGG GAC TTC ATT GCT CAC ACC TGC AGC CAG	1980
M L L T D M W M G R L R M R K N P G L K	680
ATG CTG CTG ACC GAT ATG TGG ATG GGA AGA CTG CGG ATG CGG AAG AAC CCC GGC CTG AAG	2040
V I M G I L L P P T I L F L E F R T Y D	700
GTT ATC ATG GGG ATT CTT CTA CCC CCC ACC ATC TTG TTT TTG GAA TTT CGC ACA TAT GAT	2100
D F S Y Q T S K E N E D G K E K E E E N	720
GAT TTC TCG TAT CAA ACA TCC AAG GAA AAC GAG GAT GGC AAA GAA AAA GAA GAG GAA AAT	2160
T D A N A D A G S R K G D E E N E H K K	740
ACG GAT GCA AAT GCA GAT GCT GGC TCA AGA AAG GGG GAT GAG GAG AAC GAG CAT AAA AAA	2220
Q R I I P I G T K I C K F Y N A P I V K	760
CAA AGA ATT ATC CCC ATC GGA ACA AAA ATC TGT AAA TTC TAT AAC GCG CCC ATT GTC AAG	2280
F W F Y T I S Y L G Y L L L F N Y V I L	780
TTC TGG TTT TAC ACA ATA TCA TAC TTG GGC TAC CTG CTG CTG TTT AAC TAC GTC ATC CTG	2340
V R M D G W P S L Q E W I V I S Y I V S	800
GTG CGG ATG GAT GGC TGG CCG TCC CTC CAG GAG TGG ATC GTC ATC TCC TAC ATC GTG AGC	2400
L A L E K I R E I L M S E P G K L S Q K	820
CTG GCG TTA GAG AAG ATA CGA GAG ATC CTC ATG TCA GAA CCA GGC AAA CTC AGC CAG AAA	2460

FIGURE 5 cont.

I K V W L Q E Y W N I T D L V A I S T F	840
ATC AAA GTT TGG CTT CAG GAG TAC TGG AAC ATC ACA GAT CTC GTG GCC ATT TCC ACA TTC	2520
M I G A M A T R S V M M I G K M M I D M	860
ATG ATT GGA GCA ATG GCC ACG AGA TCT GTG ATG ATG ATT GGA AAG ATG ATG ATC GAC ATG	2580
L Y F V V I M L V V L M S F G V A R Q A	880
CTG TAC TTT GTG GTC ATC ATG CTG GTC GTG CTC ATG AGT TTC GGA GTA GCC CGT CAA GCC	2640
I L H P E E K P S W K L A R N I F Y M P	900
ATT CTG CAT CCA GAG GAG AAG CCC TCT TGG AAA CTG GCC CGA AAC ATC TTC TAC ATG CCC	2700
Y W M I Y G E V F A D Q I D L Y A M E I	920
TAC TGG ATG ATC TAT GGA GAG GTG TTT GCA GAC CAG ATA GAC CTC TAC GCC ATG GAA ATT	2760
N P P C G E N L Y D E E G K R L P P C I	940
AAT CCT CCT TGT GGT GAG AAC CTA TAT GAT GAG GAG GGC AAG CGG CTT CCT CCC TGT ATC	2820
P G A W L T P A L M A C Y L L V A N I L	960
CCC GGC GCC TGG CTC ACT CCA GCA CTC ATG GCG TGC TAT CTA CTG GTC GCC AAC ATC CTG	2880
L V N L L I A V F N N T F F E V K S I S	980
CTG GTG AAC CTG CTG ATT GCT GTG TTC AAC AAT ACC TTC TTT GAA GTA AAA TCA ATA TCC	2940
N Q V W K F Q R Y Q L I M T F H D R P V	1000
AAC CAG GTG TGG AAG TTC CAG CGA TAT CAG CTG ATT ATG ACA TTT CAT GAC AGG CCA GTC	3000
L P P P M I I L S H I Y I I I M R L S G	1020
CTG CCC CCA CCG ATG ATC ATT TTA AGC CAC ATC TAC ATC ATC ATT ATG CGT CTC AGC GGC	3060
R C R K K R E G D Q E E R D R G L K L F	1040
CGC TGC AGG AAA AAG AGA GAA GGG GAC CAA GAG GAA CGG GAT CGT GGA TTG AAG CTC TTC	3120
L S D E E L K R L H E F E E Q C V Q E H	1060
CTT AGC GAC GAG GAG CTA AAG AGG CTG CAT GAG TTC GAG GAG CAG TGC GTG CAG GAG CAC	3180
F R E K E D E Q Q S S S D E R I R V T S	1080
TTC CGG GAG AAG GAG GAT GAG CAG CAG TCG TCC AGC GAC GAG CGC ATC CGG GTC ACT TCT	3240
E R V E N M S M R L E E I N E R E T F M	1100
GAA AGA GTT GAA AAT ATG TCA ATG AGG TTG GAA GAA ATC AAT GAA AGA GAA ACT TTT ATG	3300
K T S L Q T V D L R L A Q L E E L S N R	1120
AAA ACT TCC CTG CAG ACT GTT GAC CTT CGA CTT GCT CAG CTA GAA GAA TTA TCT AAC AGA	3360
M V N A L E N L A G I D R S D L I Q A R	1140
ATG GTG AAT GCT CTT GAA AAT CTT GCG GGA ATC GAC AGG TCT GAC CTG ATC CAG GCA CGG	3420
S R A S S E C E A T Y L L R Q S S I N S	1160
TCC CGG GCT TCT TCT GAA TGT GAG GCA ACG TAT CTT CTC CGG CAA AGC AGC ATC AAT AGC	3480
A D G Y S L Y R Y H F N G E E L L F E D	1180
GCT GAT GGC TAC AGC TTG TAT CGA TAT CAT TTT AAC GGA GAA GAG TTA TTA TTT GAG GAT	3540
T S L S T S P G T G V R K K T C S F R I	1200
ACA TCT CTC TCC ACG TCA CCA GGG ACA GGA GTC AGG AAA AAA ACC TGT TCC TTC CGT ATA	3600
K E E K D V K T H L V P E C Q N S L H L	1220
AAG GAA GAG AAG GAC GTG AAA ACG CAC CTA GTC CCA GAA TGT CAG AAC AGT CTT CAC CTT	3660
S L G T S T S A T P D G S H L A V D D L	1240
TCA CTG GGC ACA AGC ACA TCA GCA ACC CCA GAT GGC AGT CAC CTT GCA GTA GAT GAC TTA	3720

FIGURE 5 cont.

K N A E E S K L G P D I G I S K E D D E	1260
AAG AAC GCT GAA GAG TCA AAA TTA GGT CCA GAT ATT GGG ATT TCA AAG GAA GAT GAT GAA	3780
R Q T D S K K E E T I S P S L N K T D V	1280
AGA CAG ACA GAC TCT AAA AAA GAA GAA ACT ATT TCC CCA AGT TTA AAT AAA ACA GAT GTG	3840
I H G Q D K S D V Q N T Q L T V E T T N	1300
ATA CAT GGA CAG GAC AAA TCA GAT GTT CAA AAC ACT CAG CTA ACA GTG GAA ACG ACA AAT	3900
I E G T I S Y P L E E T K I T R Y F P D	1320
ATA GAA GGC ACT ATT TCC TAT CCC CTG GAA GAA ACC AAA ATT ACA CGC TAT TTC CCC GAT	3960
E T I N A C K T M K S R S F V Y S R G R	1340
GAA ACG ATC AAT GCT TGT AAA ACA ATG AAG TCC AGA AGC TTC GTC TAT TCC CGG GGA AGA	4020
K L V G G V N Q D V E Y S S I T D Q Q L	1360
AAG CTG GTC GGT GGG GTT AAC CAG GAT GTA GAG TAC AGT TCA ATC ACG GAC CAG CAA TTG	4080
T T E W Q C Q V Q K I T R S H S T D I P	1380
ACG ACG GAA TGG CAA TGC CAA GTT CAA AAG ATC ACG CGC TCT CAT AGC ACA GAT ATT CCT	4140
Y I V S E A A V Q A E Q K E Q F A D M Q	1400
TAC ATT GTG TCG GAA GCT GCA GTG CAA GCT GAG CAA AAA GAG CAG TTT GCA GAT ATG CAA	4200
D E H H V A E A I P R I P R L S L T I T	1420
GAT GAA CAC CAT GTC GCT GAA GCA ATT CCT CGA ATC CCT CGC TTG TCC CTA ACC ATT ACT	4260
D R N G M E N L L S V K P D Q T L G F P	1440
GAC AGA AAT GGG ATG GAA AAC TTA CTG TCT GTG AAG CCA GAT CAA ACT TTG GGA TTC CCA	4320
S L R S K S L H G H P R N V K S I Q G K	1460
TCT CTC AGG TCA AAA AGT TTA CAT GGA CAT CCT AGG AAT GTG AAA TCC ATT CAG GGA AAG	4380
L D R S G H A S S V S S L V I V S G M T	1480
TTA GAC AGA TCT GGA CAT GCC AGT AGT GTA AGC AGC TTA GTA ATT GTG TCT GGA ATG ACA	4440
A E E K K V K K E K A S T E T E C (SEQ ID NO:7)	1497
GCA GAA GAA AAA AAG GTT AAG AAA GAG AAA GCT TCC ACA GAA ACT GAA TGC TAG TCTGTT	4991

TTGTTTCTTTAATTTTTTTTTTAAACAGTCAGAACCACCTAATGGGTGTCATCTTGCCATCCTAAACATCCATCCAAT

TTCTTAAAAACATTTTCCCT (SEQ ID NO:6)

FIGURE 6

ACTCATTATAGGGNTCGAGCGGCCCGCCGGGCGAGGTTTGTAGCTGTGCCCTCTCCATTCCACTGCTGTGGCAGGGTCAGA
 AATCTTGGATAGAGAAAACCTTTTGCAAACGGGAATGTATCTTTGTAAATCCTAGCACGAAAGACTCTAACAGGTGTG
 CTGTGGCCAGTTTACCAACCAGCATATCCCCCTCTGCCAAGTGCAACACCCAGCAAAATGAAGAGGAAAGCAAACAG
 GTGGAGACTCAGCCTGAGAAATGGTCTGTGCAAGCACACCCAGAGCTACCCAACAGATTCTATGGAGTTCTTGAATT
 CCAGGGTGGCGATATTCCAATAAGCC

M	Y	I	R	V	S	Y	D	T	K	P	D	S	L	L	H	L	M	V	K	20
ATG	TAT	ATC	CGT	GTA	TCC	TAT	GAC	ACC	AAG	CCA	GAC	TCA	CTG	CTC	CAT	CTC	ATG	GTG	AAA	60
D	W	Q	L	E	L	P	K	L	L	I	S	V	H	G	G	L	Q	N	F	40
GAT	TGG	CAG	CTG	GAA	CTC	CCC	AAG	CTC	TTA	ATA	TCT	GTG	CAT	GGA	GGC	CTC	CAG	AAC	TTT	120
E	M	Q	P	K	L	K	Q	V	F	G	K	G	L	I	K	A	A	M	T	60
GAG	ATG	CAG	CCC	AAG	CTG	AAA	CAA	GTC	TTT	GGG	AAA	GGC	CTG	ATC	AAG	GCT	GCT	ATG	ACC	180
T	G	A	W	I	F	T	G	G	V	S	T	G	V	I	S	H	V	G	D	80
ACC	GGG	GCC	TGG	ATC	TTC	ACC	GGG	GGT	GTC	AGC	ACA	GGT	GTT	ATC	AGC	CAC	GTA	GGG	GAT	240
A	L	K	D	H	S	S	K	S	R	G	R	V	C	A	I	G	I	A	P	100
GCC	TTG	AAA	GAC	CAC	TCC	TCC	AAG	TCC	AGA	GGC	CGG	GTT	TGT	GCT	ATA	GGA	ATT	GCT	CCA	300
W	G	I	V	E	N	K	E	D	L	V	G	K	D	V	T	R	V	Y	Q	120
TGG	GGC	ATC	GTG	GAG	AAT	AAG	GAA	GAC	CTG	GTT	GGA	AAG	GAT	GTA	ACA	AGA	GTG	TAC	CAG	360
T	M	S	N	P	L	S	K	L	S	V	L	N	N	S	H	T	H	F	I	140
ACC	ATG	TCC	AAC	CCT	CTA	AGT	AAG	CTC	TCT	GTG	CTC	AAC	AAC	TCC	CAC	ACC	CAC	TTC	ATC	420
L	A	D	N	G	T	L	G	K	Y	G	A	E	V	K	L	R	R	L	L	160
CTG	GCT	GAC	AAT	GGC	ACC	CTG	GGC	AAG	TAT	GGC	GCC	GAG	GTG	AAG	CTG	CGA	AGG	CTG	CTG	480
E	K	H	I	S	L	Q	K	I	N	T	R	L	G	Q	G	V	P	L	V	180
GAA	AAG	CAC	ATC	TCC	CTC	CAG	AAG	ATC	AAC	ACA	AGA	CTG	GGG	CAG	GGC	GTG	CCC	CTC	GTG	540
G	L	V	V	E	G	G	P	N	V	V	S	I	V	L	E	Y	L	Q	E	200
GGT	CTC	GTG	GTG	GAG	GGG	GGC	CCT	AAC	GTG	GTG	TCC	ATC	GTC	TTG	GAA	TAC	CTG	CAA	GAA	600
E	P	P	I	P	V	V	I	C	D	G	S	G	R	A	S	D	I	L	S	220
GAG	CCT	CCC	ATC	CCT	GTG	GTG	ATT	TGT	GAT	GGC	AGC	GGA	CGT	GCC	TGC	GAC	ATC	CTG	TCC	660
F	A	H	K	Y	C	E	E	G	G	I	I	N	E	S	L	R	E	Q	L	240
TTT	GCG	CAC	AAG	TAC	TGT	GAA	GAA	GGC	GGA	ATA	ATA	AAT	GAG	TCC	CTC	AGG	GAG	CAG	CTT	720
L	V	T	I	Q	K	T	F	N	Y	N	K	A	Q	S	H	Q	L	F	A	260
CTA	GTT	ACC	ATT	CAG	AAA	ACA	TTT	AAT	TAT	AAT	AAG	GCA	CAA	TCA	CAT	CAG	CTG	TTT	GCA	780
I	I	M	E	C	M	K	K	K	E	L	V	T	V	F	R	M	G	S	E	280
ATT	ATA	ATG	GAG	TGC	ATG	AAG	AAG	AAA	GAA	CTC	GTC	ACT	GTG	TTC	AGA	ATG	GGT	TCT	GAG	840
G	Q	Q	D	I	E	M	A	I	L	T	A	L	L	K	G	T	N	V	S	300
GGC	CAG	CAG	GAC	ATC	GAG	ATG	GCA	ATT	TTA	ACT	GCC	CTG	CTG	AAA	GGA	ACA	AAC	GTA	TCT	900
A	P	D	Q	L	S	L	A	L	A	W	N	R	V	D	I	A	R	S	Q	320
GCT	CCA	GAT	CAG	CTG	AGC	TTG	GCA	CTG	GCT	TGG	AAC	CGC	GTG	GAC	ATA	GCA	CGA	AGC	CAG	960
I	F	V	F	G	P	H	W	T	P	L	G	S	L	A	P	P	T	D	S	340
ATC	TTT	GTC	TTT	GGG	CCC	CAC	TGG	ACG	CCC	CTG	GGA	AGC	CTG	GCA	CCC	CCG	ACG	GAC	AGC	1020
K	A	T	E	K	E	K	K	P	P	M	A	T	T	K	G	G	R	G	K	360
AAA	GCC	ACG	GAG	AAG	GAG	AAG	AAG	CCA	CCC	ATG	GCC	ACC	ACC	AAG	GGA	GGA	AGA	GGA	AAA	1080
G	K	G	K	K	K	G	K	V	K	E	E	V	E	E	E	T	D	P	R	380
GGG	AAA	GGC	AAG	AAG	AAA	GGG	AAA	GTG	AAA	GAG	GAA	GTG	GAG	GAA	GAA	ACT	GAC	CCC	CGG	1140
K	I	E	L	L	N	W	V	N	A	L	E	Q	A	M	L	D	A	L	V	400

FIGURE 6 cont.

AAG ATA GAG CTG CTG AAC TGG GTG AAT GCT TTG GAG CAA GCG ATG CTA GAT GCT TTA GTC	1200
L D R V D F V K L L I E N G V N M Q H F	420
TTA GAT CGT GTC GAC TTT GTG AAG CTC CTG ATT GAA AAC GGA GTG AAC ATG CAA CAC TTT	1260
L T I P R L E E L Y N T R L G P P N T L	440
CTG ACC ATT CCG AGG CTG GAG GAG CTT TAT AAC ACA AGA CTG GGT CCA CCA AAC ACA CTT	1320
H L L V R D V K K S N L P P D Y H I S L	460
CAT CTG CTG GTG AGG GAT GTG AAA AAG AGC AAC CTT CCG CCT GAT TAC CAC ATC AGC CTC	1380
I D I G L V L E Y L M G G A Y R C N Y T	480
ATA GAC ATC GGG CTC GTG CTG GAG TAC CTC ATG GGA GGA GCC TAC CGC TGC AAC TAC ACT	1440
R K N F R T L Y N N L F G P K R P K A L	500
CGG AAA AAC TTT CGG ACC CTT TAC AAC AAC TTG TTT GGA CCA AAG AGG CCT AAA GCT CTT	1500
K L L G M E D D E P P A K G K K K K K K	520
AAA CTT CTG GGA ATG GAA GAT GAT GAG CCT CCA GCT AAA GGG AAG AAA AAA AAA AAA AAG	1560
K K E E E I D I D V D D P A V S R F Q Y	540
AAA AAG GAG GAA GAG ATC GAC ATT GAT GTG GAC GAC CCT GCC GTG AGT CGG TTC CAG TAT	1620
P F H E L M V W A V L M K R Q K M A V F	560
CCC TTC CAC GAG CTG ATG GTG TGG GCA GTG CTG ATG AAA CGC CAG AAA ATG GCA GTG TTC	1680
L W Q R G E E S M A K A L V A C K L Y K	580
CTC TGG CAG CGA GGG GAA GAG AGC ATG GCC AAG GCC CTG GTG GCC TGC AAG CTC TAC AAG	1740
A M A H E S S E S D L V D D I S Q D L D	600
GCC ATG GCC CAC GAG TCC TCC GAG AGT GAT CTG GTG GAT GAC ATC TCC CAG GAC TTG GAT	1800
N N S K D F G Q L A L E L L D Q S Y K H	620
AAC AAT TCC AAA GAC TTC GGC CAG CTT GCT TTG GAG TTA TTA GAC CAG TCC TAT AAG CAT	1860
D E Q I A M K L L T Y E L K N W S N S T	640
GAC GAG CAG ATC GCT ATG AAA CTC CTG ACC TAC GAG CTG AAA AAC TGG AGC AAC TCG ACC	1920
C L K L A V A A K H R D F I A H T C S Q	660
TGC CTC AAA CTG GCC GTG GCA GCC AAA CAC CGG GAC TTC ATT GCT CAC ACC TGC AGC CAG	1980
M L L T D M W M G R L R M R K N P G L K	680
ATG CTG CTG ACC GAT ATG TGG ATG GGA AGA CTG CGG ATG CGG AAG AAC CCC GGC CTG AAG	2040
V I M G I L L P P T I L F L E F R T Y D	700
GTT ATC ATG GGG ATT CTT CTA CCC CCC ACC ATC TTG TTT TTG GAA TTT CGC ACA TAT GAT	2100
D F S Y Q T S K E N E D G K E K E E E N	720
GAT TTC TCG TAT CAA ACA TCC AAG GAA AAC GAG GAT GGC AAA GAA AAA GAA GAG GAA AAT	2160
T D A N A D A G S R K G D E E N E H K K	740
ACG GAT GCA AAT GCA GAT GCT GGC TCA AGA AAG GGG GAT GAG GAG AAC GAG CAT AAA AAA	2220
Q R I I P I G T K I C K F Y N A P I V K	760
CAA AGA ATT ATC CCC ATC GGA ACA AAA ATC TGT AAA TTC TAT AAC GCG CCC ATT GTC AAG	2280
F W F Y T I S Y L G Y L L L F N Y V I L	780
TTC TGG TTT TAC ACA ATA TCA TAC TTG GGC TAC CTG CTG CTG TTT AAC TAC GTC ATC CTG	2340
V R M D G W P S L Q E W I V I S Y I V S	800
GTG CGG ATG GAT GGC TGG CCG TCC CTC CAG GAG TGG ATC GTC ATC TCC TAC ATC GTG AGC	2400
L A L E K I R E I L M S E P G K L S Q K	820
CTG GCG TTA GAG AAG ATA CGA GAG ATC CTC ATG TCA GAA CCA GGC AAA CTC AGC CAG AAA	2460

FIGURE 6 cont.

I K V W L Q E Y W N I T D L V A I S T F	840
ATC AAA GTT TGG CTT CAG GAG TAC TGG AAC ATC ACA GAT CTC GTG GCC ATT TCC ACA TTC	2520
M I G A I L R L Q N Q P Y M G Y G R V I	860
ATG ATT GGA GCA ATT CTT CGC CTA CAG AAC CAG CCC TAC ATG GGC TAT GGC CGG GTG ATC	2580
Y C V D I I F W Y I R V L D I F G V N K	880
TAC TGT GTG GAT ATC ATC TTC TGG TAC ATC CGT GTC CTG GAC ATC TTT GGT GTC AAC AAG	2640
Y L G P Y V M M I G K M M I D M L Y F V	900
TAT CTG GGG CCA TAC GTG ATG ATG ATT GGA AAG ATG ATG ATC GAC ATG CTG TAC TTT GTG	2700
V I M L V V L M S F G V A R Q A I L H P	920
GTC ATC ATG CTG GTC GTG CTC ATG AGT TTC GGA GTA GCC CGT CAA GCC ATT CTG CAT CCA	2760
E E K P S W K L A R N I F Y M P Y W M I	940
GAG GAG AAG CCC TCT TGG AAA CTG GCC CGA AAC ATC TTC TAC ATG CCC TAC TGG ATG ATC	2820
Y G E V F A D Q I D L Y A M E I N P P C	960
TAT GGA GAG GTG TTT GCA GAC CAG ATA GAC CTC TAC GCC ATG GAA ATT AAT CCT CCT TGT	2880
G E N L Y D E E G K R L P P C I P G A W	980
GGT GAG AAC CTA TAT GAT GAG GAG GGC AAG CGG CTT CCT CCC TGT ATC CCC GGC GCC TGG	2940
L T P A L M A C Y L L V A N I L L V N L	1000
CTC ACT CCA GCA CTC ATG GCG TGC TAT CTA CTG GTC GCC AAC ATC CTG CTG GTG AAC CTG	3000
L I A V F N N T F F E V K S I S N Q V W	1020
CTG ATT GCT GTG TTC AAC AAT ACT TTC TTT GAA GTA AAA TCA ATA TCC AAC CAG GTG TGG	3060
K F Q R Y Q L I M T F H D R P V L P P P	1040
AAG TTC CAG CGA TAT CAG CTG ATT ATG ACA TTT CAT GAC AGG CCA GTC CTG CCC CCA CCG	3120
M I I L S H I Y I I I M R L S G R C R K	1060
ATG ATC ATT TTA AGC CAC ATC TAC ATC ATC ATT ATG CGT CTC AGC GGC CGC TGC AGG AAA	3180
K R E G D Q E E R D R G L K L F L S D E	1080
AAG AGA GAA GGG GAC CAA GAG GAA CGG GAT CGT GGA TTG AAG CTC TTC CTT AGC GAC GAG	3240
E L K R L H E F E E Q C V Q E H F R E K	1100
GAG CTA AAG AGG CTG CAT GAG TTC GAG GAG CAG TGC GTG CAG GAG CAC TTC CGG GAG AAG	3300
E D E Q Q S S S D E R I R V T S E R V E	1120
GAG GAT GAG CAG CAG TCG TCC AGC GAC GAG CGC ATC CGG GTC ACT TCT GAA AGA GTT GAA	3360
N M S M R L E E I N E R E T F M K T S L	1140
AAT ATG TCA ATG AGG TTG GAA GAA ATC AAT GAA AGA GAA ACT TTT ATG AAA ACT TCC CTG	3420
Q T V D L R L A Q L E E L S N R M V N A	1160
CAG ACT GTT GAC CTT CGA CTT GCT CAG CTA GAA GAA TTA TCT AAC AGA ATG GTG AAT GCT	3480
L E N L A G I D R S D L I Q A R S R A S	1180
CTT GAA AAT CTT GCG GGA ATC GAC AGG TCT GAC CTG ATC CAG GCA CGG TCC CGG GCT TCT	3540
S E C E A T Y L L R Q S S I N S A D G Y	1200
TCT GAA TGT GAG GCA ACG TAT CTT CTC CGG CAA AGC AGC ATC AAT AGC GCT GAT GGC TAC	3600
S L Y R Y H F N G E E L L F E D T S L S	1220
AGC TTG TAT CGA TAT CAT TTT AAC GGA GAA GAG TTA TTA TTT GAG GAT ACA TCT CTC TCC	3660
T S P G T G V R K K T C S F R I K E E K	1240
ACG TCA CCA GGG ACA GGA GTC AGG AAA AAA ACC TGT TCC TTC CGT ATA AAG GAA GAG AAG	3720

FIGURE 6 cont.

D V K T H L V P E C Q N S L H L S L G T	1260
GAC GTG AAA ACG CAC CTA GTC CCA GAA TGT CAG AAC AGT CTT CAC CTT TCA CTG GGC ACA	3780
S T S A T P D G S H L A V D D L K N A E	1280
AGC ACA TCA GCA ACC CCA GAT GGC AGT CAC CTT GCA GTA GAT GAC TTA AAG AAC GCT GAA	3840
E S K L G P D I G I S K E D D E R Q T D	1300
GAG TCA AAA TTA GGT CCA GAT ATT GGG ATT TCA AAG GAA GAT GAT GAA AGA CAG ACA GAC	3900
S K K E E T I S P S L N K T D V I H G Q	1320
TCT AAA AAA GAA GAA ACT ATT TCC CCA AGT TTA AAT AAA ACA GAT GTG ATA CAT GGA CAG	3960
D K S D V Q N T Q L T V E T T N I E G T	1340
GAC AAA TCA GAT GTT CAA AAC ACT CAG CTA ACA GTG GAA ACG ACA AAT ATA GAA GGC ACT	4020
I S Y P L E E T K I T R Y F P D E T I N	1360
ATT TCC TAT CCC CTG GAA GAA ACC AAA ATT ACA CGC TAT TTC CCC GAT GAA ACG ATC AAT	4080
A C K T M K S R S F V Y S R G R K L V G	1380
GCT TGT AAA ACA ATG AAG TCC AGA AGC TTC GTC TAT TCC CGG GGA AGA AAG CTG GTC GGT	4140
G V N Q D V E Y S S I T D Q Q L T T E W	1400
GGG GTT AAC CAG GAT GTA GAG TAC AGT TCA ATC ACG GAC CAG CAA TTG ACG ACG GAA TGG	4200
Q C Q V Q K I T R S H S T D I P Y I V S	1420
CAA TGC CAA GTT CAA AAG ATC ACG CGC TCT CAT AGC ACA GAT ATT CCT TAC ATT GTG TCG	4260
E A A V Q A E Q K E Q F A D M Q D E H H	1440
GAA GCT GCA GTG CAA GCT GAG CAA AAA GAG CAG TTT GCA GAT ATG CAA GAT GAA CAC CAT	4320
V A E A I P R I P R L S L T I T D R N G	1460
GTC GCT GAA GCA ATT CCT CGA ATC CCT CGC TTG TCC CTA ACC ATT ACT GAC AGA AAT GGG	4380
M E N L L S V K P D Q T L G F P S L R S	1480
ATG GAA AAC TTA CTG TCT GTG AAG CCA GAT CAA ACT TTG GGA TTC CCA TCT CTC AGG TCA	4440
K S L H G H P R N V K S I Q G K L D R S	1500
AAA AGT TTA CAT GGA CAT CCT AGG AAT GTG AAA TCC ATT CAG GGA AAG TTA GAC AGA TCT	4500
G H A S S V S S L V I V S G M T A E E K	1520
GGA CAT GCC AGT AGT GTA AGC AGC TTA GTA ATT GTG TCT GGA ATG ACA GCA GAA GAA AAA	4560
K V K K E K A S T E T E C * (SEQ ID NO:9)	1533
AAG GTT AAG AAA GAG AAA GCT TCC ACA GAA ACT GAA TGC TAG TCTGTTTTGTTTCTTTAATTTTT	4599
TTTTTTAACAGTCAGAAACCCACTAATGGGTGTCATCTTGCCCATCCTAAACACATMTCCAATTTCTCTAAAAACATTT	
TCCCT (SEQ ID NO:8)	

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled COMPOSITIONS AND METHODS FOR THE DIAGNOSIS, PREVENTION AND TREATMENT OF TUMOR PROGRESSION, the specification of which

☒ is attached hereto.

☐ was filed on _____ as Application Serial No. _____ and was amended on _____.

☐ was described and claimed in PCT International Application No. _____ filed on _____ and as amended under PCT Article 19 on _____.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose all information I know to be material to patentability in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose all information I know to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56(a) which became available between the filing date of the prior application and the national or PCT international filing date of this application:

U.S. SERIAL NO.	FILING DATE	STATUS
08/412,431	March 29, 1995	<input checked="" type="checkbox"/> Pending <input type="checkbox"/> Issued <input type="checkbox"/> Abandoned

I hereby appoint the following attorneys and/or agents to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: J. Peter Fasse, Reg. No. 32,983, William E. Booth, Reg. No. 28,933; Barry E. Bretschneider, Reg. No. 28,055; John W. Freeman, Reg. No. 29,066; Timothy A. French, Reg. No. 30,175; Alan H. Gordon, Reg. No. 26,168; John F. Land, Reg. No. 29,554; John B. Pegram, Reg. No. 25,198; Rene D. Tegtmeyer, Reg. No. 33,567; Hans R. Troesch, Reg. No. 36,950; Dorothy P. Whelan, Reg. No. 33,814; Charles C. Winchester, Reg. No. 21,040.

Address all telephone calls to J. Peter Fasse at telephone number 617/542-5070.

Address all correspondence to J. Peter Fasse, Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patents issued thereon.

COMBINED DECLARATION AND POWER OF ATTORNEY CONTINUED

Full Name of Inventor: Andrew W. Shyjan

Inventor's Signature: _____ Date: _____

Residence Address: 58 Colby Way, Nahant, Massachusetts 01908

Citizen of: U.S.A.

Post Office Address: 58 Colby Way, Nahant, Massachusetts 01908

266250 24429880

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled COMPOSITIONS AND METHODS FOR THE DIAGNOSIS, PREVENTION AND TREATMENT OF TUMOR PROGRESSION, the specification of which

■ was filed on March 29, 1996 as Application Serial No. 08/623,679 and was amended on _____.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose all information I know to be material to patentability in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose all information I know to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56(a) which became available between the filing date of the prior application and the national or PCT international filing date of this application:

U.S. SERIAL NO.	FILING DATE	STATUS
<u>08/412,431</u>	<u>March 29, 1995</u>	■ Pending □ Issued □ Abandoned

I hereby appoint the following attorneys and/or agents to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: J. Peter Fasse, Reg. No. 32,983, Mark F. Boshar, Reg. No. 35,456, William E. Booth, Reg. No. 28,933; Barry E. Bretschneider, Reg. No. 28,055; John W. Freeman, Reg. No. 29,066; Timothy A. French, Reg. No. 30,175; Alan H. Gordon, Reg. No. 26,168; John F. Land, Reg. No. 29,554; John B. Pegram, Reg. No. 25,198; Rene D. Tegtmeyer, Reg. No. 33,567; Hans R. Troesch, Reg. No. 36,950; Dorothy P. Whelan, Reg. No. 33,814; Charles C. Winchester, Reg. No. 21,040.

Address all telephone calls to J. Peter Fasse at telephone number 617/542-5070.

Address all correspondence to J. Peter Fasse, Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patents issued thereon.

COMBINED DECLARATION AND POWER OF ATTORNEY CONTINUED

Full Name of Inventor: Andrew W. Shyjan

Inventor's Signature: Andrew W. Shyjan Date: 5/9/90

Residence Address: 58 Colby Way, Nahant, Massachusetts 01908

Citizen of: U.S.A.

Post Office Address: 58 Colby Way, Nahant, Massachusetts 01908

003344 003344 003344